

Preparation and Evaluation of Oral Extended Release Tablet of Acelophenac for Better Efficacy

Shailendra Sriwastava¹ and Dr. Sangamesh B. Puranik²

Research Scholar, Department of Pharmacy¹

Professor, Department of Pharmacy²

OPJS University, Churu, Rajasthan, India

Abstract: There is need of extended release matrix tablet because of the reason that there is some short high life time and maximum therapeutic efficacy within the body so here we planed to prepare the acelophenac as an active ingradient with polymer and some necessary ingredients for the preparation of garnual and then after tablet preparation we find that there is good behaviour shown by the formulation and the best result is evaluated further for the study

Keywords: Extended release matrix

REFERENCES

- [1]. Yum, S. I., Schoenhard, G., Tipton, A. J., Gibson, J. W., Middleton, J. C., Fu, R., &Zamloot, M. S. (2016). *U.S. Patent No. 9,517,271*. Washington, DC: U.S. Patent and Trademark Office.
- [2]. Omprakash, B., Ajay, S., Santosh, G., & Amin, P. (2012). Formulation development of venlafaxine hydrochloride extended release tablet and invitro characterizations. *International Journal of PharmTech Research*, 4(4), 1777-1784.
- [3]. Khan, G. M. (2001). Controlled release oral dosage forms: Some recent advances in matrix type drug delivery systems. *The sciences*, 1(5), 350-354.
- [4]. Higuchi, T. (1963). Mechanism of sustained-action medication. Theoretical analysis of rate of release of solid drugs dispersed in solid matrices. *Journal of pharmaceutical sciences*, 52(12), 1145-1149.
- [5]. Davis, S. S., Hardy, J. G., &Fara, J. W. (1986). Transit of pharmaceutical dosage forms through the small intestine. *Gut*, 27(8), 886-892.
- [6]. Siepman, J., &Peppas, N. A. (2001). Modeling of drug release from delivery systems based on hydroxypropyl methylcellulose (HPMC). *Advanced drug delivery reviews*, 48(2-3), 139-157.
- [7]. Hirani, J. J., Rathod, D. A., &Vadalia, K. R. (2009). Orally disintegrating tablets: a review. *Tropical journal of pharmaceutical research*, 8(2).
- [8]. Barloy, L., Lallier, J. P., Battioni, P., Mansuy, D., Piffard, Y., Tournoux, M., ...& Jones, W. (1993). Manganese Porphyrins Adsorbed or Intercatalated in Different Mineral Matrices: Preparation and Compared Properties as Catalysts for Alkene and Alkane Oxidation. *ChemInform*, 24(11).
- [9]. Liu, Y., Layrolle, P., de Bruijn, J., van Blitterswijk, C., & de Groot, K. (2001). Biomimetic coprecipitation of calcium phosphate and bovine serum albumin on titanium alloy. *Journal of Biomedical Materials Research Part A*, 57(3), 327-335.
- [10]. Kaneko, K., & Ishii, C. (1992). Superhigh surface area determination of microporous solids. *Colloids and surfaces*, 67, 203-212.
- [11]. Fiedler, U., &Růžička, J. (1973). Selectrode—the universal ion-selective electrode: Part VII. A valinomycin-based potassium electrode with nonporous polymer membrane and solid-state inner reference system. *Analyticachimicaacta*, 67(1), 179-193.
- [12]. Siepman, J., &Peppas, N. A. (2000). Hydrophilic matrices for controlled drug delivery: an improved mathematical model to predict the resulting drug release kinetics (the “sequential layer” model). *Pharmaceutical Research*, 17(10), 1290-1298.

- [13]. Ju, R. T., Nixon, P. R., Patel, M. V., & Tong, D. M. (1995). Drug release from hydrophilic matrices. 2. A mathematical model based on the polymer disentanglement concentration and the diffusion layer. *Journal of pharmaceutical sciences*, 84(12), 1464-1477.
- [14]. Conti, S., Maggi, L., Segale, L., Machiste, E. O., Conte, U., Grenier, P., & Vergnault, G. (2007). Matrices containing NaCMC and HPMC: 2. Swelling and release mechanism study. *International Journal of pharmaceutics*, 333(1-2), 143-151.