

International Journal of Advanced Research in Science, Communication and Technology (IJARSCT)

International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal

Volume 4, Issue 3, March 2024

A Class of Multivalent Functions Associated with. rth Differential Operator.

¹Jadhav. S. S and ²Khairnar S. L

¹Sundarrao More Arts, Commerce, and Science (Sr.) College, Poladpur ²Changu Kana Thakur, Arts, Commerce and Science College, Panvel

Abstract: This paper is concern with multivalent functions. We derived the new r^{th} differential operator. Associated with this operator new class of q-valent functions is studied. Sufficient condition for this class has been obtained.

Keywords: q-valent, multivalent, regular, operator

REFERENCES

[1]Ahuja O.P., (1985), Integral operator of certain univalent functions, Int. J.Math.Sci, 8,653-662.

[2] Albehba M., Darus M., (2015), Subclass of meromorphic multivalent functions, Acta. Univ.Math.Inf. 43,157-167.

[3] Alexander J.W., (1915), Functions which maps interior of unit circle upon simple region, Ann. Math, 17, 12-22.

[4] Arif M., Ahmad K. J. Liu, Sokol J., (2019), New classes of analytic functions associated with salagean operator, J.of Funct.Spaces, Hindawi, ID, 6157394.

[5]Chaughule V.A., Naik U.H., (2020), Some properties of new subclasses of multivalent functions, IJRAR, 6(2)[6]Goodman A.W., (1979), An invitation to study of univalent and multivalent functions. Int . J. of Math and Math.Sci, 2, 163-186.

[7] Jack I.S., (1971), Functions starlike and convex of order t, J.LondonMath.Soc, 2(3),469-474.

[8] Juma A.R., (2008), Some problems connected with geometryof univalent and multivalent functions, PhD theis, University of Pune.

[9] Kalpan W, (1952), Close to convex schlicht functions, Mich.Math.J., 1, 169-185.

[10] Nunokawa M., (1987), Onth theory of multivalent functions, Tsukuba J.Math., 11,273-286.

