

DNA-Binding and Photocleavage Studies of 9-Phenyl Carbazole

Hanchate Pallavi¹ and Oddepally Rajender²

Pingle Government College for Women, Hanmakonda, India

Government Polytechnic, Station Ghanpur, India

rpic17@gmail.com¹ and oddepallyrajender@gmail.com²

Abstract: The DNA-binding mode of 9-Phenyl carbazole with CT-DNA was investigated by absorption spectroscopy, EB-DNA displacement, circular dichroism, thermal denaturation and viscosity measurements. Results indicated that these compounds intercalate into the base pairs of CT-DNA. The effect of ionic strength on the fluorescence property of the system indicated the presence of electrostatic interaction via phosphate backbone of DNA helix. The intrinsic binding constant values suggested that compound has DNA binding propensity. This compound promote the cleavage of plasmid pBR322. These results may be useful for the design of 9-Phenyl carbazole with desired binding characteristics and useful to better understand the DNA binding mode of heterocyclic compound.

Keywords: DNA-binding.

REFERENCES

- [1]. Y. Cao and X. W. He, "Studies of interaction between safranine T and double helix DNA by spectral methods," *Spectrochim. Acta A.*, vol **54**, pp. 883-892, 1998.
- [2]. P. B. Dervan, "Molecular Recognition of DNA by Small Molecules," *Bioorg. Med. Chem.*, vol **9**, pp 2215-2235, July 2001.
- [3]. L. H. Hurley, "DNA and associated processes as targets for cancer therapy," *Nat. Rev. Cancer*, vol **2**, pp 188-200, 2002.
- [4]. B. Gold, "Effect of cationic charge localization on DNA structure," *Biopolymers*, vol **65**, pp 173-179, November 2002.
- [5]. M. F. Brana, M. Cacho, A. Gradillas, B. de Pascual-Teresa, and A. Ramos, "Intercalators as anticancer drugs," *Curr. Pharm. Des.*, vol **7**, pp 1745-1780, December 2001.
- [6]. S. Neidle, "DNA minor-groove recognition by small molecules," *Nat. Prod. Rep.*, vol **18**, pp 291-309, June 2001.
- [7]. W. B. Tan, A. Bhamhani, M. R. Duff, A. Rodger, and C. V. Kumar, "Spectroscopic identification of binding modes of anthracene probes and DNA sequence recognition," *Photochem. Photobiol.*, vol **82**, pp 20-30, January 2006.
- [8]. J. J. Chen, J. C. Drach, L. B. Townsend, "Convergent Synthesis of polyhalogenated quinololines C-nucleosides as potential antiviral agents," *J. Org. Chem.*, vol **68**, pp 4170-4178, May 2003.
- [9]. X. Song, B. S. Vig, P. L. Lorenzi, J. C. Drach, L. B. Townsend, G. L. Amidon, "Amino acid ester prodrugs of the antiviral agent 2-bromo-5,6-dichloro-1-(beta-Dribofuranosyl) benzimidazole as potential substrates of hPEPT1 transporter," *J. Med. Chem.*, vol **48**, pp 1274-1277, February 2005.
- [10]. J. D. Venable, H. Cai, W. Chai, C. A. Dvorak, C. A. Grice, J. A. Jablonowski, C. R. Shah, A. K. Kwok, K. S. Ly, B. Pio, J. Wei, P. J. Desai, W. Jiang, S. Nguyen, P. Ling, S. J. Wilson, P. J. Dunford, R. L. Thurmond, T. W. Lovenberg, L. Karlsson, N.I. Carruthers, and J. P. Edwards, "Preparation and Biological Evaluation of Indole, Benzimidazole, and Thienopyrrole Piperazine Carboxamides: Potent Human Histamine H4 Antagonists," *J. Med. Chem.*, vol **48**, pp 8289-8298, December 2005.

- [11]. M. E. Reichmann, S. A. Rice, C. A. Thomas, and P. Doty, "A further examination of the molecular weight and size of desoxypentose nucleic acid," *J. Am. Chem. Soc.*, vol **76**, pp 3047-3053, June 1954.
- [12]. W. Zhong, J. S. Yu, W. Huang, K. Ni, and Y. Liang, "Spectroscopic Studies of Interaction of Chlorobenzylidine with DNA," *Biopolymers*, vol **62**, pp 315-323, October 2001.
- [13]. J. B. Chaires, N. Dattagupta, and D. M. Crothers, "Studies on Interaction of Anthracycline Antibiotics and Deoxyribonucleic Acid: Equilibrium Binding Studies on Interaction of Daunomycin with Deoxyribonucleic Acid," *Biochemistry*, vol **21**, pp 3933-3940, August 1982.
- [14]. J. K. Barton, A. T. Danishefsky, and J. M Goldberg, "Tris(phenanthroline) Ru(II): Stereoselectivity in Binding to DNA," *J. Am. Chem. Soc.*, vol **106**, pp 2172-2176, April 1984.
- [15]. F. Gao, H. Chao, F. Zhou, Y. X. Yuan, B. Peng, and L. N. Ji, "DNA interactions of a functionalized ruthenium(II) mixed-polypyridyl complex [Ru(bpy)₂ppd]²⁺," *J. Inorg. Biochem.*, vol **100**, pp 1487-1494, September 2006.
- [16]. A. Wolfe, G. H. Shimer, and T. Meehan, "Polycyclic aromatic hydrocarbons physically intercalate into duplex regions of denatured DNA," *Biochemistry*, vol **26**, pp 6392-6396, October 1987.
- [17]. J. B. Lepecq, and C. Paoletti, "A Fluorescent Complex between Ethidium Bromide and Nucleic Acids Physical-Chemical Characterization," *J. Mol. Biol.*, vol **27**, pp 87-106, July 1967.
- [18]. F. Y. Wu, Y. L. Xiang, Y. M. Wu, and F. Y. Xie, "Study of interaction of a fluorescent probe with DNA," *J. Lumin.*, vol **129**, pp 1286-1291, November 2009.
- [19]. T. C. Michael, R. Marisol, and J. B. Allen, "Voltammetric studies of the interaction of metal chelates with DNA. 2. Tris-chelated complexes of cobalt(III) and iron(II) with 1,10-phenanthroline and 2,2'-bipyridine," *J. Am. Chem. Soc.*, vol **111**, pp 8901-8911, November 1989.
- [20]. M. Eriksson, M. Leijon, C. Hiort, B. Norden, A. Gradsland, "Binding of delta- and lambda-[Ru(phen)₃]²⁺ to [d(CGCG ATCGCG)]₂ Studied by NMR," *Biochemistry*, vol **33**, pp 5031-5040, May 1994.
- [21]. Y. Xiong, X. F. He, X. H. Zou, J. Z. Wu, X. M. Chen, L. N. Ji, R. H. Li, J. Y. Zhou, and K. B. Yu, "Interaction of polypyridyl ruthenium(II) complexes containing non-planar ligands with DNA," *J. Chem. Soc. Dalton. Trans.*, vol 1, pp 19-24, October 1999.
- [22]. A. Sitlani, E. C. Long, A.M. Pyle, and J. K. Barton, "DNA Photocleavage by Phenanthrenequinone Diimine Complexes of Rhodium (III): shape-selective recognition and reaction," *J. Am. Chem. Soc.*, vol 114, pp 2303-2312, March 1992.