

Chronic Kidney Disease Prediction

W. Priscilla Victoriya¹ M. Lincy Jacqueline²

Student, M.Sc Computer Science, Nirmala College for Women (Autonomous) Coimbatore, Tamilnadu¹

Assistant Professor, Nirmala College for Women (Autonomous) Coimbatore, Tamilnadu²

Abstract: *Early diagnosis and characterization are the important components in determine the treatment of chronic kidney disease CKD. CKD is an ailment which tends to damage the kidney ad affect their effective functioning of excreting waste and balancing body fluids. Some of the complications included are hypertension, anaemia (low blood count), mineral bone disorder, poor nutritional healthy acid base abnormalities ad neurological complications. These chronic diseases are prognosticated using various types of data mining classification approaches.*

BIBLIOGRAPHY

- [1] Matthias, H., "Models of software development", Retrieved on 5-OCT-2012, from <http://www.ccs.neu.edu/home/Matthias/670-s05/lectures/2.html>.
- [2] Pressman, R 2001, Software Engineering, 5th edition, McGraw-Hill, New York.
- [3] Ross D.T., J.B. Good enough and C.A. Irvine, Software engineering: Process, Principals, and goals. COMPUTER 8(5) (May 1975): 17-27.
- [4] Cartwright, A. (2008, November 1). Beyond the paper chase. Law Enforcement Technology, 35(11), 58, 60-62.
- [5] Geiger, B. (2004, March 4). Citizens reporting crimes online: the son Francisco experience. The Police Chief, 74(8).
- [6] Coresh J, Selvin E, Stevens LA, et al. Prevalence of chronic kidney disease in the United States. JAMA. 2007;298(17):2038–2047. doi:10.1001/jama.298.17.2038 [PubMed] [CrossRef] [Google Scholar]
- [7] Hsu CY, Vittinghoff E, Lin F, Shlipak MG. The incidence of end-stage renal disease is increasing faster than the prevalence of chronic renal insufficiency. Ann Intern Med. 2004;141(2):95–101. doi:10.7326/0003-4819-141-2-200407200-00007 [PubMed] [CrossRef] [Google Scholar]
- [8] Plantinga LC, Boulware LE, Coresh J, et al. Patient awareness of chronic kidney disease: trends and predictors. Arch Intern Med. 2008;168(20): 2268–2275. doi:10.1001/archinte.168.20.2268 [PMC free article] [PubMed] [CrossRef] [Google Scholar]
- [9] Gansevoort RT, Correa-Rotter R, Hemmelgarn BR, Jafar TH, Heerspink HJ, Mann JF, et al. chronic kidney disease and cardiovascular risk: epidemiology, mechanisms, and prevention. Lancet. 2013. PMID:23727170.
- [10] Go AS, Chertow GM, Fan D, McCulloch CE, Hsu CY. Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. The New England journal of medicine. 2004;351(13):1296–305. Epub 2004/09/24. PMID:15385656.
- [11] Matsushita K, van der Velde M, Astor BC, Woodward M, Levey AS, de Jong PE, et al. Association of estimated glomerular filtration rate and albuminuria with all-cause and cardiovascular mortality in general population cohorts: a collaborative meta-analysis. Lancet. 2010;375(9731):2073–81. Epub 2010/05/21. PMID:20483451.
- [12] Coresh J, Selvin E, Stevens LA, Manzi J, Kusek JW, Eggers P, et al. Prevalence of chronic kidney disease in the United States. JAMA. 2007;298(17):2038–47. PMID:17986697.
- [13] Abramson JL, Jurkovic CT, Vaccarino V, Weintraub WS, McClellan W. Chronic kidney disease, anemia, and incident stroke in a middle-aged, community-based population: the ARIC Study. Kidney Int. 2003;64(2):610–5. Epub 2003/07/09.