

# **Biology of *Apanteles Baoris* Wilkinson (Hymenoptera: Braconide) A Larval Parasitoid of *Chapra Maathas* Fab**

**T. M. Chougale**

Department of Zoology

Bhogawati Mahavidyalaya, Kurukali, MS, India

**Abstract:** *Apanteles baoris* Wilkinson is potential biocontrol agent of *Chapra mathias* Fab. a paddy pest in Kolhapur. It attacks about 86% caterpillars of *C. mathias*. In the present paper life cycle, longevity, Sex ratio and behavioral aspects viz; mating and oviposition have been studied in detail. The life cycle of the parasitoid egg to adult emergence have been completed within 17 days. Egg hatching period was 3 days, larval period was 8-9 days and pupal period was 5-6 days. Mating takes place within one hour of their emergence from cocoons. Mating occurred at morning hours. Oviposition is completed within 3 seconds. In a single host the parasitoid can lay on an average 55 eggs of which progeny production obtained was 46 (average). Sex ratio of adult parasitoids was found favoring females. The results are useful for mass rearing of the above parasitoids.

**Keywords:** *Apanteles Baoris*, *Chapra Mathias*, Biocontrol Agent, etc.

## REFERENCES

- [1] Broodryk., S. W., 1969. The biology of *Chilonus* (*Microchilonus*) *curvimaculatus* Cameron (Hymenoptera: Braconidae) J. ent. Soc. South Afr., 32: 169-189.
- [2] Cardona, C. and Oatman, E. R. 1971. Biology of *Apanteles dignus* (Hymenoptera: Braconidae), a primary parasite of the Tomato pin worm. Ann. ent. Soc. Am., 60: 996-1007.
- [3] Fulton, B.B. 1940. The horn worm parasite, *Apanteles congregatus* (Say) and the hyperparasite *Hypopteromalus tobacum* (Fitch) Ann. ent. Soc. Am, 33: 231-244.
- [4] Kajita. W. Drake, E.F. 1989. Biology of *Apanteles chilonus* and *A. flavipes* (Hymenoptera: Braconidae), parasites of *Chilo suppressalis*. Mushi. 42: 163-179.
- [5] Sathe, T.V. 1986. Biology of *Cotesia diurnii* Rao and Nikam (Hymenoptera: Braconidae), a larval parasitoid of *Exelastis atomosa* Walsingham. Oikoassy, 3 (2), 31-33.
- [6] Sathe T.V., N.B.Gosawi and D.V.Devgire, 1986. Parasite complex associated with *Chapra mathias* Fab., a paddy pest in Kolhapur. Geobios new reports, 5: 59-60.
- [7] Sathe, T.V. 1988. The Biology of *Cotesia orientalis* Chalikwar & Nikam (Hymenoptera: Braconidae), a larval parasitoid of plume moth in India. J. Zool. Res., 1(1): 23-37.
- [8] Sathe, T.V. 1991. The Biology of *Apanteles asawari* Sathe (Hymenoptera: Braconidae), a larval parasitoid of *Spodoptera litura* (Fab.). Oikoassy, 8(1): 15-18.
- [9] Sato, Y. 1975. Rearing *Apanteles glomeratus* L. on the larvae of *Pieris repae crucivora* Biosduval fed on an artificial diet. Kontyu (Tokyo), 43: 242-249.
- [10] Short J.R., T. 1952. The morphology of the head of larval Hymenoptera with special reference to the head of Ichneumonoidea, including a classification of the final instar of Braconidae. Trans. Roy. Entom of Soc. London, 103: 27-84.
- [11] Short J.R. T. 1953. A grouping of larval characters of some species of the genus *Apanteles* (Hymenoptera: Braconidae). Intomol. Res. Bull., 44: 327-33.