

# Total Lipid Alteration Gill and Muscle of Fingerlings of Freshwater Fish *Labeo Rohita* Fed on Formulated Feed

V.B. Nalawade<sup>1</sup>, M.P. Bhilave<sup>2</sup>, M. N. Desai<sup>3</sup>

Assistant Professor, Department of Zoology, The New College, Kolhapur<sup>1</sup>

Professor, Department of Zoology, Shivaji University, Kolhapur<sup>2</sup>

Principal, Krushibhushan Dr. Tanajirao Chorage Arts, Commerce and Science Senior College, Nandwal<sup>3</sup>

**Abstract:** Now a day's consumer wants to know and ensured the nutritional value of the products what they are eating. The biochemical composition of the whole body indicates the fish quality. In the biochemical composition lipids are of great important not only to fish nutrients but also to human nutrition. For this in the present experiment, 90 days experiment was conducted in glass aquaria to study lipid content in intestine of fingerlings of freshwater fish *Labeo rohita* fed on 100%, 75%, 50% and 25% non-conventional source of protein in formulation of fish feed i.e., Blood of bovine animals obtained from slaughter house waste and conventional feed. i.e., Groundnut oil cake. The fishes were fed at the rate of 2% of the body weight every day. After specific time intervals the fishes were weighed and sacrificed for intestine tissue. The estimation of total lipid by Floch's method. The above result demonstrated that, the maximum lipid content was observed at 25% formulated feed followed by 50% 100% and 75% formulated as compared to conventional feed and muscle at 100% formulated feed as compared with conventional feed.

**Keywords:** Lipid, Formulated Feed, Conventional Feed. *Labeo Rohita*, etc.

## REFERENCES

- [1] Rodriguez-Gonzalez, H., Kernandez-Liomas, A. Villarreal, H., Saucedo, P.E., Garcia-Ulloa M. and Rodriguez-Jaramillo, M.C: Gonadal development and biochemical composition of female cray fish *Cherax quadricarinatus* (Decapoda: Parastacidae) in relation. to the gonado somatic index at first maturation Aquaculture, 254,637-645. 2006.
- [2] Maina, R., Beames, P., Mbugua, G. I. and Kisia, S.: Digestibility and feeding value of some feed ingredients fed to tilapia (*Oreochromis niloticus* L.). Aquaculture Research 33: 853 - 862. 2002
- [3] Siddhuraju, P. and Becker, K.: Comparative nutritional evaluation of differentially processed mucuna seeds (*Mucuna pruriens* L.) DC var. utilis (Wall ex Wight) Baker ex Burck, on growth performance, feed utilization and body composition in Nile Tilapia (*Oreochromis niloticus* L.), Aquaculture, 34: 487 - 500.2003
- [4] Islam M. S and Tanaka M: Optimization of dietary protein requirement for pond-reared mahseer, *Tor putitora* Hamilton (Cypriniformes: Cyprinidae). Aquaculture Research 35, 1270-1276.2004
- [5] Nalawade., V.B and Bhilave V. B: Protein efficiency ratio and gross feed conversion efficiency (GFCE) of freshwater fish *Labeo rohita* fed on formulated feed. International Quarterly Journal of Life science The Bioscan 6(2): 301-303. 2011
- [6] NRC: Nutrition requirement of fish, Committee on Animal Nutrition, Board on Agriculture. National Research Council. National Academy Press, Washington, D.C. 1993
- [7] Bhilave, M.P., Nalawade, V.B., Nadaf, S. B and Bhosale S. V: Nutritional analysis of non-conventional feed resource, Uttar Pradesh j. Zoology.3 (1): 87-90. 2011
- [8] Robinson, E. H.; Li, M. H: Comparison of practical diets with and without animal protein at various concentrations of dietary protein on performance of channel catfish *Ictalurus punctatus* raised in earthen ponds. J. World Aquac. Soc 29, 273-280.1998

- [9] Allan G: Fish meal replacement in aquaculture feeds: Sub-program administration. Final report to Fisheries Research and Development Corporation. New South West Fisheries Final Report Series No.7 4pp.1998.
- [10] Crashaw R: Blood meal: A review of its nutritional qualities for pigs, poultry and ruminant animals. National Renderers Association Technical Review. 7p. 1994
- [11] Mohanta K.N, Subramanian S and Korikanthimath V. S: Evaluation of different animal protein sources in formulating the diets for Blue Gourami *Trichogaster Trichopterus* Fingerlings J Aquac Res Development 4-2. 2013
- [12] Agbedi O.T, Otubusin O, Orunleye F. O: Effect of different levels of substitution of fishmeal with blood meal in pelleted feeds on catfish *Clarius gariepinus* (Burchell 1822) culture in net cages. Euro J Sci Res 31:6-10.2003.
- [13] Hertrampf J. W and Piedad Pascual F: Handbook Ingredients for Aquaculture Feeds.Pp 69-78 Kluwer Academic, Dordrecht. 2000
- [14] Medale, F., P. Aguirre and S. J. Kaushik: Utilization of dietary carbohydrates by rainbow trout at two water temperatures. In: Wenk, C., Boessinger, M. (eds), Energy Metabolism of Farm Animals. Proc. 12<sup>th</sup> Symp. Energy Metabolism of Farm Animals, Kartause Ittingen, Switzerland, 1–7 September 1991. Eaap Pub., 58: 392–395.1991.
- [15] Kaushik, S. J. and A. Oliva-Teles: Effects of digestible energy on nitrogen and energy balance in rainbow trout. Aquaculture, 50: 89-101. 1985.
- [16] Barnes, H. and J. Black Stock: Estimation of lipids in marine animals, detail investigation of sulphophosphovanillin method. J. Exp. Mar. Biol. Ecol., 12:103-118. 1973.
- [17] Nadaf S.B and Bhilave M. P: Study of biochemical moieties in freshwater fish, *Labeo rohita* fed on formulated feed. Journal of Aquatic biology and fisheries Vol 2(1) 168-177. 2014.