IJARSCT



International Journal of Advanced Research in Science, Communication and Technology (IJARSCT)

Volume 2, Issue 9, June 2022

Pharmacological Review on Azadirachta indica

Pratik Y. Dhole¹, Mansi N. Gadekar², Swapnil K. Gorade³, Snehal S. Rakshe⁴ Saurabh D. Kaduskar⁵, Sachin V. Datkhile⁶

Samarth Institute of Pharmacy, Belhe, Maharashtra, India^{1,2,3,4,5} Department of Pharmaceutics, Samarth Institute of Pharmacy, Belhe, Maharashtra, India⁶

Abstract: Glashanddical practice seek merge alternative medicine with evidence had medicine for a better underdig of the metabolic process and its effects in the human body. An example is the use of complementary medicine like phytotherapy. Neem has become valuable plant in the world which shows the solutions for hundreds to thousands problems. Neem has become important in the global context today because it offers answers to the major concerns facing mankind. Neem has been extensively used in Ayurveda. Unani and Homoeopathic medicine. Neem and known as Neemba in devbhasha Sanskrit.

Keywords: Antibiotic, Antiviral, Antimicrobial and Antifungal

REFERENCES

- [1]. A. Zong, H. Cao, and F. Wang, "Anticancer polysaccharides from natural resources: a review of recent research," Carbohydrate Polymers, vol. 90, no. 4, pp. 1395–1410, 2012.
- [2]. T. Efferth and E. Koch, "Complex interactions between Phytochemicals. The Multi-Target Therapeutic concept of Phytotherapy," Current Drug Targets, vol. 12, no. 1, pp. 122–132, 201
- [3]. T. R. Govindachari, G. Suresh, G. Gopalakrishnan, B. Banumathy, and S. Masilamani, "Identification of antifungal compounds from the seed oil of Azadirachtaindica," Phytoparasitica, vol. 26, no. 2, pp. 109–116, 1998.
- [4]. N. Singh and M. S. Sastry, "Antimicrobial activity of Neem oil," Indian Journal of Pharmacology, vol. 13, pp. 102–106, 1997.
- [5]. A. Kher and S. C. Chaurasia, "Antifungal activity of essential oils of three medical plants," Indian Drugs, vol. 15, pp. 41–42, 1997.
- [6]. U. Bandyopadhyay, K. Biswas, A. Sengupta et al., "Clinical studies on the effect of Neem(Azadirachtaindica) bark extract on gastric secretion and gastroduodenal ulcer," Life Sciences, vol. 75, no. 24, pp. 2867–2878, 2004.
- [7]. K. Biswas, I. Chattopadhyay, R. K. Banerjee, and U. Bandyopadhyay, "Biological activities and medicinal properties of neem (Azadirachtaindica)," Current Science, vol. 82, no. 11, pp. 1336 1345, 2002
- [8]. Sarkar K, Bose A, Laskar S, Choudhuri SK, Dey S, Roychowdhury PK, et al. Antibody response against neem leaf preparation recognizes carcinoembryonic antigen. IntImmunopharmacol 2007; 7:306-12
- [9]. Bose A, Baral R. Natural killer cell mediated cytotoxicity of tumor cells initiated by neem leaf preparation is associated with CD40-CD40L-mediated endogenous production of interleukin-12. Hum Immunol 2007; 68:823-31.
- [10]. Chakraborty K, Bose A, Pal S, Sarkar K, Goswami S, Ghosh D, et al. Neem leaf glycoprotein restores the impaired chemotactic activity of peripheral blood mononuclear cells from head and neck squamous cell carcinoma patients by maintaining CXCR3/CXCL10 balance. IntImmunopharmacol 2008; 8:330-40
- [11]. PingaleShirish S, Hepatoprotection study of leaves powder of Azadirachtaindica A. juss, International Journal of Pharmaceutical Sciences Review and Research, Volume 3, Issue 2, July August 2010, page no-37-42.
- [12]. K. Girish, Bhat S. Shankara, Neem A Green Treasure, Electronic Journal of Biology, 2008, Vol. 4(3):102-111.
- [13]. BiswasKausik, ChattopadhyayIshita, Banerjee K Ranajit. andBandyopadhyayUday, Biological activities and medicinal properties of neem (Azadirachtaindica), Current Science, Vol-82, 10 june 2002, pp.1336-1345.
- [14]. Sergio Muñoz-Valenzuela, Alberto Arnoldo Ibarra-López, Luis Mariano RubioSilva, HumbertoValdez-

Copyright to IJARSCT DOI: 10.48175/568 15
www.ijarsct.co.in

IJARSCT



International Journal of Advanced Research in Science, Communication and Technology (IJARSCT)

Volume 2, Issue 9, June 2022

- Dávila, and JesúsBorboa-Flores, Neem Tree Morphology and Oil Content, Issues in new crops and new uses. 2007, 126-128
- [15]. Bijauliya RK, Alok S, Kumar M, Chanchal DK and Yadav S: A comprehensive review on herbal cosmetics. Int J Pharm Sci Res 2017; 8(12): 4930-49.
- [16]. Pillai NR and Santhakumari G: Hypoglycemic activity of Meliaazadirachta L.: (Neem). Indian Journal of Medical Research. 1981; 74, 931-933.
- [17]. Satyanarayana K, Sravanthi K, Shaker IA and Ponnulakshmi R: Molecular approach to identify antidiabetic potential of Azadirachtaindica. Journal of Ayurveda and Integrative Medicine. 2015; 6(3): 165-174.
- [18]. Arika WM, Nyamai DW, Agyirifo DS, Ngugi MP and Njagi ENM: In-vivo antidiabetic effect of aqueous leaf extract of Azardirachtaindica, A. juss in alloxan induced diabetic mice. J Diabetic Complications Med 2016; 1(2): 106.
- [19]. Bailey LH. The standard cyclopedia of horticulture. New York: The Macmillan Company; 1953.
- [20]. Butterworth JH, Morgan ED. Isolation of substance that suppresses feeding in locusts. Journal of the Chemical Society, Chemical Communications 1968:23e4.
- [21]. Kapoor LD. Hand book of ayurvedic medicinal plants. Boca Raton, Florida, USA: Herbal Reference Library Edition, CRC Press LLC; 2001. p. 59e61.

DOI: 10.48175/568

- [22]. https://www.greenjeeva.com/blog/top-uses-of-neem-in-the-cosmetics-industry
- [23]. https://www.ajol.info/index.php/njap/article/view/202913