# **IJARSCT**



International Journal of Advanced Research in Science, Communication and Technology (IJARSCT)

Volume 2, Issue 9, June 2022

# **Biological Control of Plant Diseases**

Dr Shri Kant<sup>1</sup> and Dr Ram Kumar Singh<sup>2</sup>

Department of Ag. Botany, C.C.R. (PG) College, Muzaffarnagar, UP<sup>1</sup> Department of Ag. Botany, R.S.M. (PG) College, Dhampur, Bijnor (UP)<sup>2</sup>

Abstract: Every organism gives birth to its offspring in an environment that is suitable for it. Optimal nutrition provides the needed energy. Organisms compete with one another for mastery over one another and for this food. In general, sources of food for the strongest were discovered to be lower creatures or weaker individuals. In the current state of nature, a reversible cycle between microbes and large species is ongoing. Here is a case study with Trichoderma harzianum and Sclerotium rolfsii. Both belong to the Deuteromycetes phylum. However, careful examination revealed that trichoderma consumed sclerotium mycelium, providing relief from the sclerotium-related sickness. Since January 2017, these observations have been taken into consideration. repeated ten times in a row.

Keywords: Biological Control, Phytoremediation, and Plant Diseases

## REFERENCES

- [1]. Alexopolus(1907- 1979) classification of fungi & bergey'smanual (1970).
- [2]. Baum, J.A.; Johnson, T.B. and Carlton, B.C. (2003). Bacillus thuringiensis natural and recombinant bioinsecticide products through vartous Methods.
- [3]. R. and Wyllie, S.G.(2000). The mode of antimicrobial action of essential oil of Melaleuca alternifolia (tea tree oil). J.Appl. Microbiol., 88: 170-175.
- [4]. Crick, H. (2000). The poisoned heart of Africa. New Scientist, 24: 39-42.
- [5]. De Boer, M.; Bom, P.; Kindt, F.; Keurentjes, J.J.B.; van der Sluis, I.; van Loon, L.C. and Bakker, P.A.H.M. (2003). Control of Fusarium wilt of radish by combining Pseudomonas putidastrains that have different disease suppressive mechanisms. Phytopathology, 93: 626-632.
- [6]. Deising, H. and Siegrist, J. (1995). Chitin Deacetylase Activity of the Rust Uromycesviciae-fabae is Controlled by Fungal Morphogenesis. FEMS Microbiol. Lett., 127: 207-212.
- [7]. Dekaboruah, H.P. and Dileep Kumar, B.S. (2003). Rhizoctonia wilt

DOI: 10.48175/568

#### **IJARSCT**



#### International Journal of Advanced Research in Science, Communication and Technology (IJARSCT)

## Volume 2, Issue 9, June 2022

- suppression of brinjal (Solanum melongena L) and plant growth activity by Bacillus BS2. Ind. J. Exp. Biol., 41(6): 627-631.
- [8]. Deshpande, M.V. (2000). Mycopesticide production: Contribution of fungal morphologies in the biopesticide formulations. Nat. Bot. Soc., 55: 11-17.
- [9]. Devoto, A.; Musket, P.R. and Shirasu, K. (2003). Role of ubiquitination in the regulation of plant defence against pathogens. Current Opinion in Plant Biology, 6: 307-311.
- [10].A.M. (2001). Impact of biocontrol Pseudomonas ûuorescens CHA0 and a genetically modiûed derivative on the diversity of culturable fungi in the cucumber rhizosphere. Appl. Environ. Microbiol.,67: 1851-18.

DOI: 10.48175/568