IJARSCT



International Journal of Advanced Research in Science, Communication and Technology (IJARSCT)

Volume 2, Issue 2, July 2022

Formulation and Evaluation of Embilica officinalis

Bhalerao Pooja A¹, Wakchaure Sayali M², Inamdar Sanobar M³, Bhasale Sakshi S⁴, Prof. Shelke D S⁵ Students, B. Pharm, Samarth Institute of Pharmacy, Pune, India^{1,2,3,4}

Department of Pharmacognosy, Samarth Institute of Pharmacy, Pune, India⁵

Abstract: As earlier studies has stated amla is having rich hair nourishing activity and is beneficial for treatment of dandruff problem. The present study is based on combining the capacity of amla with Hibiscus, Henna, Neem, Shikakai and other herbs to obtain enhanced property which can be used to treat the dandruff and problems of whitening of hairs at early age in youth due to exposure to pollutants, improper care and many other factors. The present research aims at formulating and evaluating polyherbal shampoo. The shampoo was tested with different parameters such as anti-dandruff activity, Dirt dispersion test, Foaming ability, Patch test and Stability studies were carried out for 3 months. And based on the all the evaluation test the formulation was found to be stable and can be considered good for applying to scalp and was found to be effective to treat dandruff and also nourish hairs and enhance its appearance.

Keywords: Amla, Anti-dandruff, Henna, Hair tonic

REFERENCES

- [1]. Badi KA, Khan SA. Formulation, evaluation and comparison of the herbal shampoo with the commercial shampoo. Beni-Suef Univ J Basic Appl Sci 2014;3:301-5.
- [2]. Gaud RS, Gupta GD. Practical Physical Pharmacy. 1st ed. New Delhi: C.B.S. Publisher and Distributer; 2001. p. 81-105.
- [3]. Mainkar AR, Jolly CI. Evaluation of commercial herbal shampoos. Int J Cosmet Sci 2000;22:385-91.
- [4]. Klein K. Evaluation of shampoo foam. Cosmet Toilet Mag 2004;119:32-5.
- [5]. Ali HS, Kadhim RB. Formulation and evaluation of herbal shampoo from Ziziphus spina leaves extract. Int J Res Appl Pharm 2011;2:1802-6.
- [6]. Baran R, Maibah HI. Cosmetic dermatology in children. In: Text Book of Cosmetic Dermatology. 2nd ed. London: CRC Press; 1998. p. 507-8.
- [7]. Ireland S, Carlino K, Gould L, Frazier F, Haycock P, Ilton S, et al.Shampoo after craniotomy: A pilot study. Can J Neurosci Nurs 2007;29:14-9.
- [8]. Boonme P, Pakpayat N, Yotmanee K, Kunlawijitrungsee S, Maneenuan D. Evaluation of shampoos containing silicone quaternary microemulsion. J App Pharm Sci 2011;1:59-63.
- [9]. Hahn G. In: Koch HP, Lawson LD, eds. Garlic: the science and therapeutic application of Allium sativum L and related species (2ndedn). Baltimore Williams and Wilkins, 1996: 1–24.
- [10]. Cavallito C, Bailey JH. Allicin, the antibacterial principal of Allium sativum. Isolation, physical properties and antibacterial action. J Am Chem Soc. 66; 1944: 1944–52.
- [11]. Aghel N., Moghimipour B. and Dana R.A., Formulation of a Herbal Shampoo Using Total Saponins of Acanthophyllum squarrosum, Iranian Journal of Pharmaceutical Research. 6(3); 2007: 167-172.
- [12]. Mainkar, A. and Jolly, C. Formulation of natural shampoos. Int. J. Cosmet. Sci. 23(1); 2001: 59–62.
- [13]. Mainkar and Jolly. Evaluation of commercial herbal shampoos. Int. J. Cosmet. Sci. 22(5); 2000: 385–391.
- [14]. Kokate CK, Purohit AP, Gokhale SB "Pharmacognosy" Twenty Fourth Edition published by Nirali Prakashan; (2003); 343.
- [15]. Sorkin M. Shapiro B. and Kass G.S. The practical evaluation of shampoos. J.Soc. Cosmet Chem. 7; 1996: 539-551.

DOI: 10.48175/568