IJARSCT



International Journal of Advanced Research in Science, Communication and Technology (IJARSCT)

Volume 2, Issue 2, July 2022

Pharmacological Activity of Mentha Piperita –A Review

Auti Pranali S¹, Dethe Akanksha A², Bhosale Srushti S³, Atole Shubhangi V⁴, Gadge Shubham C⁵ Samarth Institute of Pharmacy, Belhe, Maharashtra, India^{1,2,3,4}

Department of Pharmacology, Samarth Institute of Pharmacy, Belhe, Maharashtra, India ⁵ pranaliauti7030@gmail.com

Abstract: Mentha piperita L. (Family: Lamiaceae; Synonym: M. balsamea Wild; commonly known as peppermint) is a natural hybrid (M. aquatica ×M. spicata) with immense therapeutic uses (oldest known medicinal plant species, "medicinal plant of the year 2004", known as "heirba Buena" meaning good herb) apart from possessing other potential uses (as flavouring agent from chewing gum to after dinner mints, in cosmetics and pharmaceutical products). Mentha piperita Linn. emend. Huds. is widely used in food, cosmetics and medicines. It has been proven helpful in symptomatic relief of the common cold. It may also decrease symptoms of irritable bowel syndrome and decrease digestive symptoms such as dyspepsia and nausea, although more research is needed. It is used topically as an analgesic and to treat headaches..

Keywords: Mentha piperita, overview, Lamiaceae Peppermint, Peppermint oil, Menthol, Medicine, Toxicity.

REFERENCES

- [1]. Sparks MJ, et al, does peppermint oil relieve spasm during barium enema Br J Radiol, 1995, 68(812), 841-843.
- [2]. Leicester RJ and Hunt RH, Peppermint oil to reduce colonic spasm during endoscopy [letter], Lancet, 1982, II (No.8305), 989.
- [3]. Barbalho SM, et al Metabolic profile of offspring from diabetic Wistar rats treated with Mentha piperita (peppermint). Evid Based Complement Alternate Med 2011; 2011:430237
- [4]. Kokate CK et al, Antibacterial activity of volatile oil of seven species of Mentha, Indian J Microbiol, 1970, 10(2), 45
- [5]. Sawhney SS, et al, Antimicrobial efficacy of some essential oils in vitro, Indian Drugs, 1977, 15, 30.
- [6]. Patil SR, et al Phytochemical, antibacterial and dipterian adulticidal approach. International Journal of Pharmacy and Pharmaceutical Sciences., 2016; 8(3): 352-355.
- [7]. Radaelli M et al. Antimicrobial activities of six essential oils commonly used as condiments in Brazil against Clostridium perfringens. Braz J Microbiol., 2016 Apr-Jun; 47(2):424
- [8]. Behnam S, et al. Composition and antifungal activity of essential oils of Mentha piperita and Lavendula angustifolia on post-harvest phytopathogens. Commun Agric Appl BiolSci 2006; 71(3 Pt B): 1321-6.
- [9]. Magro A, et al. Efficacy of plant extracts against stored-products fungi. Rev Iberoam Micol2006; 23(3): 176-8.
- [10]. Agarwal V, et al Prevention of Candida albicansbiofilm by plant oils. Mycopathologia 2008; 165(1): 13-9.
- [11]. Stanev S, et al Study on essential oil and free menthol accumulation in 19 cultivars, populations, and clones of peppermint (Mentha X piperita).
- [12]. Jullien F, et al. An optimizing protocol for protoplast regeneration of three peppermint cultivars (Mentha x piperita). Plant Cell Tissue and Organ Culture 1998; 54 (3): 153–9.
- [13]. Dorman H J,et al Antioxidant property and composition of aqueous extract from mentha species, hybride variety and cultivars. JAgricFood Chem2003;51(16):4563-9

DOI: 10.48175/IJARSCT-5864