

# Review on Wound Healing Cream of *Jatropha Curcas* Plant Latex

Inamdar Saniya N<sup>1</sup>, Badhekar Akansha S<sup>2</sup>, Shinde Pallavi M<sup>3</sup>, Datkhile Sachin V<sup>4</sup>

Students, Samarth Institute of Pharmacy, Belhe, Maharashtra., India<sup>1,2,3</sup>

Department of Pharmaceutics, Samarth Institute of Pharmacy, Belhe, Maharashtra, India<sup>4</sup>

Department of Pharmacology, Samarth Institute of Pharmacy, Belhe, Maharashtra, India<sup>5</sup>

**Abstract:** *Jatropha curcas* is widely employed in traditional folk medicine. Every a part of the plant has its own advantages in treating diseases. The latex of this plant has strong antimicrobial activity. Secondary metabolites (tannins, saponins and flavonoids) that are present within the latex make it such an honest antibacterial agent. an antibiotic cream is formulated using the dried latex of *Jatropha* because the active ingredient. The potential of antimicrobial activities within the *Jatropha* latex is proven. Thus, antibiotic cream will be wont to kill microorganisms and forestall infections. The cream is additionally shown good physical stability at temperature and normal weather conditions after one month. Since the *Jatropha* latex has medicinal properties, it's potentially to be commercialized in industry.

**Keywords:** *Jatropha*, Latex, Flavonoids, Saponins, *Jatropha curcas*

## REFERENCES

- [1]. Abdelgadir, H.A. and Staden, J.V. (2013) Ethnobotany, ethnopharmacology and toxicity of *Jatropha curcas* L. (Euphorbiaceae): A review. South Afr. J. Bot., 88: 204-218
- [2]. Igbinsosa, O.O., Igbinsosa, E. and Aiyegoro, O.A. (2009) Antimicrobial activity and phytochemical screening of stem bark extracts from *Jatropha curcas* (Linn). Afr. J. Pharm. Pharmacol., 3(2): 58-62.
- [3]. Soares, A.M.S., Carvalho, L.P., Melo, E.J.T., Costa, H.P.S., Vasconcelos, I.M. and Oliveira, J.T.A. (2015) A protein extract and a cysteine protease inhibitor enriched fraction from *Jatropha curcas* seed cake have in vitro anti-Toxoplasma gondii activity. Exp. Parasitol., 153: 111-117
- [4]. Sangeetha, J., Divya, K., Prashanth, M.V., Vamshikrishna, A. and Rani, G.L. (2009) Anti-inflammatory and antibacterial activity of *Jatropha curcas* Linn. J. Pharm. Res. Health Care, 2: 258-262.
- [5]. Warra, A.A. (2012) Cosmetic potentials of physic nut (*Jatropha curcas* Linn.) seed oil: A review. Am. J. Sci. Ind. Res., 3(6): 358-366.
- [6]. Shetty, S., Udupa, S.L., Udupa, A.L. and Venkata, V.R. (2006) Wound healing activities of bark extract of *Jatropha curcas* Linn in albino rats. Saudi Med. J., 27(10): 1473-1476.
- [7]. Esimone, C.O., Nworu, C.S. and Jackson, C.L. (2008) Cutaneous wound healing activity of a herbal ointment containing the leaf extract of *Jatropha curcas* L. (Euphorbiaceae). Int. J. Appl. Res. Nat. Prod., 1: 1-4.
- [8]. Wongkrajang, P., Nurlaly, P.S., Tamsiririrkkul, R., Thongpraditchote, S., Ruangwises, N., Mitrevej, A., Khammanit, R. and Hanyongyuth, R. (2015) Hemostatic effect of n-hexane extracts of *Jatropha curcas* Linn leaf. Mahidol Univ. J. Pharm. Sci., 42(3): 110-117.
- [9]. Oskoueian, E., Abdullah, N., Saad, W.Z., Omar, A.R., Ahmad, S., Kuan, W.B., Zolkifli, N.A., Hendra, R. and Ho, Y.W. (2011) Antioxidant, anti-inflammatory and anti-cancer activities of methanolic extracts from *Jatropha curcas* Linn. J. Med. Plants Res., 5(1): 49-57.
- [10]. Darmawi D, Darniati Manaf Z.H, Syafruddin Sayuti A. Isolation of gram-negative bacteria from metacarpal injury of *Panthera tigris sumatrae* trapped in Subulussalam, Proc. Ann. Int. Con. (AIC) Syiah Kuala Univ. 2011;1:131-135.
- [11]. Raica M, Cimpean A.M. Platelet-derived growth factor (PDGF)/PDGF receptors (PDGFR) axis as target for antitumor and antiangiogenic therapy. Pharmaceutical. 2010;3:572-599.
- [12]. Yu J, Ustach C, Kim H.R.C. Platelet-derived growth factor signaling and human cancer. J. Biochem. Mol. Biol. 2003;36(1):49--59.

- [13]. Priyandari Y, Umatjina SATM: Effect of jarak tree topical increase wound healing excision period of mice. *Journal of Ners Community*. 2015; 6 (2): 198-206.
- [14]. Pratiwi AD, Ratnawati R, Kristianto H: Effect of clove flower bud extract (*Syzygium aromaticum*) on increasing the thickness of epithelialization of incision wounds in white rats (*Rattus norvegicus*) Wistar strain. *FKUB Health Magazine*. 2015; 2 (3): 135-143.
- [15]. Takahashi M, Ogihara H, Nomingerel T, et al.: Repair of the surface epithelium after saponin-induced colonic mucosal injury in the rat. *Medical Electron Microscopy*. 1998; 31: 1–9.
- [16]. Ricardo M, Uribe G, Saldivar S: Anti-inflammatory glycosylated flavonoids as therapeutic agents for the treatment of diabetes-impaired wounds. *Curr Top Med Chem*. 2015; 15 (23): 2456-2463.
- [17]. Li K, Diao Y, Zhang H, et al.: Tannin extracts from immature fruits of *Terminalia chebula* Fructus Retz. promote cutaneous wound healing in rats. *BMC COMPLEMENT ALTERN M*. 2011; 11 (86): 1-9.
- [18]. Muntiaha M.C, Yamlean P.V.Y, Lolo W.A. Test effectivity *Jatropha multifida*, to heal wound infection cause by *Staphylococcus aureus* in rabbit (*Oryctolagus cuniculus*) *Pharmacon*. 2014;3(3):294-302.