

Recent Developments in the Antimicrobial Activity of Heterocyclic Compounds

K Vanaja¹ and Dr. Swapnila²Research Scholar, Department of Chemistry¹Research Guide, Department of Chemistry²

NIILM University, Kaithal, Haryana, India

Abstract: A vast development in multidrug resistance in antimicrobial action enhances the hunt for high-potential novel medicines. Due to their chelation, fine structural flexibility, and chemotherapeutic characteristics, Schiff base heterocyclic metal complexes containing imine moiety are useful antimicrobials. Schiff base compounds from heterocyclic platforms have been well reviewed.

Keywords: heterocyclic, antimicrobial.

REFERENCES

- [1]. Al Mulla A. A review: Biological importance of heterocyclic compounds. *Pharm Chem* 2017;9:141-7.
- [2]. Borad MA, Bhoi MN, Prajapati NP, Patel HD. Review of synthesis of multSpiro heterocyclic compounds from isatin. *Int J Rapid Commun Synth Org Chem* 2013;43:1057.
- [3]. Sönmez M, Şekerci M. A new heterocyclic schiff base and its metal complexes. *Synth Reactivity Inorg Metal-Organic Chem* 2010;34:489-502.
- [4]. Kajal A, Bala S, Kamboj S, Sharma N, Saini V. Schiff bases: A versatile pharmacophore. *J Catalysts* 2013;2013:Article ID: 893512, 14.
- [5]. Tobriya SK. Biological applications of schiff base and its metal complexes-a review. *Int J Sci Res (IJSR)* 2014;3:1254-6.
- [6]. Tadele KT. Antioxidant activity of Schiff bases and their metal complexes: A recent review. *J Pharm Med Res* 2017;3:73-7.
- [7]. Parasha RK, Sharma RC, Govind M. Biological activity of some schiff bases and their metal complexes. *Biol Trace Elel Res* 1989;23:145-50.
- [8]. Bernadette SC, Brian D, Denise A, Egana KK, Georgina R. Anticancer and antifungal activity of copper (II) complexes of quinolin-2(1H)-one-derived schiff bases. *Inorg Chim Acta* 2010;363:4048-58.
- [9]. Singh WM, Dash BC. Synthesis of some new Schiff bases containing thiazole and oxazole nuclei and their fungicidal activity. *Pesticides* 1988;22:33-7.
- [10]. Shaikh AA, Raghuwanshi MG, Khurshid I, Molvi K, Nazim S, Ahmed A. Schiff's bases and amides of selected five membered heterocyclic compounds: A review. *J Chem Pharm Res* 2013;5:14-25.
- [11]. Spinu C, Pleniceanu M, Cristian T. Biologically active transition metal chelates with a 2-thiophenecarboxaldehyde derived Schiff base: Synthesis, characterization, and antibacterial properties. *Turk J Chem* 2008;32:487-93.
- [12]. Mehmet G, Mehmet S, Ismet B. Synthesis, characterization, and antimicrobial activity of a new pyrimidine schiff base and its Cu(II), Ni(II), Co(II), Pt(II), and Pd(II) complexes. *Turk J Chem* 2012;36:189-200.
- [13]. Pratibha MS, Vatsala P, Uma V. Biologically active Co (II), Ni (II), Cu (II) and Mn(II) complexes of schiff bases derived from vinyl aniline and heterocyclic aldehydes. *Int J Chem Technol Res* 2009;1:225-32.
- [14]. Nazk MA, Shaalanand ND, Sahar S. Synthesis, spectroscopic, thermodynamic and biological activity studies of Schiff base and metal complexes derived from 2-[1h-pyrrol-2-ylimino methyl]-5-phenyl-1,3,4-oxadiazole. *Glob J Sci Front Res B Chem* 2015;15:14-8.



- [15]. Asif NK, Ajay P, Sharad T, Jagannath JK, Lokhande MV. Antibacterial activity of 2-[(2-chloro-4-methylbenzylidene) amino] pyridin-4-ol and its some transitional metal ion complexes. IOSR J Appl Chem 2014;7:14-20.
- [16]. Abhishek K, Fernandes J, Pankaj K. Synthesis, antimicrobial and anti-inflammatory studies of some novel schiff base derivatives. Int J Drug Dev Res 2014;6:165-7.
- [17]. Ahmed RM, Yousif EI, Al-Jeboori MJ. Co(II) and Cd(II) complexes derived from heterocyclic schiff-bases: Synthesis, structural characterization, and biological activity, Hindawi publishing corporation. Sci World J 2013;2013:Article ID: 754868, 6.
- [18]. Nair MS, Arish D, Joseyphus RS. Synthesis, characterization, antifungal, antibacterial and DNA cleavage studies of some heterocyclicSchiff base metal complexes. J Saudi Chem Soc 2012;16:83-8.
- [19]. Joseyphus RS, Nair MS. Synthesis, characterization and biological studies of some Co(II), Ni(II) and Cu(II) complexes derived from indole-3-carboxaldehyde and glycylglycine as schiff base ligand. Arabian J Chem 2010;3:195-204.
- [20]. Abd-Elzaher MM, Labib AA, Mousa HA, Moustafa SA, Ali MM, El- Rashedy AA. Synthesis, anticancer activity and molecular docking study of Schiff base complexes containing thiazole moiety, Beni-Suef university. J Basic Appl Sci 2016;5:85-96.
- [21]. Andiappan K, Sanmugam A, Deivanayagam E, Karuppasamy K, Kim HS, Vikraman D, *et al.* *In vitro* cytotoxicity activity of novel schiffbase ligand-lanthanide complexes. Sci Rep 2018;8:3054.
- [22]. Ikram M, Rehman S, Subhan F, Akhtar MN, Sinnokrot MO. Synthesis, characterization, thermal degradation and urease inhibitory studies of the new hydrazide based schiff base ligand 2-(2-hydroxyphenyl)-3- {[(E)-(2-hydroxyphenyl) methylidene] amino}-2,3- dihydroquinazolin- 4(1H)-one. Open Chem 2017;15:308-19.
- [23]. Karabasannavar S, Allolli P, Shaikh IN, Kalshetty MB. Synthesis, characterization and antimicrobial activity of some metal complexes derived from thiazole schiff bases with *in-vitro* cytotoxicity and DNA cleavage studies. Indian J Pharm Educ Res 2017;51:490-501.
- [24]. Chaudhary NK, Mishra P. Metal complexes of a novel schiff base based on penicillin: Characterization, molecular modeling, and antibacterial activity study Hindawi. Bioinorg Chem Appl 2017;2017:Article ID: 6927675, 13.
- [25]. Thakare AP, Mandlik PR. Synthesis, spectroscopic and thermal studies of Fe(III) and VO (IV) complexes of heterocyclic schiff base ligand. Indian J Adv Chem Sci 2017;5:318-23.
- [26]. Nasrin D, Alam MA, Hossain MN, Nazimuddin M. Synthesis, characterization and antimicrobial activity of metal complexes of schiff's base derived from S-benzylthiocarbamate with 2-hydroxyacetophenone. Chem J 2013;3:13-9.
- [27]. Sheheryar,ParveenZ,Rahman T, Zeb mMA, Hassan Z,RehmanW.Synthesisand antibacterial activity of schiff basemetals complexes. Int J Biosci 2017;10:259-64.
- [28]. Selvi ET, Mahalakshmi S. Synthesis and characterization of heterocyclic schiff base ligand derived from 4-aminiantipyrene and thiophene-2- carbaldehyde. Int J Adv Res Dev 2017;2:51-6.
- [29]. Joshi SR, Habib SI. Co (II) and Zn (II) metal complexes of heterocyclic schiff bases: A synthesis, spectral and antimicrobial study. Orient J Chem 2014;30:1343-13.
- [30]. Yamgar RS, Nivid Y, Nalawade S, Mandewale M, Atram RG, Sawant SS, *et al.* Novel zinc(II) complexes of heterocyclic ligands as antimicrobial agents: Synthesis, characterisation, and antimicrobial studies. Bioinorg Chem Appl 2014;2014:276598.
- [31]. Hasan A, Ameer A, Ahmed A, Yousif E. Synthesis and characterization of some transition metal (II) complexes with 1,2,4-triazole schiff base.J Chem Pharm Res 2015;7:531-5.
- [32]. Saadeh SM. Synthesis, characterization and biological properties of Co(II), Ni(II), Cu (II) and Zn (II) complexes with an SNO functionalizedligand. Arabia J Chem 2013;6:191-6.
- [33]. Dikio CW, Ejidike IP, Mtunzi FM, Klink MJ, Dikio ED. Hydrazide schiff bases of acetylacetone metal Complexes: Synthesis, spectroscopic and biological studies. Int J Pharm Pharm Sci 2017;257-67.
- [34]. Manjula B, Antony SA, Dhanaraj CJ. Synthesis, spectral characterization, and antimicrobial activities of schiff base complexes derived from 4-aminoantipyrene. Spectrosc Lett 2014;47:1-9.



- [35]. Jogi P, Padmaja M, Kumar KP, Gyanakumari C. Studies on DNA cleavage and antimicrobial screening of transition metal complexes of a schiff base derived from 2-(aminomethyl)-benzimidazole and p-chloro benzaldehyde. *J Chem Pharm Res* 2012;4:1389-97.
- [36]. Dawood ZF, Al-Bustani RR, Taha M. Biological activity of the complexes of Hg(II), Zn(II) and Cd(II) mixed ligands (thiosemicarbazone and azine) Part II. *Natl J Chem* 2009;36:760-8.
- [37]. Gonewar NR, Jadhav VB, Jadhav KD, Sakure SS, Killedar AA, Sarawadekar RG. Synthesis, characterisation and antimicrobial activity of bivalent metal (Zn, Cd, Hg, Pb and Ag) chelates of 1,2-naphthoquinone dioxime. *IOSR J Pharm* 2012;2:25-33.
- [38]. Mahmoud HW, Mohamed GG, El-Dessouky MM. Synthesis, characterization and *in vitro* biological activity of mixed transition metal complexes of lornoxicam with 1, 10-phenanthroline. *Int J Electrochem Sci* 2014;9:1415-38.
- [39]. Mangaiyarkkarsi P, Arulantony S. DNA cleavage, cytotoxic activities, and antimicrobial studies of some novel schiff base transition metal complexes derived from 4-aminoantipyrine and dihydropyrimidone of vanillin. *Int J Curr Pharm Res* 2016;8:43-7.
- [40]. Ergene E, Sivas H, Benkl K. Biological activities of Cu (II) and Hg (II) complexes of a heptadentate schiff base ligand. *Turk J Biol* 2010;34:379-387.
- [41]. Rajasekar M, Sreedaran S, Prabhu R, Narayanan V, Jagadeesh R. Synthesis characterization and antimicrobial activities of Ni (ii) and cu(ii) schiff base complexes. *J Coord Chem* 2010;63:136-46.