

The Review Article on “Pharmacognostical and Pharmacological Activity of *Pongamia Pinnata*”

Kadam Samiksha S., Sanap Sumedha S., Arote Akshada R., Tapkir Vaibhav S.,
Pokharkar Piyush D., Gaikwad S. D.

Samarth Institute of Pharmacy, Belhe, Maharashtra, India

Abstract: Medicinal plants are important in the traditional medicine and as well as in modern pharmaceutical drugs.^[1] *Pongamia pinnata* exhibits many pharmacological attributes. The plant extract shows anti Diarrhoeal, anti-fungal, anti-plasmodial, anti-ulcer, anti-inflammatory, anti-nociceptive, Anti-hyperglycaemic, anti-hyperammonic, anti-lipoxidative, anti-oxidant and analgesic Activities. For gums, teeth and ulcers, roots are used for cleaning. For bleeding piles, bark Is used internally.^[2] It contains several phyto constituents belonging to category flavonoids And fixed oils. The fruits and sprouts of *P. pinnata* were used in folk remedies for abdominal tumours in India, the seeds for keloid Tumours in Sri Lanka and a powder derived from the plant for tumours in Vietnam. In Sanskrit India, seeds were used for skin Aliments

Keywords: Pongamia pinnata, physicochemical parameters, pharmacological activities

I. INTRODUCTION

The *Pongamia pinnata* tree is richest and brightest trees of India, the name *Pongamia* has conclude from the Tamil name, *pinnata* That indicate to the *pinnata* leaves.^[4] belonging to the family Fabaceae (Papilionaceae). It is also called *Derris indica* and *Pongamia glabra*.^[5] *Pongamia pinnata* is a medium sized evergreen tree with a spreading crown and short stems. It is a shade plant. It is an indo-malaysian species from sea level to 1200m from India to Fiji, is common in alluvial and coastal conditions. Now found in Philippines, Seychelles, Florida, Hawaii, India, Malaysia, Oceania, Australia.^[6] the plant have been used as a crude drug for the treatment of tumours, piles, skin Diseases, itches, abscess, wounds, ulcers, cleaning teeth, dermatopathi, vagainopathi, painful Rheumatic joints^[7]. The seed powder of the plant is given as expectorant in the treatment of bronchitis. An Infusion of *Pongamia* leaves is used to relieve rheumatism. In the treatment of dyspepsia the *Pongamia* seed oil is given as stomachic and cholagogue. By the process of transesterification The seed oil of *Pongamia pinnata* can be converted to biodiesel. The activities such as anti-Diarrhoeal, anti- plasmodium, anti-inflammatory, anti-ulcer, wound healing properties were Reported^[8]. It includes alkaloids, flavonoids, tannins, Hormones, glycosides, karangin, glabrin, kanugin, and fixed Oils, as well as other phytoconstituents. Historically, *P. Pinnata* has been used as a folk medicinal plant, mostly in the Indian medicine systems of Ayurveda and Siddha. The anti-Inflammatory, anti-nociceptive, antioxidant, anti-diarrhoeal, Anti-fungal, anti-plasmodial, anti-ulcer, anti-hyperglycaemic, Anti lipoxidative, anti-hyperammonic and analgesic functions Are available in plant extracts.^[9]

Synonyms:

Pongamia pinnata Linnier is also known by various synonyms such as

- *Millettia pinnata* (L.) Panigrahi
- *Pongamia glabra* vent
- *Derrisindica* (Lam)

Taxonomical classification

Kingdom : Plantae

Order: Fabales Family : Fabaceae

Genus : *Pongamia* (*Millettia*)

Species : *Pinnata*

Binomial name : *Pongamia pinnata* Linn pierre



Fig. Name : Pongamia pinnata

Vernacular names Different vernacular names of P. pinnata have been reported as follow

Languages	Common names
Telugu	Kanuga
Tamil	Punagai,dalkarmacha,pongam,punku
Hindi	Karanja,pongam oil tree,paper,kanji
Bengali	Karach
Sanskrit	Naktamala
Kannada	Honge,hulagilu
Urdu	Karanj
Malayam	Pungu,ungu,unu,avital
Oriya	Koranjo

Origin and geographical distribution: It was naturally distributed in Asia, now this is found in Australia, Florida, Hawaii, India, Malaysia, Oceania, Philippines and Seychelles [4]. It was commonly grown in coastal forests over India and near the streams and rivers.

Ecology:

The best growth is found in well drained sandy loams with assured moisture. It does not grow well on dry sands even though it tolerates salinitic conditions, alkalinity and water locked soils. It will also grow on heavy swelling clay soils. These species have PH higher than 7.5 they become nutrient deficient.

Propagation:

In-situ germination is preferred for this seeds and time limit is within 1-5 weeks of sowing. Planting to the field should occur at the beginning of the next rainy season, when the seedlings are about 60 cms in height. As young plants tolerate shade well a spacing of 7.5x15cms is recommended. Natural reproduction is profuse by seed and commonly by root suckers. Spontaneous seedlings and root suckers may cause critical weed problems.

Temperature: This plant is ready to grow in the temperature of maximum 27-38°C to minimum of 1-16°C. The withstanding temperature of this plant is slightly below 0°C (32°F) and up to about 50°C.

Morphological characteristics

Millettia pinnata is a legume plant that grows to about 15-25 meters in height with large canopy. It has straight or curved trunk and its diameter is about 50-80 cm.**Leaves:**

They are soft shiny reddish-purple when young and mature to a glossy. They are deep green as the season progresses with prominent veins underneath. Imparipinnate leaves of the tree alternate and are short stalked, rounded or cuneate at the base, ovate or oblong along the length. **Flowers:** Flowers are generally with small clusters of white, purple and pink flowers blossoming throughout the year. The raceme like inflorescence bears 2 to 4 flowers which are strongly fragrant and grow to be 15 -18 mm long. Flowering generally starts after 3-4 year.

Calyx

These are bell shaped and truncate

Corolla:

It is rounded ovate shape with basal auricles with a central spot of green colour.

Seeds:

Seeds are about 1.5-2.5 centimeter long with a brittle, oily coat and are unpalatable to herbivores. Brown seed pods appear immediately after flowering and mature in 10-11 months. The pods are thick-walled, smooth, somewhat flattened and elliptical, but slightly curved with a short, curved point. Pod production commences when seedlings are 5-7 years old. The pods generally do not open naturally, and must decay before the seeds can germinate

Stem:

Stem is light green in colour with some irritating odour. It is herbaceous and slightly hard to break. It has smooth texture on its surface.

Root:

The plant has a long, thick taproot and widely spreading lateral roots. The spread of roots on this species, about 9 meters in 18 years, is greater than most other species; moreover it produces root suckers profusely. Because of these characteristics, pongamia is unsuitable for agroforestry and has the potential to become a weed if not managed carefully

Phytochemistry:

Pongamia pinnata seeds contain six compounds (two sterols, three sterol derivatives and one disaccharide) together with the eighty fatty acids (three saturated & five unsaturated). Their structures were elucidated with the help of physicochemical methods and spectroscopic techniques. Karangin, pongagalabrone and pongapin, pinnatin and kanjone have been isolated from seeds. The leaves and stem of the plant consists of flavones and chalcone derivatives such as Pongone, Galbone, Pongalbol, Pongagallone A and B5. The isolation and characterization of five flavonoids pongamones A-E was done from stems of Pongamia pinnata. Their structures were elucidated on the basis of spectroscopic analysis and by comparing their spectroscopy data with related compounds reported in the literature. Around 18 flavonoid compounds including 9 new ones, pongames III-XI from root bark of P. pinnata. The structures were determined to be (2S)-3',4'-dimethoxy-6'',6''-dimethylpyrano[2'',3'':7,8]-flavanone (III), (2S)-6,3',4'-trimethoxy-6''',6'''-dimethylpyrano [2'',3'':7,8]-βavanone (IV), (2S)-7-methoxy-6-O-γ,γ-dimethylallyl-3',4'-methylenedioxyβavanone (V), 2'-hydroxy-3,4,5'-trimethoxy-6'',6''-dimethylpyrano[2''3'':4'3'] chalcone (VI), 2',4'-dimethoxy-3,4-methylene dioxydioxidihydrochalcone (VII), 2',5',β-trimethoxy-3,4-methylenedioxy-6'',6''-dimethylpyrano[2'',3'':4',3'] dihydrochalcone (VIII), 2,β-dimethoxy-3,4-methylenedioxy-furano[2'',3'':4',3']-dihydrochalcone (IX), βhydroxy-2',4',6'-trimethoxy-3,4-methylenedioxychalcone (X) and 3-methoxy-furano-[2'',3'':7,6] flavone (XI), respectively, by means of spectral analysis and synthesis. The fruits of Pongamia pinnata possess three new furanoflavonoid glucosides, pongamosides A-C and a new flavanol glucoside, pongamoside D. The structures of these compounds were established on the basis of spectroscopic studies. This is the first time that furanoflavone glucosides have been found naturally occurring compounds

Pharmacological activities

(i) Antioxidant activity:

The protective role of Pongamia pinnata leaf extract was studied on oxidative stress during ammonium chloride induced hyperammonemia by measuring the extent of oxidative damage as well as antioxidant status. Ethanolic extract of Pongamia pinnata (PPEt) leaves was administered orally (300 mg/kg body weight) and the effects of PPEt on the levels of thiobarbituric acid reactive substances (TBARS), hydroperoxides (HP), conjugated diene (CD), superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx), and reduced glutathione (GSH) were studied in liver and kidney of ammonium chloride-induced hyperammonemic rats. On treatment with PPEt, a significant reduction in the levels of TBARS, HP, and CD and a significant increase in the levels of SOD, CAT, GPx, and GSH in liver and kidney of ammonium chloride-induced hyperammonemia rats were observed, which clearly shows the antioxidant property of PPEt. These findings show the protective role of PPEt against lipid peroxidation and suggest that PPEt possesses antioxidant potential that may be used for therapeutic purposes. The antioxidant property may be due to the presence of flavonoids and polyphenol in the extract. Antioxidant property have been observed that

effect of Pongamia leaf extract on circulatory lipid peroxidation, antioxidant status was evaluated in ammonium chloride – induced hyper ammonium rats enhanced lipid peroxidation in the circulatory ammonium chloride –treated rats was accounted by a significant decrease in the levels of vitamin-C, vitamin-E reduced glutathione peroxidase, superoxide dismutacatalase. It showed that PPEt modulates by reversing the oxidant – anti oxidant imbalance during chloride-induced hyperammonemia and this could be due to its antihyperammonemia effect by means of detoxifying excess ammonia, urea and creatinine and antioxidant property

(ii) Antimicrobial activity:

The plant materials leaf, bark and seeds were extracted successively with petroleum ether and ethyl acetate. The extracts tested for antimicrobial activity against *Bacillus subtilis*, *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Candida albicans* by disc diffusion method. Both the extracts of leaf, bark and seeds showed a good zone of inhibition. Both the extracts showed maximum zone inhibition against *Bacillus subtilis* while *E. coli* and *C. albicans* were unaffected by any of the extracts except petroleum extract . Various extracts of the plant exhibited antibacterial activity against a broad spectrum of gram negative and gram positive bacteria, such as *Proteus vulgaris*, *Staphylococcus epidermidis*, *Staphylococcus aureus*, *Enterobacter aerogenes*, *Bacillus subtilis*, *Salmonella typhimurium*, *Escherichia coli*, *Propionibacterium acne*, *Yersinia enterocolitica*, *Listeria monocytogenes*, *Shigella flexneri* and *Vibrio cholera* . The plant possesses numerous phytoconstituents such as flavones, flavones, chalcone, triterpenes and aromatic carboxylic acids. These compounds seem to be responsible for antibacterial activity of various extracts of parts of *Pongamia pinnata*. The seed oil exhibited significant antifungal activity against tested fungi. The maximum inhibition against *Aspergillus niger* followed *Aspergillus terreus* and *Candida albicans*. The pure oil (100%) showed maximum inhibition and a minimum inhibition by 40-45% of extracted oil against all tested fungi . The first antifungal compound isolated from plant is Triterpene 118. It showed strong antifungal activity against yeast and low activity against molds and no activity against *Penicillium notatum* . The antimicrobial efficacy of ethanolic extract of stems of *Pongamia pinnata* (PPEE) was carried by agar well diffusion method at concentrations of 250µg, 500 µg, 750 µg and 1000 µg against selective Gram positive pathogens such as *Staphylococcus aureus*, *Bacillus subtilis*, *Bacillus megaterium*, *Enterococcus faecalis*, Gram negative *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Proteus vulgaris* and fungi *Aspergillus niger*. The PPEE possess appreciable levels of tannin and flavonoid contents, 39.7±2.6 mg of GAE/gm and 41.2±1.7 mg of rutin/gm of extract respectively. The plant extract was exhibited maximum activity against *Bacillus subtilis* followed by *Escherichia coli* and then against *Bacillus megaterium*. The plant also exhibited significant activity against *Aspergillus niger*. The MIC lies between 31.2 to 62.5 mg/ml . The antimicrobial efficacy of stems of *Pongamia pinnata* against dental caries pathogens like *Streptococcus mutans*, *Staphylococcus aureus*, *Enterococcus faecalis*, *Escherichia coli*, *Lactobacillus acidophilus*, *Pseudomonas aeruginosa* and *Candida albicans* was performed²² . The diameters of zone of inhibition range from 15.33±0.57 to 28.0±1.0 mm at different concentrations. The antimicrobial activity was compared with the standards tetracycline and fluconazole. The crude ethanolic stem extract exhibited maximum activity against *S. aureus* (26.0±1.0 mm) and moderate activity against *E. faecalis* (25.66±1.15 mm) and minimum activity against *P. aeruginosa* (21.66±1.52 mm).The plant also effectively inhibited *C. albicans* (28.0±1.0 mm). The results of the current study demonstrate the antimicrobial activity of stems of *Pongamia pinnata* to prevent dental caries. MIC was performed by agar dilution method and the range was found to be 31.2 mg/ml to 62.5 mg/ml.

(iii) Antiviral activity

The crude aqueous seed extract showed antiviral activity. It completely inhibited growth of herpes simplex virus type1(HSV-1) and (HSV-2) at the concentration of 1 and 20 mg /ml (w/v) respectively and showed complete absence of cytopathic effect . The crude dried leaves extract showed no activity against rota virus. *Pongamia pinnata* seed extract²⁴ showed the safety effect in acute and chronic toxicological studies conducted in swiss albino rats. The antiviral activity of bis (2-methylheptyl) phthalate isolated from *Pongamia pinnata* leaves against White Spot Syndrome Virus (WSSV) of *Penaeus monodon* Fabricius was evaluated. The oral administration of ethanolic extract and purified compound bis (2-methylheptyl) phthalate has increased the survival of WSSV infected *Penaeus monodon*. The shrimp were fed with the pelletized feed impregnated with ethanolic extract of the leaves of *Pongamia pinnata* prior and after WSSV infection at 200 and 300 mcg/g of body weight of shrimp/day. The survival rate for the WSSVinfected shrimp that were fed with 200 and 300 mcg extract/g were 40% and 80%, respectively .

(iv) Antidiarrhoeal Activity:

This activity was determined by evaluating antimicrobial effect of crude decoction of dried leaves of *Pongamia pinnata*. It also evaluated for its effect on production and action of Enterococcus (cholera toxin, *Escherichia coli* labile toxin, stable toxin) and adherence of enteropathogenic *E. coli* and invasion of enteroinvasive *E. coli* and *Shigella flex* epithelial cells. This study concludes that decoction of *Pongamia pinnata* had selective anti-diarrhoeal action with against cholera and enteroinvasive bacterial causing bloody diarrhoeal episode .

(v) Antiprotozoal activity

The plant was reported for its anti-plasmodial activity against *Plasmodium falciparum* . The bark and leaf extract with low IC50 values of 9-43 mcg/ml has been shown to be potential as anti-malaria by possessing anti-plasmodial activity. This activity might be due to the presence of lupeol (120), which blocked the invasion of *Plasmodium falciparum* merozoites into erythrocytes at IC50 1.5 mcg/ml. It also reported the inhibition of growth of *Trypanosoma cruzi* and *Leishmania* with an IC90 at the dose of 100 mcg/ml . Crude decoction of dried leaves had no activity against trophozoites.

(vi) Anti-Inflammatory

activity It has been reported that the 70% ethanolic leaf extract of *P. pinnata* possess potent anti-inflammatory activity against different phases (acute, sub-acute and chronic) of inflammation without side effect on gastric mucosa. It also showed significant anti pyretic action of the extract against brewer's yeast-induced pyrexia

(vii) Anti-Ulcer activity

The methanolic extract of roots of *P. pinnata* reported for significant protection against mucosal damage induced by aspirin and has a tendency to decrease acetic-acid induced ulcer after 10-days treatment. The extract showed ulcer protective effect with cessation of mucosal defensive factors like mucin secretion, life span of mucosal cells, mucosal cell glycoproteins, cell proliferation and prevention of lipid peroxidation . The methanolic extract of seeds was evaluated for the ulcer protective and healing effect in rats. When administered orally, the extract showed dose-dependent (12.5-50mg/kg for 5 days) ulcer protective effect against gastric ulcer induced by 2h cold resistant stress. Optimal effective dose of PPSM (25mg/kg) showed anti-ulcerogenic activity against acute gastric ulcers (GU) induced by pylorous ligation and aspirin and duodenal ulcer induced by cysteamine but not against ethanol-induced GU .

viii) Anti-hyperglycemic & anti-lipid peroxidative activity

It has been reported that oral administration of ethanolic extract of *P. pinnata* flower shows significant antihyperglycemic and anti-lipid peroxidative effect and enhancement in antioxidant defense system in alloxan-induced diabetic rats. Oral administration of the ethanolic extract of the flower (300mg/kg bw) showed significant antihyperglycemic activity which considerably reduce the blood glucose concentration in a similar extent to that of the reference drug glibenclamide (600microgram/kg bw) in alloxan -induced diabetic rats. The results suggested that the treatment of *Pongamia pinnata* extract could be used as a safe alternative anti-hyperglycaemic drug for diabetic patients .

II. CONCLUSION

In traditional system of Ayurvedic medicine *Pongamia pinnata* has been widely used as curative agents for variety of ailments. In the traditional systems of medicines, such as Ayurveda and Unani, the plant is used for anti-inflammatory, anti-plasmodial, anti-nociceptive, anti-hyperglycaemic, antilipidperoxidative, anti-diarrhoeal, anti-ulcer, antihyperammonic, anti- oxidant and antibacterial. The extensive literature survey revealed that *Pongamia Pinnata L.* is an important versatile medicinal plant with diverse pharmacological spectrum. The plant shows the presence of many chemical constituents which are responsible for varied pharmacological and medicinal properties. However, evaluation needs to be carried out on *Pongamia Pinnata L.* in order to explore the concealed areas and their practical clinical applications, which can be used for the welfare of the mankind.

ACKNOWLEDGEMENT

We acknowledge profound gratitude to the management of Vignan Pharmacy College, Guntur for providing infrastructure facilities to conduct this work.

REFERENCES

- [1]. A phyto pharmacological review on a versatile medicinal plant: Pongamia pinnata (L.) pierreSowjanyaPulipati, P Srinivasa Babu, D Neelima Lakshmi, N NavyasriHarshini, J Vyshnavi and M PrasanthJournal of Pharmacognosy andPhytochemistry 2018;journal of Pharmacognosy and Phytochemistry; 7(4): 459-463
- [2]. A Review on Pongamia pinnata (L.): Traditional Uses, Phytochemistry and Pharmacological Properties ;Akshay G. Fugare , Rajkumar V. Shete , Vishal S. Adak , Krishna Murthy G. Journal of Drug Delivery and Therapeutics 2021: 207
- [3]. Pongamia pinnata: Phytochemical constituents, Traditional uses and Pharmacological properties: A reviewV.V. Chopade, A.N. Tankar1, V.V. Pandel, A.R. Tekade1, N.M. Gowekar1S.R.Bhandari1,S.N. Khandake1; International Journal of Green Pharmacy; 2008 : 72
- [4]. Pramila Ghumare, D B Jirekar, Mazahar Farooqui and S D Naikwade, "A review of Pongamia pinnata – An Important medicinal plant," Current Research in Pharmaceutical Sciences 2014; 04(02): 44-47.
- [5]. Allen ON, Allen EK. The leguminosae. The UniversityofWisconsinPress; 1981. p. 812.
- [6]. S. K. Sarje, Shagufta Farooqui, Shinde Punam, Sontakke Shital and Narote Mayur, "Phytochemical, Pharmacognostic and Quantitative estimation of Pongamia Pinnata Leaves Extract-A Preliminary study To Identified Phytoconstituents," World Journal of Pharmaceutical Research, Volume 9, Issue 3, 2020, 1096-1112.
- [7]. Tanaka T, Iinuma M, Fujii Y, Yuki K, Mizuno M. Flavonoids in root bark of Pongamia pinnata. Phytochemistry. 1992; 31:993-98.Essa MM, Subramanian P. Protective role of Pongamia pinnata leaf extract on tissue antioxidant status and lipid peroxidation in ammonium chloride-induced hyperammonemic rats. Toxicology Mechanisms and Methods. 2006; 16(9):477-483
- [8]. Essa MM, Subramanian P. Protective role of Pongamia pinnata leaf extract on tissue antioxidant status and lipid peroxidation in ammonium chloride-induced hyperammonemic rats. Toxicology Mechanisms and Methods. 2006; 16(9):477-483
- [9]. Meera B., Kumar S., Kalidhar S.B., 2003. A review of the chemistry and biological activity of Pongamia pinnata. J. Med. Aromatic. Plant Sci. 2003; 25:441-65