

# Antibacterial and Antioxidant Study of *Zingiber Officinale*

**R. M. Kadam\*, W. N. Jagirdar and N. S. Bagul**

Department of Botany, Mahatma Gandhi Mahavidyalaya, Ahmedpur-413512, (M.S.), India

Department of Botany, Mahatma Gandhi Mahavidyalaya, Ahmedpur-413512, (M.S.), India

Department of Biotechnology, Modern College, Ganeshkind, Pune-411007, (M.S.), India

Corresponding Author: rmk76@rediffmail.com

**Abstract:** Ethanol, chloroform, carbon tetrachloride methanol, and hexane extracts from and root of *Zingiber officinale* were investigated for their in vitro antimicrobial properties. a disk-diffusion and minimal inhibition concentration (MIC) method . All extract of *Zingiber officinale* (Bel) were different in terms of their antibacterial activities. The ethanol act showed a stronger and broader spectrum of antibacterial activity, study was also carried out to evaluate the in-vitro antioxidant activities of ethanol ,chloroform and carbon tetrachloride extract of *Zingiber officinale* species namely This was achieved by screening the two plant extracts at varying concentrations (10-50ig/ml)using DPPH radical scavenging activity. The results were analyzed statistically which showed that ethanol extract *Zingiber officinale* has more antioxidant activity than standard antioxidant.

**Keywords:** plant, Antioxidant activity, Antibacterial activity, DPPH Test

## I. INTRODUCTION

Ginger is a tropical plant that was spread around the tropical world during the colonial days. In Africa, Asia, or South America, and many hot spots between those continents, locals have taken to using ginger widely in medicine and food. The entire ginger family is rich in oils that both kill micro-critters and stimulate the immune system to do the same. The ways it stimulates the body are many, and, whether added to food or taken as a medicinal tea<sup>1-5</sup>, ginger makes your body a little bit stronger and a little bit better able to resist the damaging forces of nature. Here are the basics for using ginger as a medicine.

Damage to cells caused by free radicals is believed to play a central role in the aging process and in disease progression. Antioxidants are our first line of defense against free radical damage, and are critical for maintaining optimum health and wellbeing. The need for antioxidants becomes even more critical with increased exposure to free radicals<sup>6-10</sup>. Pollution, cigarette smoke, drugs, illness, stress, and even exercise can increase free radical exposure. Because so many factors can contribute to oxidative stress<sup>11-16</sup>, individual assessment of susceptibility becomes important. Many experts believe that the Recommended Dietary Allowance (RDA) for specific antioxidants may be inadequate and, in some instances, the need may be several times the RDA<sup>17-18</sup>.

## II. MATERIAL AND METHOD

### 2.1 Plant Material

The plant materials used in this study, *Zingiber officinale* roots, seed and leaves of were collected from the field in panvel region of Raigad District, identified by Dr. Varkhed KBP College Vashi Navi Mumbai Maharashtra, India. A voucher specimen of the collected sample was deposited in our institutional herbarium for the reference. Preparation of various extract of *Zingiber officinale*.

In present study we use dry stem of the plant collected from western ghat region Maharashtra. Dried Fruit, Leaves and Seeds are cut into small pieces these pieces are then grinded. The grinded sample is dark or gray brown in colour with a special smell.

This powder of all three separately stirred in non-polar solvent Carbon tetra chloride, for 1/2 hour & then it is refluxed for 1/2 hour this is performed for extraction of non-polar component from powder. After extraction the Carbon tetra

chloride layer is distilled to recover solvent & to get a brown colored liquid fraction which shows single spot on thin layer chromatography.

The residue of Carbon tetra chloride extraction is used for further study. This residue is mixed with Chloroform and stirred for 1/2 hour & then refluxed for 1 hour. After filtration the filtrate is distilled to get Chloroform Fraction which is Red-brown colored liquid.

Then the Residue of Chloroform is used for extraction with Ethyl acetate stirred well & refluxed for 1 hour then filtered. Filtrate is then distilled & fraction of Ethyl acetate is collected it shows no spot on TLC plate. Conclusion is that no organic compound is present.

The Ethyl acetate residue is further mixed with ethanol & stirred for 1/2 hr and refluxed for 1hr. Then it is filtered & filtrate is distilled out. Ethanol fraction is yellow brown in colour & show three spot on TLC plate .All extract Collected in Bottle and studied for various activities.

### 2.2 Antioxidant Activity

DPPH Scavenging Test: Quantitative measurement of radical scavenging property was carried out in a universal bottle. The reaction mixture contained 50 µL of test samples (or 80% MeOH as blank) and 5 mL of a 0.004% (w/v) solution of DPPH in methanol. Different known antioxidants, vitamin E, and butylated hydroxytoluene (BHT, Sigma) were used for comparison or as a positive control. Discoloration was measured at 517 nm after incubation for 30 min. Measurements was taken at least in triplicate. DPPH radical's concentration was calculated using the following equation: DPPH scavenging effect (%) = [(Ao – A1) / Ao] × 100; Where Ao was the absorbance of the control and A1 was the absorbance in the presence of the sample. The actual decrease in absorption induced by the test compounds was compared with the positive controls. The mean OD 517 results of DPPH scavenging activity were recorded.

### 2.3 Antimicrobial Activity

The agar diffusion method<sup>[11]</sup> was used to evaluate the antimicrobial activity. Bacteria were cultured overnight at 37 ° C in Mueller Hinton 10 µl Broth and fungi at 28 ° C for 72h in Potato Dextrose Broth (PDB, Oxide) and used as inoculums. A final inoculums, using 100 µl of suspension containing 10<sup>8</sup> CFV/ml of bacteria 10<sup>4</sup> spore/ml of fungi spread on Mueller Hinton Agar (MHA) and Potato Dextrose Agar (PDA) medium respectively.

The disc (6 mm in diameter) was impregnated with 10 µl of 75 µl/ml, 50 µl/ml, 25 µl/ml, 10 µl/ml and 5 µl/ml of each extracts and for each organism placed on seeded agar. Ciprofloxacin and Fluconazole (75 µl/ml, 50 µl/ml, 25 µl/ml, 10 µl/ml and 5 µl/ml) were used as positive control bacteria and fungi respectively.

The test plates were incubated at 37 ° C for 24h for bacteria and at 28 ° C for 72h for fungi depending on the incubation time required for a visible growth.

MIC values were also studied for microorganisms by turbid metric method, which were determined as sensitive to the extracts in cup plate method. MIC was defined as the lowest concentration of extract that inhibit visible growth.

## III. OBSERVATION TABLES

**Table-1 Antibacterial activity of Ethanolic extracts seeds *Zingiber officinale***

Bacterial	Extract Ethanolic	Extract CCl <sub>4</sub>	Cefotax	Pencil	Tetrax
G (+)	In mm	In mm	In mm	In mm	In mm
1. <i>Staphylococcus epidermidis</i>	10	9.2	10	10	9
2. <i>Staphylococcus aureus</i>	08	10.6	10	8	8
3. <i>bacillus paludisme</i>	12	6.4	11	10	8
4. <i>Bacillus subtilis</i>	14	9.4	11	10	7
G (-)					
1. <i>Escherichia coli</i>	5	8.7	6	5.5	5.5
2. <i>Pseudomonas aeruginosa</i>	8	7.8	6	7	4.5
3. <i>Bacillus subtilis</i>	7	8.5	4.5	8.5	9
4. <i>Enterobacter aerogenes</i>	4	5.7	6	5	6

**Table-II: Antifungal activity of Ethanolic extracts from extract from seeds *Zingiber officinale***

Fungus	Extract Ethanolic	Extract CCl <sub>4</sub>	Cefotax	Penicil	Tetrax
1. <i>Candida albicans</i>	6.5	6.8	7	5	9
2. <i>Aspergillus fumigatus</i>	8.3	3.6	5	7	4
3. <i>Aspergillus niger</i>	4.7	10.5	10	8	9

**Table-III :Antibacterial activity of methanolic extract from root of *Zingiber officinale***

Bacteria	Extract Ethanolic	Extract CCl <sub>4</sub>	Cefotax	Penicil	Tetrax
G (+)	In mm	In mm			
1. <i>Staphylococcus epidermidis</i>	6.6	7.2	10	10	9
2. <i>Staphylococcus aureus</i>	8.8	7	10	8	8
3. <i>bacillus paludisme</i>	6.8	9.7	11	10	8
4. <i>Bacillus subtilis</i>	9.8	5.4	11	10	7
G (-)					
1. <i>Escherichia coli</i>			6	5.5	5.5
2. <i>Pseudomonas aeruginosa</i>	7.3	3.4	6	7	4.5
3. <i>Bacillus subtilis</i>	8.5	6.3	4.5	8.5	9
4. <i>Enterobacter aerogenes</i>	8.2	7.4	6	5	6
1. <i>Staphylococcus epidermidis</i>	8.3	4.3	10	10	9

**Table-IV: Antifungal activity of Ethanolic extract from root of *Zingiber officinale***

Fungus	Extract ethanolic	Extract CCl <sub>4</sub>	Cefotax	Penicil	Tetrax
1. <i>Candida albicans</i>	4.0	5	7	5	4
2. <i>Aspergillus fumigatus</i>	7.2	8	5	6	4
3. <i>Aspergillus niger</i>	4.2	8	6	4	5

**Table-V: Antioxidant activity of Fruits**

Extract Conc. Mg/ml	BHT	Ethanol	CHCl <sub>3</sub>	CCl <sub>4</sub>
0.05	45.1	20.60	21.53	17.47
0.1	46.91	36.64	14.53	10 .70
0.2	49.24	24.24	27.50	24 .50
0.3	57.57	40.12	44.00	30 .50

**TABLE VI: Antioxidant activity of root**

Extract Conc. Mg/ml	BHT	Ethanol	CHCl <sub>3</sub>	CCl <sub>4</sub>
0.05	45.1	18	15	17
0.1	46.91	21	23	25
0.2	49.24	23	27	29
0.3	57.57	45	30	267

#### IV. RESULTS AND DISCUSSION

There is a strong need for effective antioxidants from natural sources as alternatives to synthetic antioxidant in order to prevent the free radicals implicated diseases which can have serious effects on the cardiovascular system, either through lipid per oxidation or vasoconstriction

The extracts and essential oils of many plants have been investigated for their antioxidant activity<sup>5-7</sup>. Secondary metabolites such as polyphenols are not required for plant development and growth, but are involved in plant communication and defense<sup>8-9</sup>. Polyphenols interact with pathogens, herbivores, and other plants; they protect from

ultraviolet radiation and oxidants, repel or poison predators and attract beneficial insects or microbes<sup>10-11</sup>. Therefore, in this study, the antioxidant properties of the methanol extracts of leaves and stems of plant like of reexamined for DPPH radical scavenging activity according to the method described and the results of the screening are shown in table 1 & table 2 as comparable with known antioxidant BHT.

In terms of antioxidant activity, all the extracts investigated exhibited a rather high degree of activity (more than 40%). In particular, fruits (ethanol extract) of *Zingiber officinale* displayed the highest activities as antioxidant activity as removal of the stable radical DPPH and the lowest activity were found in CCl<sub>4</sub> extract of stem. As expected, the overall activity of the raw extracts was lower than that of commercial antioxidant BHT, the reference antioxidant.

The results of Antimicrobial activity were done for all the five, pet ether, chloroform, acetone, and methanol and aqueous extracts. During antimicrobial study methanolic extracts showed maximum zone of inhibition against almost all organisms in cup plate method.

### V. CONCLUSION

The ethanolic extract from roots of *Zingiber officinale* showed a good inhibition against all the bacterial Strains tested (MIC between 10&80 ug/ml). The gram (+) bacteria were sensitive with gram (-) bacteria and some common fungi.

### REFERENCES

- [1]. Parr, A.; Bolwell, G. P. Phenols in the plant and in man: The potential for possible nutritional enhancement of the diet by modifying the phenols content or profile. *J. Sci. Food Agric.* 2000, 80,985–1012.
- [2]. Basile, A.; Ferrara, L.; Del Pozzo, M.; Mele, G.; Sorbo, S.; Bassi, P.; Montesano, D. Antibacterial and antioxidant activities of ethanol extract from *Paullinia cupana* Mart. *J. Ethnopharmacol.* 2005, 102, 32-36.
- [3]. Lorenzi, H.; Matos, F. J. A. *Plantas medicinais no Brasil: nativas e exóticas*. Instituto Plantarum: São Paulo, 2002; pp 544-546.
- [4]. Coelho, L. P.; Reis, P. A.; Castro, F. L.; Gayer, C. R. M.; Lopes, C. S.; Silva, M. C. C.; Sabino, K.C. C.; Todeschini, A. R.; Coelho, M. G. P. Antinociceptive properties of ethanolic extract and fractions of *Pterodon pubescens* Benth. seeds. *J. Ethnopharmacol.* 2005, 98, 109-119.
- [5]. Brito, A. R. M. S.; Brito, A. A. S. Forty years of Brazilian medicinal plant research. *J. Ethnopharmacol.* 1993, 39, 53-67.
- [6]. Duarte, I. D. G.; Ferreira-Alves, D. L.; Veloso, D. P.; Nakamura-Craig, M. Evidence of the involvement of biogenic amines in the antinociceptive effect of a vouacapan extracted from *Pterodon polygalaeflorus* Benth. *J. Ethnopharmacol.* 1996, 55, 13-18.
- [7]. Finkel, T.; Holbrook, N. J. Oxidants, oxidative stress and the biology of ageing. *Nature* 2000, 408,239-247.
- [8]. Halliwell, B. The antioxidant paradox. *The Lancet* 2000, 355, 1179- 1180.
- [9]. Pietta, P. Flavonoids as antioxidant. *J. Nat. Prod.* 2000, 63, 1035- 1042.
- [10]. Visioli, F.; Keaney Jr, J. F.; Halliwell, B. Antioxidants and cardiovascular disease; *panaceas or tonics for tired sheep?* *Cardiovasc. Res.* 2000, 47, 409-418.
- [11]. Husain, S. R.; Cillard, J.; Cillard, P. Hydroxyl radical scavenging activity of flavonoids. *Phytochemistry* 1987, 26, 2489-2497.
- [12]. African pharmacopoeia, 1st ed. Lagos, Organization of African Unity Technical & Research Commission, 1985.
- [13]. British pharmacopoeia. London, Her Majesty's Stationery Office, 1988.
- [14]. Deutsches Arzneibuch 1996. Stuttgart, Deutscher Apotheker Verlag, 1996.
- [15]. European pharmacopoeia, 2nd ed. Strasbourg, Council of Europe, 1995.
- [16]. Pharmacopée française. Paris, Adrapharm, 1996.
- [17]. Pharmacopoea hungarica VII. Budapest, Medicina konyvkiado, 1986.
- [18]. The Japanese pharmacopoeia XIII. Tokyo, Ministry of Health and Welfare, 1996.
- [19]. Morazzoni P, Bombardelli E. Valeriana officinalis: traditional use and recent evaluation of activity. *Fitoterapia*, 1995, 66:99–112.
- [20]. Youngken HW. Textbook of pharmacognosy, 6th ed. Philadelphia, Blakiston, 1950.

- [21]. Bisset NG. Max Wichtl's herbal drugs & phytopharmaceuticals. Boca Raton, FL, CRC, Press, 1994.
- [22]. Farnsworth, NR. ed. NAPRALERT database. Chicago, University of Illinois at Chicago, IL, March 15, 1995 production (an on-line database available directly through the University of Illinois at Chicago or through the Scientific and Technical Network (STN) of Chemical Abstracts Services).
- [23]. Bruneton J. Pharmacology, phytochemistry, medicinal plants. Paris, Lavoisier 1995.
- [24]. Jackson BP, Snowden DW. Atlas of microscopy of medicinal plants, culinary herbs and spices. Boca Raton, FL, CRC Press, 1990.