

Biological Activity of Transition Metal Schiff Base Complexes

Krishan Kumar Arya

Department of Chemistry,
Government Model Degree College Arniya Bulandshahr

Abstract: *Schiff bases — organic compounds formed by the condensation of primary amines with carbonyl compounds — have attracted sustained research interest for decades, largely because of their versatile coordination chemistry and their remarkable range of biological activities. When these ligands bind transition metals such as copper, cobalt, nickel, zinc, and vanadium, the resulting complexes frequently exhibit biological potency that exceeds the free ligand alone, a phenomenon that has driven extensive investigation into their antimicrobial, antifungal, anticancer, antioxidant, and antidiabetic properties. This article reviews the current state of knowledge on the biological activity of transition metal Schiff base complexes, examining the structural features that govern activity, the mechanisms through which these complexes interact with biological targets, and the evidence base for each major activity class. Particular attention is given to the role of the central metal ion in modulating biological behavior, the influence of donor atom sets and geometric configuration, and the emerging evidence for DNA binding and enzyme inhibition as mechanistic pathways. The article also addresses current limitations in the field — including gaps between in vitro activity and in vivo efficacy — and identifies the directions where the chemistry is most likely to produce clinically relevant outcomes.*

Keywords: *Schiff base complexes, antimicrobial activity, DNA binding, coordination chemistry, transition metals, anticancer activity*

I. INTRODUCTION

There is something elegant about Schiff base chemistry that keeps drawing chemists back to it. The reaction is simple — mix a primary amine with an aldehyde or ketone, lose a molecule of water, and you get an imine linkage — but what comes out of that reaction can be structurally sophisticated, electronically tunable, and biologically active in ways that took decades of research to begin unpacking. Hugo Schiff described the condensation reaction that bears his name in 1864, and the compounds he inadvertently opened the door to have been studied ever since.

The real excitement, at least from a medicinal chemistry perspective, comes when these ligands meet transition metals. Schiff bases are excellent chelating agents. The imine nitrogen, combined with oxygen, sulfur, or additional nitrogen donor atoms elsewhere in the molecule, creates binding pockets that transition metal ions occupy readily, forming stable five- or six-membered chelate rings. Copper, cobalt, nickel, zinc, vanadium, manganese — all of them form complexes with Schiff base ligands, and those complexes behave biologically in ways that neither the free metal ion nor the free ligand predicts particularly well.

Why does complexation change biological activity so dramatically? Several reasons have been proposed and explored. The metal center can enhance membrane permeability of the ligand. Coordination geometry and the electronic environment around the metal affect how the complex interacts with enzymes, DNA, and cell membranes. Redox-active metal centers like copper and cobalt can generate reactive oxygen species under biological conditions, adding cytotoxic mechanisms that purely organic compounds cannot replicate. The lipophilicity of the complex — which governs how readily it crosses cell membranes — is tunable through ligand design in ways that give medicinal chemists real synthetic control over biological behavior.



This article reviews the biological activities that have been documented for transition metal Schiff base complexes, with particular focus on the activities that have generated the most research interest and the most convincing mechanistic evidence: antimicrobial and antifungal activity, anticancer activity, antioxidant capacity, and antidiabetic properties. Along the way, it examines what structural features tend to produce the strongest activities and where the gap between laboratory findings and clinical application remains wide.

II. CHEMISTRY OF SCHIFF BASE LIGANDS AND THEIR METAL COMPLEXES

2.1 Synthesis and Structural Features

Schiff bases form readily under mild conditions — typically by refluxing an amine and a carbonyl compound in ethanol or methanol, often with a catalytic amount of acid. The reaction is reversible in principle but driven to completion by removal of water. The resulting imine bond (C=N) is the defining structural feature, but what makes Schiff bases so useful as ligands is the arrangement of donor atoms that typically flanks it.

The most biologically studied Schiff base ligands are bidentate or tridentate, offering the metal ion two or three coordination sites. An N,O-donor Schiff base — where the imine nitrogen and a phenolic or alcoholic oxygen both bind to the metal — is perhaps the most common structural motif in the literature. Adding a second phenolic arm creates an N,N,O or O,N,O donor set, giving access to octahedral coordination geometries with first-row transition metals. Sulfur-containing Schiff bases incorporating thiosemicarbazone or thiol groups add S-donor atoms that have particular relevance for biological activity, since sulfur coordinates strongly to soft metal ions and many biologically important metal sites.

The geometry around the metal center — tetrahedral, square planar, octahedral — influences biological activity in ways that are still being worked out. Square planar copper(II) and nickel(II) complexes, for instance, tend to intercalate more effectively into DNA double helix structures than tetrahedral complexes, because their planar geometry allows them to stack between base pairs. This geometric selectivity has direct consequences for the anticancer and antimicrobial activities discussed later.

2.2 The Role of the Metal Ion

Choosing which metal to incorporate into a Schiff base complex is not arbitrary — different metals bring fundamentally different biological properties. Copper(II) complexes are among the most extensively studied, partly because copper is an essential biological metal with established roles in enzyme catalysis, and partly because Cu(II) complexes tend to show strong DNA binding and can participate in oxidative DNA damage through Fenton-like chemistry. Cobalt(III) complexes are kinetically inert, which makes them attractive for targeted delivery applications where controlled release of the metal is desirable. Nickel(II) complexes show interesting DNA binding properties and moderate cytotoxicity. Zinc(II) complexes are generally less toxic than copper or cobalt analogues, which makes them more interesting for antimicrobial applications where selectivity for microbial over mammalian cells is important. Vanadium complexes, particularly oxovanadium(IV) species, have attracted attention specifically for antidiabetic activity because of vanadium's known insulin-mimetic properties.

III. ANTIMICROBIAL AND ANTIFUNGAL ACTIVITY

3.1 Evidence and Trends

Researchers have done extensive studies about how transitional metal Schiff base complex compounds kill microorganisms because this research method provides simple testing methods. Most chemistry laboratories have access to standard disc diffusion testing and minimum inhibitory concentration testing, which enables researchers to generate antimicrobial research results at a fast pace. The literature shows a consistent pattern which provides usable results because metal complexation improves antimicrobial activity of Schiff base ligands, while the extent of this enhancement depends on metal, ligand structure, and target organism. The broadest and strongest antimicrobial profiles emerge from Copper(II) and Zinc(II) complexes, which show effectiveness against *Staphylococcus aureus*, *Bacillus*



subtilis, Escherichia coli, and Pseudomonas aeruginosa. Nickel(II) complexes demonstrate slightly lower effectiveness than other treatments, but they always deliver better results than unbound ligands. Cobalt(II) complexes display different activity levels because their effectiveness depends on the particular ligand structure, which distinguishes them from other metals in the series (Pahontu et al., 2013).

The chelation theory introduced by Tweedy in 1964 explains chemical mechanisms through its most cited framework which continues to receive scholarly references in current academic publications. The framework shows that when metal ions interact with ligands their charge decreases because the metal shares some of its charge with donor atoms while their electron density moves throughout the chelate ring structure. The complex's decreased charge together with its higher lipophilicity enables better bacterial cell membrane penetration than the free ion and free ligand. The complex inside the cell can disrupt metal-dependent enzyme function by damaging membrane structures and causing oxidative damage.

This section needs to highlight Thiosemicarbazone Schiff base complexes as important compounds. The thiosemicarbazone arm of these ligands which provides N,N,S coordination enables them to create highly effective antimicrobial metal complexes which appear throughout scientific literature. According to multiple studies copper and nickel thiosemicarbazone complexes display antibacterial properties against methicillin-resistant Staphylococcus aureus (MRSA) at dosages that match or exceed the effectiveness of standard antibiotics which creates an important medical finding because of the worldwide antibiotic resistance emergency (Lobana et al. 2009).

The patterns of antifungal effectiveness show broad similarities with each other. Copper(II) and zinc(II) Schiff base complexes establish their superior antifungal performance against Candida albicans and Aspergillus niger through in vitro testing which exceeds the efficacy of fluconazole and other standard antifungal drugs although their effectiveness in living organisms remains untested. The mechanism likely involves disruption of ergosterol biosynthesis — the fungal cell membrane equivalent of bacterial lipopolysaccharide disruption — though direct evidence for this specific mechanism in metal Schiff base complexes is less developed than the antimicrobial mechanistic literature.

IV. ANTICANCER ACTIVITY

4.1 DNA Binding and Cytotoxicity

The field of transition metal Schiff base literature reaches its maximum expansion when researchers study cancer because this area shows the greatest divide between what researchers discover in laboratories and what actually occurs in medical practice. Researchers have studied the anticancer properties of these complexes through various methods which include testing their ability to kill cancer cells and studying their interaction with DNA and their capacity to block enzymes and their recent research into how they trigger cell death and halt cell division. The most researched method through which transition metal Schiff base complexes fight cancer operates through DNA binding because this approach relies on the earlier established geometric principle. Through their planar aromatic Schiff base complexes square planar copper(II) and nickel(II) complexes with extensive conjugated ligand structures achieve intercalation between the DNA double helix base pairs. The process of intercalation causes the helix to unwind and develop distortions which create obstacles for DNA replication and transcription processes. The result, in rapidly dividing cancer cells, is inhibition of proliferation and induction of apoptosis.

The research used Binding studies through UV-visible absorption spectroscopy and fluorescence quenching and viscometry and circular dichroism to investigate DNA interactions with different complexes. The intrinsic binding constants for calf thymus DNA binding of copper(II) Schiff base complexes control between 10^4 and 10^6 M^{-1} according to established scientific standards which demonstrate high binding power when compared to ethidium bromide which is a standard intercalator (Shiju et al., 2015). The complexes exhibit two distinct binding patterns because they interact with DNA through groove binding and electrostatic forces which connect with the phosphate backbone of the DNA structure.

The research presents extensive cytotoxicity data which demonstrates the effects of treatment on cancer cell lines. The studies have reported IC_{50} values in the low micromolar range which represent the concentration needed to achieve



50% cell growth inhibition for copper and cobalt and nickel Schiff base complexes against HeLa (cervical), MCF-7 (breast), HepG2 (liver), and A549 (lung) cell lines. The research compares these values to cisplatin which serves as the standard platinum anticancer drug and demonstrates that some complexes match or exceed the performance of cisplatin in laboratory tests which has created strong optimism in the scientific community (Gupta & Suresh, 2016).

The positive outlook requires a more measured approach because laboratory tests which assess cell line toxicity show no ability to forecast actual drug effectiveness in live organisms. Numerous compounds which display potential during cell culture experiments exhibit failure in animal testing because of their inadequate pharmacokinetic properties and their unmanageable behavior under normal physiological conditions and their capacity to harm healthy body parts. The field of Schiff base complexes continues to experience this issue because existing research fails to establish strong links between laboratory test results and actual therapeutic outcomes.

4.2 Apoptosis and Cell Cycle Mechanisms

Beyond simple cytotoxicity measurements, researchers have increasingly examined how transition metal Schiff base complexes kill cancer cells at the mechanistic level. Apoptosis induction — the triggering of programmed cell death — is the most desirable mechanism from a therapeutic standpoint, since it is the same pathway that effective anticancer drugs like cisplatin exploit and is generally preferable to necrosis from a toxicity standpoint.

Flow cytometry studies using annexin V-FITC/propidium iodide staining have shown that several copper and cobalt Schiff base complexes induce early and late apoptosis in cancer cell lines at concentrations close to their IC_{50} values, with minimal necrosis at those concentrations. Cell cycle analysis in the same studies often reveals G1 or G2/M phase arrest — exactly the checkpoints where DNA damage signaling would be expected to halt cell division in cells with double-strand breaks caused by intercalation or oxidative damage. These mechanistic findings add biological plausibility to the DNA binding data and suggest that at least some complexes are working through coherent, therapeutically relevant pathways rather than simply being non-specifically cytotoxic.

V. ANTIOXIDANT AND ENZYME INHIBITION ACTIVITY

5.1 Radical Scavenging Properties

Oxidative stress the condition which occurs when reactive oxygen species (ROS) generation exceeds antioxidant protection systems leads to various medical conditions which range from cardiovascular problems and neurodegenerative disorders to cancer and diabetes. The therapeutic potential of free radical scavenging compounds which stop free radical formation explains the common practice of antioxidant testing in studies about Schiff base complex research. Transition metal Schiff base complexes show antioxidant activity through multiple mechanisms. The Schiff base ligand itself — particularly if it contains phenolic hydroxyl groups or aromatic systems capable of stabilizing radical intermediates — can donate hydrogen atoms to quench radicals like the DPPH (2,2-diphenyl-1-picrylhydrazyl) radical used in standard assays. When metal ions bind to the system this process becomes more effective because the metal atoms create extra electron movement which helps stabilize the oxidized ligand radical intermediate. Copper complexes again tend to show the strongest radical scavenging activity among first-row transition metal series which matches copper's established biological function as a superoxide dismutase enzyme component (Nair et al., 2012).

The scientific community disputes whether DPPH and ABTS radical scavenging tests accurately measure biological antioxidant defense mechanisms which creates a valid challenge to most research studies in this area. The chemical tests assess chemical reactivity with permanent stable radicals during conditions that do not simulate normal biological processes. The assessment of antioxidant activity requires research studies which examine both cellular systems and whole organism models but such studies occur less frequently. The antioxidant data presents valuable information which researchers should investigate further but they must treat it as initial screening results instead of proof that the treatment provides effective antioxidant therapeutic benefits.



5.2 Carbonic Anhydrase and Urease Inhibition

Researchers now study how transition metal Schiff base compounds inhibit enzymes because this research area has grown to become a vital part of their investigation into biological activity. Researchers focus on two particular enzymes which are carbonic anhydrase (CA) and urease. The pharmacological importance of carbonic anhydrase inhibition lies in the fact that diuretic and antiglaucoma and anticancer drugs target specific CA isoforms. The study found that various vanadium and copper and zinc Schiff base complexes inhibit carbonic anhydrase activity with IC₅₀ values between nanomolar and low micromolar range, which matches the effectiveness of acetazolamide, a known carbonic anhydrase inhibitor (Akincioglu et al. 2013).

Researchers consider urease inhibition important for developing treatments that target *Helicobacter pylori* infections because this bacterial pathogen requires urease to survive in the acidic environment of the human stomach. Multiple studies have demonstrated that nickel and copper Schiff base complexes exhibit strong urease inhibitory activity, which creates the potential for these compounds to function as dual antimicrobial and enzyme inhibitory agents against this medically significant pathogen.

VI. ANTIDIABETIC ACTIVITY

6.1 Vanadium Complexes and Insulin Mimesis

Research has established the antidiabetic effects of vanadium compounds since 1980 when sodium orthovanadate demonstrated its ability to reduce blood sugar levels in diabetic rats. The issue with basic vanadium salts arises from their high toxicity which leads to serious gastrointestinal problems at doses needed for treatment. The vanadium toxicity issue gets resolved through Schiff base complexation because this method provides three benefits: it keeps the oxidation state constant, it facilitates better oral absorption, and it decreases harmful effects, while most cases show increased insulin-mimetic effects compared to the unbound vanadyl cation.

The oxovanadium(IV) Schiff base complexes display antidiabetic effects in streptozotocin-induced diabetic rat models because they use O₂N and O₂N₂O donor systems to achieve this effect. The process consists of blocking protein tyrosine phosphatase 1B (PTP1B) which functions as an insulin signaling pathway inhibitor. The PTP1B inhibitor blocks PTP1B which increases insulin receptor sensitivity and produces insulin effects on patients who show insulin-resistant conditions (Thompson et al., 2009).

Researchers study Zinc Schiff base complexes for their potential to treat diabetes because zinc functions in the body to maintain insulin storage and insulin release. Research shows that zinc complexes with Schiff base ligands, which scientists derived from both amino acids and salicylaldehyde derivatives, produce glucose-lowering effects in animal studies. However, researchers still understand the exact mechanisms of this compound less than they do for vanadium. The compound operates through various mechanisms that include α -glucosidase inhibition and enhancement of pancreatic β -cell activity.

VII. STRUCTURE-ACTIVITY RELATIONSHIPS: WHAT ACTUALLY DRIVES BIOLOGICAL ACTIVITY

The field generates its most valuable practical knowledge by studying which structural features lead to the most powerful biological activities. The research has developed multiple structure-activity relationships that demonstrate reliable predictive power from existing scientific literature. The studies show that lipophilicity depends on the ligand's aromatic substituents which create a parabolic connection to biological activity. The complex requires specific lipophilicity levels because it needs to maintain membrane crossing ability while avoiding aggregation and uncontrolled distribution into fat compartments. Halogen substituents which include chloro and bromo groups on aromatic rings enhance antimicrobial and cytotoxic activity by increasing lipophilicity and creating electronic effects that stabilize metal-ligand connections. The effects of methyl and methoxy substituents show greater variation.

The metal-to-ligand ratio matters more than is sometimes acknowledged. Mononuclear complexes — one metal per ligand — tend to show different activity profiles than binuclear complexes even when the ligand framework is similar.



Binuclear copper complexes in particular have shown enhanced DNA binding relative to mononuclear analogues, consistent with cooperative binding mechanisms where two metal centers interact simultaneously with the double helix. Figure summarizes the key structural determinants of biological activity across the major activity classes, showing how ligand substituent effects, metal identity, and coordination geometry each contribute to the observed biological profile of a given complex




		Transition Metals						Key Ligand Structural Features for Activity
		Cu	Co	Ni	Zn	V	Mn	
 Biological Activity Scales Antimicrobial Antifungal Anticancer Antioxidant Antidiabetic Enzyme Inhibition	 Antimicrobial				Strong / Well-Documented Activity	Moderate or Variable Activity	Weak or Rarely Reported Activity	Planarity, Planar Lipophilic Rings, Bulky Conjugated systems
	Antifungal		Moderate or Variable Activity					Planarity, Tilted Lipophilic Rings, Bulky Conjugated systems
	Anticancer	Strong / Well-Documented Activity		Moderate or Variable Activity	Moderate or Variable Activity	Weak or Rarely Reported Activity		Planarity, Planar Lipophilic Rings, Bulky Conjugated systems
	Antioxidant							Planarity, Planar Lipophilic Rings, Bulky Conjugated systems
	Antidiabetic				Moderate or Variable Activity	Moderate or Variable Activity		Planarity, lipophilic Ring, Bulky Conjugated systems
	Enzyme Inhibition				Moderate or Variable Activity	Weak or Rarely Reported Activity		Planarity, Planar Lipophilic Ring, Bulky Conjugated systems
 Key Ligand Structural Features for Metal Activity		Planar Coordination Geometry Pi-Conjugation Bulky Lipophilic Groups	Coordination Geometry Pi-Conjugation Bulky Lipophilic Groups	Coordination Geometry Pi-Conjugation Bulky Lipophilic Groups	Coordination Geometry Pi-Conjugation Bulky Lipophilic Groups	Planar Coordination Geometry Pi-Conjugation Bulky Lipophilic Groups	Coordination Geometry Pi-Conjugation Bulky Lipophilic Groups	

Fig: Structural Determinants of Biological Activity in Transition Metal Schiff Base Complexes — A Comparative Summary Across Activity Classes, Source: Author Generated

VIII. CONCLUSION

Transition metal Schiff base complexes occupy an interesting and productive niche in medicinal inorganic chemistry. They combine synthetic accessibility with structural diversity, tunable electronic properties, and a track record of biological activity across multiple pharmacological areas. Antimicrobial applications, particularly against drug-resistant organisms, represent the area where the evidence is strongest and the path to practical application most direct. Anticancer applications are the most actively studied but also the most demanding in terms of the additional mechanistic and in vivo work needed before clinical relevance can be assessed. Antidiabetic activity of vanadium complexes represents perhaps the most mechanistically coherent area, with a clearer understanding of the relevant target — PTP1B — and a body of animal data supporting proof of concept.

The field's next productive decade will probably belong to groups that commit to the harder work: solving complex stability questions under physiological conditions, establishing genuine selectivity indices, conducting careful in vivo pharmacology, and pushing the most promising candidates toward the kinds of preclinical studies that allow serious evaluation of therapeutic potential. The chemistry is good. The biology is interesting. What the field needs now is the patience and resources to find out which of these many promising compounds actually have a future in medicine — and the scientific honesty to accept that many will not, which is exactly how drug discovery is supposed to work.

REFERENCES

- [1]. Akıncıoğlu, A., Akıncıoğlu, H., Göçer, H., Gülçin, İ., & Supuran, C. T. (2013). Novel carbonic anhydrase inhibitors: Bisphosphonate, thioureido indane and Schiff base metal complexes as inhibitors of the cytosolic and membrane-associated isoforms. *Bioorganic and Medicinal Chemistry*, 21(5), 1379–1385. <https://doi.org/10.1016/j.bmc.2012.12.045>
- [2]. Ali, M. A., Mirza, A. H., Butcher, R. J., Tarafder, M. T. H., Keat, T. B., & Ali, A. M. (2008). Biological activity of palladium(II) and platinum(II) complexes of the acetone Schiff bases of S-methyl- and S-



- benzylidithiocarbazate and the X-ray crystal structure of the palladium(II) complex. *Polyhedron*, 27(1), 25–31. <https://doi.org/10.1016/j.poly.2007.08.038>
- [3]. Chandra, S., & Gupta, L. K. (2009). EPR, mass, electronic and IR spectral studies on mononuclear complexes of chromium(III) with macrocyclic ligands derived by condensation of Schiff bases. *Spectrochimica Acta Part A*, 72(3), 750–754. <https://doi.org/10.1016/j.saa.2008.11.020>
 - [4]. Chohan, Z. H., Sumrra, S. H., Youssoufi, M. H., & Hadda, T. B. (2010). Metal based biologically active compounds: Design, synthesis, and antibacterial/antifungal/cytotoxic properties of triazole-derived Schiff bases and their oxovanadium(IV) complexes. *European Journal of Medicinal Chemistry*, 45(7), 2739–2747. <https://doi.org/10.1016/j.ejmech.2010.02.056>
 - [5]. Dutta, S., Bhattacharyya, D., De, A., Banerjee, P., & Pal, S. (2012). Synthesis, characterization and biological activities of copper(II) complexes of Schiff bases derived from amino acids. *Journal of Molecular Structure*, 1007, 154–161. <https://doi.org/10.1016/j.molstruc.2011.10.025>
 - [6]. El-Sawaf, A. K., El-Essawi, M., Nassar, A. A., & West, D. X. (2007). Synthesis, characterization, and biological activity of copper(II) complexes with tridentate Schiff base ligands derived from 2-acetylpyridine. *Journal of Molecular Structure*, 833(1–3), 96–106. <https://doi.org/10.1016/j.molstruc.2006.09.013>
 - [7]. Gupta, K. C., & Suresh, E. (2016). Synthesis, spectral characterization, and biological activity of cobalt(II), nickel(II), copper(II) and zinc(II) complexes with bidentate Schiff base ligands containing pyrazolone ring. *Applied Organometallic Chemistry*, 30(4), 263–271. <https://doi.org/10.1002/aoc.3428>
 - [8]. Keypour, H., Shayesteh, M., Rezaeivala, M., Chalabian, F., Elerman, Y., & Buyukgungor, O. (2012). Synthesis, structural characterization and antimicrobial studies of new Schiff base transition metal complexes derived from condensation of 2-acetylpyridine and 2-amino-5-bromobenzophenone. *Journal of Molecular Structure*, 1027, 83–89. <https://doi.org/10.1016/j.molstruc.2012.05.072>
 - [9]. Lobana, T. S., Sharma, R., Bawa, G., & Khanna, S. (2009). Bonding and structure trends of thiosemicarbazone derivatives of metals — an overview. *Coordination Chemistry Reviews*, 253(7–8), 977–1055. <https://doi.org/10.1016/j.ccr.2008.07.004>
 - [10]. Mendes, I. C., Soares, M. A., dos Santos, R. G., Vilela, C., & Beraldo, H. (2009). Gallium(III) complexes of 2-acetylpyridine-derived thiosemicarbazones: Antimicrobial activity against *Pseudomonas aeruginosa* and *Staphylococcus aureus*. *European Journal of Medicinal Chemistry*, 44(4), 1870–1877. <https://doi.org/10.1016/j.ejmech.2008.06.023>
 - [11]. Nair, M. S., Joseyphus, R. S., & Devi, P. S. (2012). Synthesis, characterization, and biological activity of Cu(II) and Ni(II) complexes of tridentate Schiff bases derived from vanillin. *Journal of Coordination Chemistry*, 65(1), 135–145. <https://doi.org/10.1080/00958972.2011.640397>
 - [12]. Pahontu, E., Fala, V., Gulea, A., Poirier, D., Tapcov, V., & Rosu, T. (2013). Synthesis and characterization of some new Cu(II), Ni(II) and Zn(II) complexes with salicylaldehyde-derived Schiff bases: Antibacterial, antifungal and in vitro antiproliferative activity. *Molecules*, 18(8), 8812–8836. <https://doi.org/10.3390/molecules18088812>
 - [13]. Patil, S. A., Unki, S. N., Kulkarni, A. D., Naik, V. H., & Badami, P. S. (2011). Spectroscopic characterization, in vitro biological studies and fluorescence investigations of Co(II), Ni(II), Cu(II) and Zn(II) complexes with ONNO donor Schiff bases. *Journal of Molecular Structure*, 1005(1–3), 11–21. <https://doi.org/10.1016/j.molstruc.2011.08.055>
 - [14]. Ramesh, G., Daravath, S., Ganji, N., Rambabu, A., Venkateswarlu, K., & Shanker, K. (2020). Synthesis, structural characterization, DNA interaction, antioxidant, antimicrobial, and in vitro cytotoxic studies of Co(II), Ni(II), and Cu(II) complexes of 3-amino-5-(4-fluorophenyl)isoxazole Schiff base derivatives. *Journal of Molecular Structure*, 1206, 127660. <https://doi.org/10.1016/j.molstruc.2019.127660>
 - [15]. Shiju, C., Arish, D., Kumaresan, S., & Bhuvanesh, N. (2015). Synthesis, spectral characterization, DNA binding, nuclease activity, antioxidant and antimicrobial studies of metal complexes derived from 2-



- acetylthiophene-based Schiff base. *Spectrochimica Acta Part A*, 138, 795–806. <https://doi.org/10.1016/j.saa.2014.11.089>
- [16]. Srinivasan, S., Vairavelu, L., & Murugavel, S. (2019). Synthesis, spectral characterization and anticancer study of transition metal complexes of Schiff base ligand derived from amino acid with indole moiety. *Journal of Chemical Sciences*, 131(2), 18. <https://doi.org/10.1007/s12039-019-1587-0>
- [17]. Thompson, K. H., Lichter, J., LeBel, C., Scaife, M. C., McNeill, J. H., & Orvig, C. (2009). Vanadium treatment of type 2 diabetes: A view to the future. *Journal of Inorganic Biochemistry*, 103(4), 554–558. <https://doi.org/10.1016/j.jinorgbio.2008.12.003>
- [18]. Vijayan, P., Viswanathamurthi, P., Silambarasan, V., Velmurugan, D., Butcher, R. J., Pallepogu, R., & Hashimoto, T. (2014). Design and synthesis of new organotellurium Schiff base compounds and their palladium(II), platinum(II) complexes: Structural investigation and in vitro cytotoxic evaluation. *Journal of Organometallic Chemistry*, 759, 1–13. <https://doi.org/10.1016/j.jorganchem.2014.02.012>
- [19]. Zhang, X. M., Tian, H., Guo, A. L., Li, Y. T., Wu, Z. Y., & Yan, C. W. (2017). Biological activity of new ternary copper(II) complexes with Schiff base and nucleobase derivatives: Synthesis, spectroscopic characterization, antioxidant, antimicrobial, and anticancer studies. *Journal of Inorganic Biochemistry*, 173, 8–17. <https://doi.org/10.1016/j.jinorgbio.2017.04.016>

