

Spodias Tuberosa: A Review

Kalyani Chande¹, Shelar Shubhangi², Chavan Ashok², Karande Vaishnavi², Komal Kalnar³

Assistant Professor, Dr. D.Y. Patil College of Pharmacy, Akurdi, Pune, Maharashtra, India¹

Students, Vidya Niketan Institute of Pharmacy and Research Centre Bota, Maharashtra, India²

Student, Matoshri Miratai Aher College of Pharmacy Karjule Harya, Parner, Maharashtra, India³

Abstract: *The review aimed to know the essential aspects of the phytochemical component and pharmacological activities. Based on the ethnobotanical survey states that it can be edible and it exhibits major pharmacological activities which play a key role in drug discovery. Medicinal plant preparations are more demanding due to their fewer side effects and affordable price. Spondiastuberosa is commonly known as imbu. Various phytochemical and pharmacological investigations have been done by various researchers. The phytochemical investigation has been done on different parts of plants. each part of plant has some new components which or not present in other parts of plants. Different extracts of spondiastuberosa has been used by various researchers in order to find the bioactive compounds. Different analytical techniques such as chromatography, spectroscopic, and hyphenated techniques have separated and identified bioactive compounds in spondiastuberosa.*

Keywords: Spondiastuberosa (Anacardiaceae), Anti-inflammatory, Anti-fungal activity, Anti-diabetic activity

I. INTRODUCTION

Natural products which are derived from plant extracts were an abundant source of bioactive compounds. Investigation of natural products for identification of different bioactive compounds which play a key role in novel drug discovery. There is evidence that from post-genomic era more than 80% of the drug substance has been acquired from natural products. [1]. Above 20,000 plant species are used in various traditional medicine around the world and they are reservoirs for the discovery of new drugs [2]. Forest is considered as a storehouse of medicinal plants. Medicinal properties of plants were 1st introduced by dioscorides (40 – 90 AD). Plants have also been used for the treatment of mental disorders and CNS diseases, cardiovascular and respiratory problems, and anticancer activities by plant components [3]. The theory of electronic solutions was developed more in the 20th century. Dissociation of electrolytes by Arrhenius (1859-1927). The concept of PH was published in the year 1909 by Sorensen. Heyrovsky got the Nobel Prize in the year 1959 for the invention of polarography. In 1941 introduction of spectrophotometer in to the market. In the year 1943 IR instruments were introduced, in 1951 self-recording emission spectroscopy. In the year 1953 NMR and gas chromatography was evolved. In the year 1960 atomic absorption spectroscopy was invented. In 1960's hyphenated techniques were invented by linking the chromatographic techniques with spectroscopy (GC-MS).[4]. Advantages and disadvantages of various chromatographic techniques. Gas chromatography was used to find the volatile components in the materials. gas chromatography advantages are linearity is good, it is used in preparative scale because the sample is not destroyed. Disadvantages include low sensitivity, changes occur due to effect of different temperatures, biological samples cannot be examined. HPLC advantages include it requires small sample for the analysis, sample will not disturb during the process. HPLC disadvantages it requires experienced person for the operation. Paper chromatography advantages include the sensitivity (small amount of the compounds which can be detected with routine reagents. Disadvantages include volatile components cannot be separated). Advantages and disadvantages of various spectroscopic techniques such as UV, IR, FTIR, NMR, Mass. Main advantage of UV spectroscopy was it is used for detecting the individual electron transfer between orbitals or atomic, ionic and molecular bands. It helps in the detection of aromatic and multiple bands. Disadvantage of UV highly saturated compounds (single bonds) hydrocarbons (alkanes) and sugars are unable to detect in UV-visible spectroscopy. Advantages of near infrared spectroscopy is samples can be analysed with different modes (e.g., diffuse reflectance, transmittance, transfectance). Disadvantages of this instrument low concentration below 0.1-1 ppm may not be visible. This

technique is not suitable for trace component analysis. Other disadvantage is bands that attached with hydrogen produce intense bands. (Overtone and combination region). Advantages of mid IR include KBr pellet technique was used. nowadays ATR is widely used because it gives the direct measurement solids in their native state. The main disadvantage of MIR is it absorbs more water in the IR region which in turn leads to the hiding of important information due to absorption bands of water. [5]. Applications of analytical techniques in pharmaceutical field are separation and identification of new lead molecules from plants [6]. Calculating the purity of drugs, calculating the impurities that are present in drugs and solvents, development of method validation. Forensic application is drug abuse detection, murder by poison intake, analysis of narcotic intake by sports person [7]. Food analysis application is food authentication, fraud, trade laws, adulteration, contamination, product tampering. [8]

II. TAXONOMIC DESCRIPTION OF SPONDIASTUBEROSA

Common name is imbu. Imbu is deciduous tree with a low, flat topped profusely branched, wide and very dense crown. It grows around 6 meters tall with a very short bole 40-60cm in diameter. It is an edible fruit. [14]

fig 1.15[15]



Kingdom	Planae
Subkingdom	Viridiplantae
Super division	Embryophyte
Division	Tracheophyte
Subdivision	Spermatophytina
Class	Mangnoliopsida
Superorder	Rosanae
Order	Sapindales
Family	anacardiaceae
Genus	Spondias
Species	Spondiastuberosa

Table 1.1[16]

III. PHYTOCHEMICAL INVESTIGATION

Phytochemical analysis by TLC revealed the presence of flavonoids and hydrolysable tannins and HPLC revealed compounds similar to gallic acid and hyperosid hexane extract of spondiastuberosa leaves. 1H NMR showed spectrum showed saturated and unsaturated fatty acids.[17]chlorogenic acid, caffeic acid, rutin and isoquercitrin compounds were identified through HPLC- DAD and HPLC-MS analysis.[18] 3-n-pentadecylphenol (44.1%) from bark and tetratetracontane (38.17 and 28.57%) from leaves and stem was analysed by GC-MS[19] Ultra -performance liquid chromatography coupled to quadruple / time of flight (UPLC-MS-ESI-QTOF) revealed that information about compounds like dehydroascorbic acid, dehydrophaseic acid hexose quinic acid, galloylquinic acid isomer, mangiferin, penta-O-galloylhexoside, caffeoyl-D-glucose, digalloyl glucose may be present based on their molecular weight and retention time [20]. Myricetin-O- rhamoside, anacardic acid compounds has been identified from leaves extract by UPLC-HRMS analysis [21]. Comparison between standard β carotene with different mixtures of pulp of fruits among those mixtures matured pulp flour has high contents of carotenoids [22].

IV. PHARMACOLOGICAL INVESTIGATION

Anti-inflammatory activity:

Anti-inflammatory activity was determined by introducing the inflammatory agents λ -carrageenan(125-500mg/kg) into the right hind paw of mice this activity is compared with response of dexamethasone(2mg/kg). based on the result 500mg/kg of *S. tuberosa* extract showed 63.3% and the response of dexamethasone was 78%. They found that anti-inflammatory effect that produced by *S. tuberosa* was almost equal to the dexamethasone which was used as standard [18]

Anti-fungal activity:

Anti-fungal activity was examined for leaf and root hydroalcoholic extract of *spondiastuberosa* and compared with the standard response of fluconazole. IC50 values for hydroalcoholic extract leaf of *spondiastuberosa* started from 5716.3 μ g/ml to 7805.8 μ g/ml and for the hydroalcoholic extract roots of *spondiastuberosa* 6175.3 to 7805.8 μ g/ml and extracts mixed with fluconazole its inhibitory concentration was values are ranged from 2,65 to 278,41 μ g/ml. Mixed fluconazole concentration showed the activity against *C. albicans*, *C. tropicalis*. root extract showed the activity against *C. tropicalis*. leaf extract showed morphological changes in the two strains [20].

Anti-diabetic activity:

Anti-diabetic activity was proved by injecting the hydroethanolic extract of *spondiastuberosa* (250mg/kg or 500kg/mg) in to the diabetic rats and total cholesterol and HDL, triglycerides, hepatic muscle glycogen, urine, total protein, alanine, glucose levels in blood were analysed. 500kg/mg dosage of the extract reduced the glucose during fasting and post- prandial glucose [23]

Antioxidant activity:

Antioxidant properties of ethyl acetate and methanol extracts of *spondiastuberosa* were evaluated by in vitro methods. ABTS (2,2 Azino-bis- (3 ethylbenzothiazoline)-6 sulfonic acid) method was used to detect the antioxidant activity oxidation reaction was prepared with 7mM ABTS solution +140mM potassium persulfate and this mixture was place at dark 12-16 hours. The ABTS solution was diluted in ethanol until it reaches the absorbance of 0.07 at 743nm. The effect of activity was carried out using aliquots of 30 μ L and mixed with 3 μ L diluted ABTS + solution. The solution was absorbed at frequent interval of time. From this examination it revealed that methanol extract of leaves showed the highest percent oxidation inhibition rate. [24]

V. CONCLUSION

Spondiastuberosa has different phytochemicals constituents like carotenoids, oils, caffeic acid, rutin, etc, hence it has shown many pharmacological activities such as antidiabetic, antifungal, antioxidant, and anti-inflammatory activities.

REFERENCES

- [1] C. Katiyar, A. Gupta, S. Kanjilal, and S. Katiyar, "Drug discovery from plant sources: An integrated approach," vol. 33, no. 1, doi: 10.4103/0974-8520.100295.
- [2] J. B. IlefLimem-Ben Amor 1, "Phytochemistry and biological activities of *Phlomis* species," J Ethnopharmacol, vol. Volume 125, no. 2, pp. 183–202, Sep. 2009.
- [3] "The haven of health Chiefly gathered for the comfort of students, and consequently of all those that have a care of their health, amplified upon five words of Hippocrates, written Epid. 6. Labour, cibus, potio, somnus, Venus. Hereunto is added a preservation from the pestilence, with a short censure of the late sicknes at Oxford. By Thomas Coghnan Master of Arts, and Batcheler of Physicke." <https://quod.lib.umich.edu/cgi/t/text/text-idx?c=eebo;idno=A19070.0001.001> (accessed Oct. 21, 2022).
- [4] M. I. Karayannis and C. E. Efstathiou, "Significant steps in the evolution of analytical chemistry - Is the todayCloseCurlys analytical chemistry only chemistry?" Talanta, vol. 102, pp. 7–15, Dec. 2012, doi: 10.1016/J.TALANTA.2012.06.003.

- [5] J. A. L. N D Lourenco, "bioreactor monitoring with spectroscopy and chemometrics," *ApplMicrobiolBiotechnol*, Apr. 2012.
- [6] M. R. Siddiqui, Z. A. AlOthman, and N. Rahman, "Analytical techniques in pharmaceutical analysis: A review," *Arabian Journal of Chemistry*, vol. 10, pp. S1409–S1421, Feb. 2017, doi: 10.1016/J.ARABJC.2013.04.016.
- [7] "Sci-Hub | Criteria for the identification of compounds by liquid chromatography–mass spectrometry and liquid chromatography–multiple mass spectrometry in forensic toxicology and doping analysis. *AnalyticaChimicaActa*, 492(1-2), 69–82 | 10.1016/s0003-2670(03)00889-4." [https://sci-hub.se/https://doi.org/10.1016/S0003-2670\(03\)00889-4](https://sci-hub.se/https://doi.org/10.1016/S0003-2670(03)00889-4) (accessed Oct. 19, 2022).
- [8] "Sci-Hub | Bioactive Compounds: Historical Perspectives, Opportunities, and Challenges. *Journal of Agricultural and Food Chemistry*, 57(18), 8142–8160 | 10.1021/jf9000132." <https://scihub.se/https://doi.org/10.1021/jf9000132> (accessed Oct. 17, 2022).
- [9] "Atomic Absorption Instrumentation." http://delloyd.50megs.com/MOBILE/atomic_absorption.html (accessed Jan. 07, 2023).
- [10] "IR Spectroscopy - Principle and Instrumentation of Infrared Spectroscopy." <https://byjus.com/chemistry/infrared-spectroscopy/> (accessed Jan. 07, 2023).
- [11] "4 Schematic diagram of the main components of a gas chromatography...Download Scientific Diagram." https://www.researchgate.net/figure/Schematic-diagram-of-the-main-components-of-a-gas-chromatography-system-Philadelphia_fig3_273134301 (accessed Feb. 03, 2023).
- [12] "Schematic presentation of a typical nuclear magnetic resonance...Download Scientific Diagram." https://www.researchgate.net/figure/Schematic-presentation-of-a-typical-nuclear-magnetic-resonance-spectrometer-showing-the_fig1_333657432 (accessed Feb. 03, 2023).
- [13] "Schematic of a GC-MS system | Download Scientific Diagram." https://www.researchgate.net/figure/Schematic-of-a-GC-MS-system_fig1_221927526 (accessed Feb. 03, 2023).
- [14] "Spondiastuberosa Arruda." <https://www.gbif.org/species/8230360> (accessed Feb. 03, 2023).
- [15] M. A. C. de Lima, S. de M. Silva, and V. R. de Oliveira, "Umbu—Spondiastuberosa," *Exotic Fruits*, pp. 427–433, Jan. 2018, doi: 10.1016/B978-0-12-803138-4.00057-5.
- [16] "IT IS Report: Spondiastuberosa." https://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=506463#null (accessed Feb. 03, 2023).
- [17] B. Maria Pereira da Costa Cordeiro et al., "Hexane extract from Spondiastuberosa (Anacardiaceae) leaves has antioxidant activity and is an anti-Candida agent by causing mitochondrial and lysosomal damages", doi: 10.1186/s12906-018-2350-2.
- [18] E. M. da Silva Siqueira et al., "Spondiastuberosa (Anacardiaceae) leaves: profiling phenolic compounds by HPLC-DAD and LC–MS/MS and in vivo anti-inflammatory activity," *Biomedical Chromatography*, vol. 30, no. 10, pp. 1656–1665, Oct. 2016, doi: 10.1002/BMC.3738.
- [19] A. L. Guimarães et al., "Gas chromatography mass spectrometry (GC-MS) analysis of the constituents of the fixed oils obtained from the barks, leaves and stems of Spondiastuberosa Arruda (Anacardiaceae)," *Journal of Medicinal Plants Research*, vol. 12, no. 8, pp. 89–95, Feb. 2018, doi: 10.5897/JMPR2018.6555.
- [20] A. T. Lucas et al., "UPLC-MS-ESI-QTOF Analysis and Antifungal Activity of the Spondiastuberosa Arruda Leaf and Root Hydroalcoholic Extracts," *Antibiotics*, vol. 8, p. 240, 2019, doi: 10.3390/antibiotics8040240.
- [21] G. Jhonyson Arruda Carvalho Guedes Yuri, "Comparative analyses of metabolic fingerprint integrated with cytotoxic activity and in silico approaches of the leaves extract of Spondiasmombin L. and Spondiastuberosa Arr. Cam. from Northeast, Brazil," vol. 40, pp. 26–36, Dec. 2020.
- [22] L. B. Cangussu, P. Fronza, A. S. Franca, L. S. Oliveira, and J. Rocha, "Chemical Characterization and Bioaccessibility Assessment of Bioactive Compounds from Umbu (Spondiastuberosa A.) Fruit Peel and Pulp Flours," 2021, doi: 10.3390/foods10112597.
- [23] H. de Moura Barbosa et al., "Spondiastuberosa inner bark extract exert antidiabetic effects in streptozotocin-induced diabetic rats," *J Ethnopharmacol*, vol. 227, pp. 248–257, Dec. 2018, doi: 10.1016/J.JEP.2018.08.038.

[24] W. F. O. Amanda D. A. Uchôa1, “Antioxidant Activity and Phytochemical Profile of SpondiastuberosaArruda Leaves Extracts,” American journal of plant science, vol. 6, Dec. 2015.