

Use of Artificial Intelligence (AI) to Identify New Species in Zoology: A Review

Prof. Rajesh Kumar Ghosle

Assistant Professor, Department of Zoology

Naveen Govt. Girls College, Mungeli, Chhattisgarh, India

*Corresponding Author E-mail: rajeshghosley2011@gmail.com

Abstract: *The identification and categorization of a new species continue to be essential zoological endeavors that advance our knowledge of biodiversity and guide efforts to conserve it. Demanding and perhaps constrained by subjective biases, conventional taxonomies mostly depend on morphological evaluation and knowledge from experts. Artificial intelligence (AI) has an opportunity to revolutionize species identification procedures by speeding up and improving them. This paper examines the methods, difficulties, and potential future uses of AI in zoology for the identification and classification of novel species. We highlight machine learning, computer vision, and deep learning methods that use environmentally conscious, genetic, and morphological characteristics to automatically identify species. Zoologists can improve accuracy and efficiency and uncover novel aspects of biodiversity by combining AI with conventional taxonomic methods. This is especially important in view of the rapid changes in the environment*

Keywords: Artificial Intelligence (AI), Zoology, Biodiversity, New Species etc

I. INTRODUCTION

The resilience of life on Earth, human well-being, and environmental stability all depend on biodiversity. Many species are still unknown despite tremendous efforts, particularly in poorly understood ecosystems including deep seas, thick forests, and underground environments (Mora et al., 2011). Although traditional taxonomy, which is based on physical traits and genetic analysis, is vital, it frequently requires a significant amount of time and professional expertise. Rapid and scalable identification techniques are required because environmental changes pose a hazard to species survival (Hobern et al., 2019).

To automate, support, and improve species discovery, artificial intelligence (AI), which includes machine learning (ML), deep learning (DL), and computer vision, has become a potent instrument. The acoustic categorization of Australian frogs using improved features and machine learning methods has been covered by Xie et al. (2016). AI systems enable the faster and more accurate identification of new species by analyzing large datasets, identifying trends, and detecting tiny morphological traits that humans cannot (Schneider et al., 2019). The methods, uses, difficulties, and opportunities of AI's revolution in zoological species discovery are examined in this review article.

II. AI METHODOLOGIES IN SPECIES IDENTIFICATION

2.1 Machine Learning and Data-Driven Classification

The machine learning algorithms categorize unidentified points of data by recognizing trends in datasets with labels. Support vector machines (SVM), random forests, and neural networks are examples of approved machine learning (ML) models used in zoology that have been trained using ecological data, genetic sequences, or morphological measures in order to differentiate between species. The process of proposing an identification to a genetic identification of any species can be accelerated by computer vision (Derkarabetian et al., 2022). ML classifiers trained on morphological characteristics, for instance, are able to distinguish minute differences that may be signs of new species.



2.2 Computer vision and Deep Learning

The quick recognition of species from images has been transformed by deep learning, especially convolutional neural networks (CNNs). Artificial intelligence, biosensors, optical sensors, physical sensors, and other technologies are utilized in smart aquaculture systems to gather data in real time. Automation and automated decision-making also play a part (Sharma et al., 2020). In order to accurately identify species from photos or videos, CNNs automatically extract hierarchical characteristics from pictures. This feature is particularly useful for field research, where large visual datasets are produced via camera traps and drone photography.

2.3 Genomic Data Analysis

AI techniques are increasingly applied to genetic data, such as DNA barcodes and whole-genome sequences, to delineate species boundaries. Clustering algorithms and neural networks can detect genetic divergence indicative of speciation events. AI-based clustering approach suggests that endemic species (i.e. fish) conservation should focus on alleviation of low flows, control of species introduction, limiting generalist expansion, and enhancing the hydrological connectivity fragmented by dams (Cheng et al., 2018). Integrating morphological and genetic data through AI models enhances reliability in discovering cryptic fish species.

III. AI APPLICATIONS IN THE SEARCH FOR NEW SPECIES

3.1 Automated Image Identification in Surveys of Biodiversity

Rapid analysis of photos taken by drones, camera traps, or field photography is made possible by AI-powered image identification systems. In order to identify bird species from photos, Schneider et al. (2019) created a CNN-based approach that was successful in identifying morphological variations that were previously unknown but suggested the existence of new species. And lastly, five machine learning algorithms—K-nearest neighbor, support vector machines, random forest, artificial neural networks, and linear discriminant analysis—are used to compare various augmented visual representations of features. The findings of the experiment indicate that, when compared to alternative approaches, our suggested visual representation of features might produce superior classification performance for twenty-four frog species (Xie et al., 2016).

3.2 Species Delimitation and Genetic Information

The discovery of species boundaries has been rendered easier by AI algorithms applied to genomic databases. By using machine learning techniques to examine genetic divergence between fish populations, Sharma et al. (2021) were able to uncover cryptic species that are not discernible by morphology alone. These genetic AI technologies speed up the process of validating novel species, particularly in morphologically related groupings.

3.3 Integration of Environmental DNA (eDNA) with AI

Genetic material in environmental samples can be used to determine the existence of species through environmental DNA sampling. Even for uncommon or elusive species, AI algorithms may detect species signatures by analyzing massive eDNA datasets. In distant settings, this method has resulted in the discovery of novel aquatic species.

3.4 Automated Acoustic Monitoring

A further method for finding new species is bioacoustics, which records animal noises and cries. In order to differentiate between species based on vocalizations, AI systems examine acoustic data. By identifying distinct call patterns, for instance, models based on machine learning and deep learning have discovered new frog species (Xie et al., 2016).

IV. CHALLENGES AND LIMITATIONS

Despite promising advances, applying AI for new species discovery faces several challenges:

- **Data Quantity and Quality:** For the development of precise AI models, high-quality, annotated datasets are necessary. Model creation for uncommon or cryptic species is hampered by a lack of data.



The ability to apply the predictions of AI models to new species may be diminished due to bias and overfitting to training data.

- **Taxonomic Validation:** Expert validation is necessary for AI predictions, and differences between AI and conventional taxonomy may spark discussions on species status.
- **Issues of Ethics and Conservation:** Rapid AI-driven identification has been combined with conservation tactics to avoid environmental degradation or illicit use.

V. FUTURE PERSPECTIVES

Combining AI with other technologies has the potential to improve species discovery even more:

- **Multimodal Data Integration:** By integrating ecological, morphological, genetic, and auditory data into AI platforms, identification of species precision can be increased.
- **AI and Citizen Science:** By combining AI analysis with public participation in data collecting, biodiversity monitoring initiatives may be expanded.
- **Automated Taxonomy:** While human monitoring is still essential, the development of AI systems that can provide taxonomic categorization suggestions on their own might speed up the discovery process.
- **Global Biodiversity Monitoring:** Real-time monitoring of biodiversity hotspots can be facilitated by AI-enabled systems, which can help with the quick identification of new species and modifications to the environment.

VI. CONCLUSION

Zoological taxonomy and biodiversity discoveries might be revolutionized by artificial intelligence. AI speeds up the discovery of new species, particularly in poorly understood or inaccessible areas, using sophisticated picture recognition, genetic analysis, auditory monitoring, and environmental DNA interpretation. Even though there are still issues with data quality, validation, and ethics, further technical advancements and multidisciplinary partnerships should make artificial intelligence (AI) a vital tool in zoology. In the face of fast environmental change, utilizing AI's potential can result in a more thorough knowledge and protection of Earth's biodiversity.

ACKNOWLEDGEMENTS

Author is greatly acknowledged to our institution Naveen Govt. Girls College Mungeli (C.G.), for valuable support.

REFERENCES

- [1]. Mora, Camilo, Derek P. Tittensor, Sina Adl, Alastair GB Simpson, and Boris Worm. "How many species are there on Earth and in the ocean?." *PLoS biology* 9, no. 8 (2011): e1001127.
- [2]. Hobern, Donald, Brigitte Baptiste, Kyle Copas, Robert Guralnick, Andrea Hahn, Edwin Van Huis, Eun-Shik Kim et al. "Connecting data and expertise: a new alliance for biodiversity knowledge." *Biodiversity data journal* 7 (2019): e33679.
- [3]. Schneider, Stefan, Graham W. Taylor, Stefan Linquist, and Stefan C. Kremer. "Past, present and future approaches using computer vision for animal re-identification from camera trap data." *Methods in Ecology and Evolution* 10, no. 4 (2019): 461-470.
- [4]. Derkarabetian, Shahan, James Starrett, and Marshal Hedin. "Using natural history to guide supervised machine learning for cryptic species delimitation with genetic data." *Frontiers in Zoology* 19, no. 1 (2022): 8.
- [5]. Sharma, Dolly, and Ranjit Kumar. "Smart aquaculture: integration of sensors, biosensors, and artificial intelligence." In *Biosensors in Agriculture: Recent Trends and Future Perspectives*, pp. 455-464. Cham: Springer International Publishing, 2021.
- [6]. Cheng, Su-Ting, Wen-Ping Tsai, Tzu-Chun Yu, Edwin E. Herricks, and Fi-John Chang. "Signals of stream fish homogenization revealed by AI-based clusters." *Scientific reports* 8, no. 1 (2018): 15960.
- [7]. Xie, Jie, Michael Towsey, Jinglan Zhang, and Paul Roe. "Acoustic classification of Australian frogs based on enhanced features and machine learning algorithms." *Applied Acoustics* 113 (2016): 193-201.

