

Clinical and Conceptual Study of Sutika Vishada (Postpartum Depression) and its Management Protocols through Ayurveda

Dr. Rajgauree Appasaheb Mengawade¹ and Dr. Limbanappa Garthe²

¹P. G. Scholar. Dept of Streeroga and Prasutitantra

²HOD, Dept of Streeroga and Prasutitantra

Matoshree Asarabai Darade Ayurved College, Babhulgaon, Yeola

Abstract: *Background: The postpartum period (Sutika Kala) represents a significant physical, emotional and social change. In some cases, mothers experience ongoing sadness, anxiety, tiredness, a loss of self-confidence, disrupted sleep and problems with bonding to the infant. If these symptoms persist and are of a clinical significance, they can be diagnosed as postpartum depression (PPD). In Ayurveda, great emphasis is placed on looking after and restoring the mother during Sutika Kala.*

Objective: The aim is to examine the symptoms of postpartum depression using the Ayurvedic concept of Sutika Vishada and to consider various supportive measures such as Sutika Paricharya, Satvavajaya Chikitsa, certain external therapies, appropriate nourishment and the use of Medhya Rasayana.

Materials and Methods: The conceptual review was created on the basis of the submitted manuscript and the classical and more recent literature that it references. The Ayurvedic concepts concerning Sutika Kala, Vata, Agni, Dhatu, Manas, Manovaha Srotas and Satvavajaya were arranged together with the modern factors such as hormonal transition, sleep disruption, psychosocial stress and the gut-brain relationship.

Results: It can be suggested that the exertion associated with childbirth, loss of tissue, insufficient rest, disruption of diet and psychological stress may all lead to Agnimandya and Dhatu Kshaya, after which there is an aggravation of Vata and a disturbance of mental function. Sutika Paricharya offers a preventive and supportive approach through adequate nourishment, rest, emotional support and by following the Satvavajaya principles. Although some Ayurvedic medicines may have potential, individual clinical supervision is necessary regarding their use after childbirth and while breastfeeding.

Conclusion: In conclusion, Ayurveda provides a holistic and preventative approach to a woman's emotional well-being after giving birth; although its supportive principles could supplement modern care for maternal mental health, clinically significant or severe depression still needs proper professional evaluation and treatment.

Keywords: Sutika Vishada, Postpartum Depression, Sutika Kala, Sutika Paricharya, Satvavajaya Chikitsa, Manovaha Srotas, Medhya Rasayana, maternal mental health

I. INTRODUCTION

Becoming a mother is marked by major biological, psychological and social changes. While the mother is recovering from childbirth, she is at the same time getting used to feeding, to interrupted sleep, to physical discomfort and to her new responsibilities. In some cases this period is associated with ongoing emotional distress. Postpartum depression is not merely ordinary tiredness or a short phase of emotional sensitivity. It may include a persisting low mood, a loss of



interest, excessive guilt, hopelessness, fatigue, anxiety, disturbances of sleep or appetite, impaired concentration and difficulty in forming a comfortable relationship between mother and infant.

The current understanding is that postpartum depression is due to a number of factors. The sudden changes in reproductive hormones, the process of physical recovery, sleep deprivation, pain, difficulties with feeding, financial stress, problems in the relationship, and a lack of adequate social support can all combine with any previous psychological vulnerability. Hence, the condition must take into account both biological and psychosocial factors.

The postpartum period in Ayurveda is known as Sutika Kala and is regarded as a stage during which the mother needs special care and recovery. The traditional principles concerning the postpartum period stress nourishment, support for the digestive system, rest and the avoidance of overexertion. Although Ayurveda does not have an exact equivalent to the modern psychiatric diagnosis of PPD, its symptoms can be related in concept to Vishada (mental disturbance), Vata aggravation, depletion of the Dhatu and a disturbance of the Manovaha Srotas.

This review makes use of Sutika Vishada as a conceptual link rather than asserting that the two systems define the same diagnosis. It is still essential to carry out modern clinical assessment whenever depression is suspected.

II. AIM AND OBJECTIVES

Objective: To gain a clear conceptual understanding of postpartum depression from the Ayurvedic viewpoint of Sutika Vishada and to outline the principles relating to supportive management during the postpartum period.

Objectives: (1) To explain the physical, psychological and social factors that are associated with postpartum depressive symptoms; (2) to describe the Ayurvedic ideas that are relevant to Sutika Kala; (3) to set out a conceptual Samprapti of Sutika Vishada; (4) to review Sutika Paricharya and Satvavajaya Chikitsa; and (5) to determine the areas in which Ayurvedic supportive care and modern maternal mental-health services might complement each other.

III. MATERIALS AND METHODS

The article is a conceptual review which is mainly based on the source manuscript. The source material includes a review of the classical Ayurvedic literature such as the Brihatrayi, Laghutrayi and Kashyapa Samhita, as well as more recent literature obtained through searches on PubMed, Google Scholar and DHARA up to June 2026. The following concepts were examined: Sutika Paricharya, Sutika Unmada, maternal mental health, Medhya Rasayana and Shatavari.

In this version the material has been restructured and rewritten in original prose, keeping the key Ayurvedic terminology. The article does not include a new clinical trial and does not assert that traditional medicines have been proven to be effective; herbal preparations and treatments are stated to be supportive in nature or still under investigation and should therefore be chosen by qualified clinicians, in particular while breastfeeding.

IV. POSTPARTUM DEPRESSION AND SUTIKA KALA

A woman in the period after giving birth might have to deal at the same time with physical exhaustion, the need to adjust emotionally and major changes in her daily routine. Contemporary theories emphasise the hormonal shift, changed stress responses, loss of sleep and psychosocial pressures. These factors can exacerbate each other: lack of sleep can lead to greater fatigue and increased emotional reactivity, while exhaustion can make it more difficult to feed and look after the infant and can in turn increase distress.

In Ayurvedic terms, Vata is given particular significance after childbirth. Traditional views link increased Vata with the exertion, loss of fluids and tissues, and the changes that take place following delivery. Weakness and instability may become more noticeable if nourishment and rest are not sufficient. Emotional stress can also disrupt mental equilibrium.

Sutika Paricharya's preventive value is therefore significant. As part of the early postpartum period one can provide nourishing food, ensure adequate rest, establish a gentle routine, give emotional reassurance, offer practical support from the family and keep a close watch on any changes in mood and behaviour.



V. ETIOLOGICAL FACTORS (NIDANA)

The source manuscript classifies various factors under three categories: the Aharaja, the Viharaja, and the Manasika pathways. The factors belonging to the Aharaja category are those that involve inadequate or unsuitable food during a time when the digestive system is vulnerable. Those which fall within the Viharaja category consist of excessive physical activity, disturbed sleep, exposure to cold, and a failure to follow an appropriate postpartum routine. As for the Manasika factors, they are fear, emotional trauma, isolation, unmet expectations, interpersonal difficulties, and a feeling of being neglected.

They should be seen as factors that interact rather than as separate causes. For instance, poor sleep can lead to fatigue and anxiety; fatigue in turn can lower a woman's confidence when looking after her infant; and diminished confidence can then cause greater emotional distress. It follows that a thorough approach to the postpartum period must include physical, psychological and social support.

VI. CONCEPTUAL SAMPRAPTI OF SUTIKA VISHADA

A conceptual sequence might be explained in this way: the physical effort and loss involved in childbirth, together with dietary, behavioural or psychological stressors, may lead to Agnimandya and Dhatu Kshaya. This condition is linked with an aggravation of Vata. If psychological stress is added, then Rajas and Tamas may affect mental function. The resulting disturbance can then be examined in connection with the Manovaha Srotas and may appear as Vishada or other mental disturbances after childbirth.




Figure 1. The conceptual Samprapti pathway of Sutika Vishada.

VII. CLINICAL FEATURES

The features reported are persistent sadness, frequent crying, a feeling of helplessness, marked fatigue, lethargy, disturbed or fragmented sleep, excessive worry, guilt, self-doubt, reduced self-confidence and difficulty in forming a bond with mother and infant. Since normal postpartum tiredness can be similar to some of these symptoms, it is necessary to take into account the persistence, severity and presence of functional impairment.

Any symptoms that are severe, last for a long time or greatly disrupt everyday activities need to be evaluated by a professional. In the case of thoughts about self-harm or suicide, thoughts of harming the baby, severe confusion, hallucinations or a loss of contact with reality, urgent medical attention is required.



SYMPTOMS OF POSTPARTUM DEPRESSION



VIII. AYURVEDIC MANAGEMENT FRAMEWORK

1. Sutika Paricharya and dietary support: The food introduced after childbirth should be nourishing, warm, easy to digest and suited to the mother's tolerance. The idea is to carry out a gradual restoration rather than impose unnecessary restrictions.
2. With regard to external supportive therapies, Abhyanga is traditionally employed for the purpose of providing comfort and for helping to create a sense of grounding; other procedures like Shirodhara may be considered by trained practitioners following an individual assessment. The suggested hormonal or neurotransmitter effects still need more solid clinical evidence.
3. Regarding Medhya Rasayana and the use of herbs: The text mentions Shatavari, Ashwagandha and Saraswatarishta. The fact that these have a traditional use does not in itself prove that they are safe to take while breastfeeding. Whenever a medicinal preparation is being used, it should be chosen and kept under the supervision of a qualified clinician.
4. Satvavajaya Chikitsa: This can be regarded as comprising supportive psychological care such as offering reassurance, promoting positive modes of thinking, providing emotional support and helping to strengthen adaptive coping mechanisms. When it is appropriate from a medical point of view, gentle yoga, breathing exercises and relaxation techniques may be used as a complement to the care, but should not take the place of professional treatment for clinically significant depression.



IX. AYURVEDIC–MODERN CORRELATION

The Ayurvedic approach and the current one are based on different systems of explanation and must not be regarded as scientifically identical. Yet it is possible to find clinically relevant areas of overlap. For example, Agnimandya and digestive disturbance can be looked at in conjunction with the modern interest in the gut-brain axis, and Vata-related depletion can be considered in relation to the modern descriptions of hormonal transition, sleep loss and changes in the stress systems.

The practical agreement is more substantial than the theoretical equivalence since adequate nutrition, protection from sleep disruption, a reduction of stress, social support and the restoration of physical strength are all relevant in both approaches.

X. DISCUSSION

Postpartum depression is due to a number of factors and therefore no one intervention will be suitable for all mothers; particularly relevant is the Ayurvedic focus on prevention since the routines needed after giving birth can be put in place before the symptoms become serious.

The principles that focus on Vata stress the importance of resting, maintaining warmth, eating properly and avoiding overexertion; those concerning Manas, Rajas and Tamas place emphasis on the emotional atmosphere, while Satvavajaya stresses the need for supportive counselling and assistance in coping. In practice, supportive care can result from family involvement, protected sleep, a reduced workload, adequate nutrition and mental-health screening.

Postpartum depression may also become severe and even life-threatening. Care according to Ayurvedic principles should

XI. LIMITS AND FUTURE POSSIBILITIES

It is not a new clinical trial but a conceptual review; the evidence cited in the source consists of classical references, case reports and therapeutic literature, so stronger clinical research is needed before any Ayurvedic interventions can be recommended as established treatments.

Future research should make use of standard diagnostic criteria and validated tools such as the Edinburgh Postnatal Depression Scale, clearly report on breastfeeding status and any adverse events, define the dose and the length of treatment, and compare Ayurvedic supportive care with suitable standard care; patient acceptability, family support, sleep and quality of life should also be assessed.

XII. CONCLUSION

Sutika Kala is a period when physical exhaustion, emotional stress, disrupted sleep and significant changes to one's way of life can all come together, and the Ayurvedic idea of Sutika Vishada offers a culturally relevant framework by means of the linked concepts of Vata, Agni, Dhatu, Manas and Manovaha Srotas.

A holistic postpartum-care model could include nourishing food, adequate rest, emotional reassurance, kind supportive measures, and the Satvavajaya principles; however, although some Ayurvedic medicines may show promise, their use—especially while breastfeeding—must be under the supervision of a qualified professional and be supported by more evidence.

The best course for the future is an integrative one, since traditional knowledge gained from postpartum care can provide preventive and supportive measures while modern methods of screening, diagnosis and treatment are still available to those mothers who need them.

Table 1. Framework for the supportive ayurvedic management

Domain Core focus

Rest, adequate nourishment, a slow recovery and protection against overexertion.

Therapies such as Abhyanga may be carried out with external assistance under the guidance of trained professionals.



Support provided internally involves traditional formulations based on an individual assessment, in particular while breastfeeding.

Satvavajaya offers reassurance, counselling, adaptive coping strategies and gentle mind-body practices.

Figure 2. Four-part supportive management framework.

REFERENCES

1. Kashyapa Samhita, Khila Sthana, Chapter 11, Verse 3.
2. Sushruta Samhita, Uttarantra, Chapter 62, Verse 2.
3. the World Health Organization's Guidelines on Maternal Mental Health.
- A case report on the ayurvedic treatment of postpartum depression. International Ayurvedic Medical Journal, 2025.
5. The phytochemical aspects and its effect on the neuro-gut axis of Asparagus racemosus (Shatavari) with regard to postpartum depression. PMC, 2025.
6. Sutika Paricharya's role concerning postpartum depression. World Journal of Pharmaceutical Research, 2024

