

The Role of Cognitive Behavioural Therapy in the Prevention of Attempted Suicide: Mechanisms, Empirical Evidence and Clinical Interventions

Dr. Ajay Gauri

Associate Professor, HOD, Deptt. of Psychology
A.R. Govt. P.G. College, Ambala Cantt. (Haryana)

Abstract: *Suicide represents a global public health crisis which is accounting for hundreds of thousands of deaths annually with non-fatal suicide attempts occurring at a significantly higher frequency. Individuals with a history of suicide attempts represent the population at highest risk for future self-harm and deadly completion. Traditional psychiatric interventions often focus on underlying diagnostic categories e.g., Major Depressive Disorder, treating suicidal ideation and behaviour merely as secondary symptoms. In contrast, suicide-focused Cognitive Behavioral Therapy (CBT) specifically Cognitive Therapy for Suicide Prevention (CT-SP) and Dialectical Behavior Therapy (DBT) targets the suicidal mode directly as a primary clinical focus. This paper examines the theoretical framework, specific clinical protocols, neurocognitive mechanisms, empirical evidence base and implementation considerations of CBT in preventing attempted suicide. By addressing core cognitive distortions, emotional dysregulation, implicit suicide associations and deficient problem-solving capacities, CBT equips high-risk individuals with functional coping strategies, demonstrably reducing repeated suicide attempts across diverse clinical trials.*

Keywords: Cognitive Behavioral Therapy, Suicide Prevention, Suicide Attempt, Cognitive Therapy for Suicide Prevention, Dialectical Behavior Therapy, Suicidal Mode, Safety Planning.

I. INTRODUCTION


Suicidal behaviour constitutes one of the most pressing challenges in contemporary psychiatry, psychology and public health. According to World Health Organization, suicide accounts for approximately 700,000 deaths annually worldwide with an estimated 20 non-fatal attempts for every recorded completion. A previous suicide attempt remains the single strongest predictor of future completed suicide across clinical populations. Consequently, identifying, refining and implementing targeted psychotherapeutic interferences specifically engineered to disturb the pathway from suicidal ideation to overt action is a clinical priority.

Historically, clinical paradigms treated suicidal behaviour as a downstream symptom of primary axis-I psychiatric diagnoses such as Major Depressive Disorder, Bipolar Disorder, Schizophrenia or Borderline Personality Disorder. Under this traditional medical model, it was assumed that successfully treating the underlying disorder would automatically resolve suicidal ideation and eliminate self-harm risk. However, epidemiological and clinical data have revealed that while depression treatment may improve mood, it does not reliably eliminate suicidal behaviour, particularly during acute crises or post-discharge transitions.

Over the past three decades, psychological research has established that suicidal behaviour purposes as a distinct and idiographic clinical entity driven by unique cognitive, emotional and behavioural processes. The emergence of cognitive-behavioural paradigms explicitly designed for suicide prevention most notably Cognitive Therapy for Suicide Prevention (CT-SP) developed by Aaron Beck, Gregory Brown and colleagues and Dialectical Behavior Therapy (DBT) developed by Marsha Linehan has revolutionized suicide risk management. This paper examines how Cognitive



Behavioral Therapy (CBT) operates as an evidence-based intervention to prevent repeated suicide attempts by examining its theoretical constructs, clinical mechanics, empirical backing, and operational implementation.



II. THEORETICAL FRAMEWORKS OF CBT IN SUICIDE PREVENTION

To understand how CBT prevents suicide attempts, one must examine the cognitive models that describe the transition from internal psychological pain to external self-directed violence.

The Suicidal Mode and Beck's Cognitive Model

Beck (1996) expanded classic cognitive theory to articulate the concept of the "**suicidal mode**." A mode is an integrated network of cognitive schemas, affective states, physiological drives and behavioural tendencies designed to process environmental demands and guide action. In high-risk individuals, the suicidal mode remains inactive until triggered by acute stress, perceived burdensomeness, severe interpersonal rejection or physical pain. When activated, the suicidal mode shifts cognitive processing into a constrained, hyper-focused state characterized by three following components:

1. **Cognitive Schema Activation:** Pervasive beliefs regarding helplessness ("I cannot change my situation"), hopelessness ("Things will never get better") and unlovability ("I am a burden to everyone").
2. **Affective Distress:** Intolerable psychological pain (often termed *psychache* by Shneidman, 1993) combined with intense shame, anxiety or emotional numbness.
3. **Behavioural Imperative:** Viewing suicide not as a tragedy but as the *only accessible or logical solution* to terminate intolerable suffering.

The primary objective of CBT is not simply to relieve general distress but to decommission the suicidal mode, deactivate its triggers and build adaptive cognitive-behavioural pathways that prevent the translation of suicidal ideation into an actual attempt.

The Interpersonal Theory of Suicide (ITS)

Joiner's (2005) Interpersonal Theory of Suicide provides a complementary framework that CBT protocols frequently incorporate. ITS asserts that a fatal or near-fatal suicide attempt occurs only when an individual possesses both the desire for suicide and the acquired capability to engage in lethal self-harm.

- **Perceived Burdensomeness:** The cognitive calculation that one's existence imposes a net positive trench on family, friends or society ("My death is worth more than my life").
- **Thwarted Belongingness:** The painful emotional state of social isolation, alienating loneliness and lack of mutual care.
- **Acquired Capability:** The adaptation to physical pain and fear of death, developed through previous self-harm, exposure to violence or bodily trauma. CBT directly addresses the subjective drivers of suicidal desire by restructuring distorted beliefs regarding burdensomeness and building actionable interpersonal skills to remedy thwarted belongingness, effectively neutralizing the cognitive engine driving suicidal intent.

The Fluid Vulnerability Model

Rudd (2006) introduced the Fluid Vulnerability Model (FVM), which conceptualizes suicide risk as a dynamic interaction between baseline vulnerability (trait-level factors such as history of trauma, poor impulse control and early



attachment ruptures) and acute risk states (state-level factors like recent job loss, acute sleep deprivation or severe relationship dissolution). CBT aligns with the FVM by distinguishing between long-term skill acquisition to reduce baseline vulnerability and rapid-response safety strategies e.g., Safety Planning Interventions to stabilize acute flare-ups.

III. CORE MECHANISMS OF CBT IN SUICIDE ATTEMPT PREVENTION

CBT systematically alters the cognitive and behavioural architecture that leads to self-directed violence. The specific healing mechanisms that mitigate suicide attempt risk include cognitive restructuring, distress tolerance, emotional regulation and problem-solving training.

Restructuring Cognitive Constriction and Tunnel Vision

During an acute suicidal crisis and individuals experience severe cognitive constriction (commonly described as "tunnel vision"). The brain's executive functioning networks struggle to generate alternative outcomes, viewing the world through binary lenses: total relief via death or perpetual agony via life (Shneidman, 1993). CBT interventions target cognitive distortions that accelerate this constriction:

Cognitive Distortion	Clinical Manifestation in Suicidal Crisis	CBT Restructuring Mechanism
Catastrophizing	"If my partner leaves, my life is completely destroyed forever."	Decatastrophizing; examining worst-case vs. realistic outcomes.
Overgeneralization	"I failed this task, proving I fail at every single thing I attempt."	Gathering empirical evidence of past successes and competence.
Emotional Reasoning	"I feel like a useless burden, so I must truly be a burden."	Separating subjective emotional states from objective situational facts.
Dichotomous Thinking	"Either I fix this problem perfectly right now, or I must end my life."	Identifying continuum states and generating intermediate options.

Through Socratic questioning and guided discovery, the clinician helps the patient identify these automated thoughts, challenge their validity and construct alternative, balanced cognitive responses. Over time, this weakens the automaticity of the suicidal mode.

Enhancing Problem-Solving Capacity

Deficits in relational and task-oriented problem-solving skills are strongly correlated with suicide attempts (Neazu & Asarnow, 2013). When faced with acute life stressors, suicidal individuals often exhibit problem-solving deficits characterized by passivity, impulsivity, cognitive rigidity and an inability to anticipate long-term consequences. CBT incorporates structured Problem-Solving Therapy (PST) and provides following course to teaching patients which is a systematic step-by-step algorithm as follows:

1. **Problem Definition:** Articulating the stressor accurately without emotional over-interpretation.
2. **Goal Generation:** Defining realistic and achievable short-term objectives.
3. **Brainstorming:** Generating multiple potential solutions without immediate criticism.
4. **Decision Analysis:** Evaluating pros and cons of each generated option.
5. **Implementation & Verification:** Executing the chosen plan and reviewing the outcome.

By transforming overwhelming, amorphous crises into discrete and manageable problems, CBT restores an individual's sense of agency, reducing the perceived necessity of suicide as an escape mechanism.

Emotional Regulation and Distress Patience

Severe emotional dysregulation is a central feature preceding a suicide attempt. When emotional intensity exceeds an individual's internal handling threshold, self-harm or suicide is often selected as a rapid and despairing attempt at emotional down-regulation (Linehan, 1993). Within Dialectical Behaviour Therapy (DBT), a specialized form of CBT, distress patience skills which are explicitly taught to help individuals to survive acute crises without making the situation worse. This is as follows:



- **TIPP Skills:** Utilizing physiological interventions namely, Temperature (cold water facial immersion to stimulate the mammalian dive reflex and slow heart rate), Intense exercise, Paced breathing and Paired muscle relaxation to lower hyper-arousal rapidly.
- **Radical Acceptance:** Cultivating psychological acceptance of painful current realities that cannot be immediately altered and thereby reducing the secondary distress generated by fighting reality.
- **Mindfulness of Current Emotions:** Observing intense emotional rushes without judgment or immediate behavioural reaction by allowing the emotional wave to peak and subside naturally.

Evidenced-Based CBT Protocols for Suicide Prevention

Multiple specialized CBT protocols have been developed and rigorously tested specifically for preventing attempted suicide. The two most established paradigms are Cognitive Therapy for Suicide Prevention (CT-SP) and Dialectical Behavior Therapy (DBT).

Cognitive Therapy for Suicide Prevention (CT-SP)

Developed by Brown, Beck and colleagues (2005), CT-SP is a targeted and short-term treatment protocol typically spanning 10 to 12 individual sessions. Unlike general cognitive therapy, every session of CT-SP explicitly targets the cognitive-behavioural factors directly linked to suicidal behaviour. The CT-SP protocol is divided into three distinct operational phases:

Phase 1: Assessment, Safety Planning, and Narrative (Sessions 1–3)

1. **The Suicidal Narrative:** The clinician conducts a detailed behavioural chain analysis of the most recent suicide attempt or acute crisis for mapping out the precise timeline of environmental triggers, automatic thoughts, emotional surges and physical actions.
2. **Stanley-Brown Safety Planning Intervention (SPI):** A prioritized and written list of personalized coping strategies and resources created collaboratively with the patient.
3. **Lethal Means Restriction:** Systematic identification and removal/securing of lethal means (e.g., firearms, prescription medications and high-location access) from the patient's immediate environment which is a crucial intervention shown to reduce suicide fatalities independently (Yip et al., 2012).

Phase 2: Skill Acquisition and Cognitive Restructuring (Sessions 4–9)

1. **Coping Cards:** Creating handheld cards containing key adaptive thoughts, cognitive reframes or action steps to utilize when suicidal ideation escalates.
2. **Behavioural Activation:** Re-engaging in pleasant or mastery-oriented activities to restore positive emotional reinforcement and diminish social isolation.
3. **Cognitive Reframing:** Directly challenging core schemas of hopelessness and burdensomeness using empirical testing methods.

Phase 3: Relapse Prevention and Consolidation (Sessions 10–12)

The Relapse Prevention Task: A crucial and highly specific behavioural test wherein the patient is guided through an intense mental simulation of a hypothetical future crisis. The patient must mentally walk through the intensifying stressor, recognize activated suicidal thoughts and successfully apply their safety plan and cognitive skills in real-time to resolve the crisis without attempting suicide.

Dialectical Behaviour Therapy (DBT)

Developed by Marsha Linehan (1993), DBT was explicitly created to treat chronically suicidal individuals diagnosed with Borderline Personality Disorder (BPD). DBT synthesizes standard cognitive-behavioural techniques with Eastern mindfulness principles and dialectical philosophy. Comprehensive DBT consists of four standard treatment components:

Individual Therapy: Weekly 50-minute sessions focused on applying skills to specific life events, structured around a hierarchically ordered Diary Card where life-threatening behaviors receive top priority.



Group Skills Training: Weekly 2-hour psychoeducational sessions covering Mindfulness, Interpersonal Effectiveness, Emotion Regulation, and Distress Tolerance.

In-the-Moment Phone Coaching: Access to phone coaching between sessions to help generalize skill use to real-world acute crises before engaging in self-harm.

Consultation Team: Weekly meetings for therapists to maintain fidelity, prevent burn-out, and manage the high emotional demands of working with high-risk populations.

Brief Cognitive Behavioral Therapy for Suicide Prevention (BCBT-SP)

Rudd et al. (2015) developed BCBT-SP, an adapted short-term variation (12 sessions) optimized for acute care settings and military populations. BCBT-SP concentrates heavily on crisis management, emotion regulation and cognitive flexibilities by making it particularly viable for high-volume psychiatric emergency contexts.

IV. STRUCTURAL INTERVENTIONS: THE SAFETY PLANNING INTERVENTION (SPI)

A core contribution of CBT to suicide prevention is the Stanley-Brown Safety Planning Intervention (SPI) (Stanley & Brown, 2012). Often integrated into CBT protocols, the SPI is a brief and step-by-step clinical intervention that produces a practical and prioritized list of coping strategies and support systems.

Unlike traditional "no-suicide contracts", which carry no empirical evidence base and can dangerously substitute for real therapeutic alliance which is the Safety Plan is an active and skill-based behavioural instrument.

STEP 1: Warning Signs

- Recognizing internal thoughts, images, moods or behaviours signaling a crisis.

STEP 2: Internal Coping Strategies

- Activities executed independently without contacting another person e.g., walk and music.

STEP 3: Social Contacts for Distraction

- People and social settings that provide healthy distraction from distress.

STEP 4: Family or Friends for Help

- Trusted individuals who can be informed explicitly that the person is in crisis.

STEP 5: Professionals and Agencies

- Clinicians, emergency rooms and crisis lines e.g., 988 Suicide and Crisis Lifeline.

STEP 6: Making the Environment Safe

- Removing or securing lethal means (firearms, medications and sharp objects).

Research demonstrates that emergency department patients who receive an SPI combined with follow-up outreach are 45% less likely to exhibit suicidal behaviour and twice as likely to engage in outpatient mental health care compared to those receiving standard care (Stanley et al., 2018).

EMPIRICAL EVIDENCE BASE AND RANDOMIZED CONTROLLED TRIALS

The effectiveness of CBT in reducing repeated suicide attempts is supported by a robust body of randomized controlled trial (RCT) evidence, systematic reviews, and meta-analyses.

Key Clinical Trials

Brown et al. (2005) - The Landmark CT-SP Study

In a seminal RCT published in JAMA, Brown and colleagues evaluated 120 adult patients admitted to an emergency department following a suicide attempt. Participants were randomized to receive either standard community care or a 10-session CT-SP protocol.

- **Primary Finding:** Over an 18-month follow-up period, patients in the CT-SP group were 50% less likely to re-attempt suicide compared to the control group (12.4% vs. 24.1%).
- **Secondary Finding:** CT-SP participants demonstrated significantly lower depression severity and reduced scores on the Beck Hopelessness Scale.

18-MONTH RE-ATTEMPT RATES (Brown et al., 2005):



Standard Care: [REDACTED] 24.1%

CT-SP Group: [REDACTED] 12.4% (50% Relative Reduction)

Linehan et al. (2006) - DBT for Chronically Suicidal Individuals

Linehan et al. conducted a multi-center RCT comparing 1 year of comprehensive DBT against therapy by non-behavioral experts in women with chronic suicidal behavior and BPD.

- **Results:** DBT reduced suicide attempts by 50%, resulted in significantly fewer psychiatric emergency department visits, and reduced inpatient psychiatric admissions.

Rudd et al. (2015) - BCBT in Military Personnel

In a trial involving 152 active-duty US Army soldiers presenting with acute suicidal ideation or attempts, soldiers were randomized to receive Brief CBT (BCBT) or Treatment As Usual (TAU).

- **Results:** Soldiers in the BCBT group were **60%** less likely to report a suicide attempt during the 24-month follow-up period compared to soldiers receiving TAU.

Meta-Analyses and Systematic Reviews

Multiple meta-analyses reinforce these individual RCT findings:

- **Tarrier et al. (2008):** Analyzed 28 studies and concluded that cognitive-behavioural interventions significantly reduced suicidal behavior, ideation and hopelessness compared to control conditions.
- **Mewton and Andrews (2016):** Evaluated CBT across age cohorts, confirming moderate-to-large effect sizes in reducing self-harm and suicide attempts in both adult and adolescent populations.
- **Riblet et al. (2017):** Conducted a systematic review on outpatient psychotherapies post-discharge, confirming CBT as one of the few modalities demonstrating explicit efficacy in reducing suicide attempt recurrence.

V. NEUROCOGNITIVE AND BIOMARKER SHIFTS ASSOCIATED WITH CBT

Advances in neuroimaging and neurobiology have begun to clarify how cognitive-behavioural interventions alter the brain networks involved in suicidal behaviour. These are as follows:

Prefrontal Cortex Regulation and Executive Functioning

Suicidal individuals frequently exhibit structural and functional alterations in the prefrontal cortex (PFC), particularly the dorsolateral PFC (dlPFC) and ventral PFC (vPFC), which regulate impulse control, risk assessment and decision-making (Jollant et al., 2011). CBT interventions strengthen top-down inhibitory control from the prefrontal cortex over lower limbic structures. As patients practice cognitive reframing and distress tolerance, neuroimaging indicates increased activation in the dlPFC during emotional stress, correlating with improved behavioral control during acute crises.

Amygdala Down-Regulation

The amygdala plays a central role in processing threat and emotional distress. Suicidal individuals demonstrate amygdala hyper-reactivity when exposed to negative interpersonal stimuli or psychological stress. By repeatedly deploying cognitive reappraisal techniques taught in CBT, patients build functional neural connections between the ventromedial PFC and the amygdala by allowing for rapid down-regulation of emotional distress before it sparks suicidal impulses.

HPA-Axis Stabilization

Chronically high stress and trauma dysregulate the Hypothalamic-Pituitary-Adrenal (HPA) axis, impairing stress recovery and increasing vulnerability to crisis states. Successful CBT interventions have been linked to normalizations in cortisol awakening responses and improved autonomic nervous system regulation (as measured by heart-rate variability) by indicating an enhanced physiological capacity to tolerate stress without escalating to self-harm.



VI. IMPLEMENTATION CHALLENGES AND CLINICAL CONSIDERATIONS

Despite its proven efficacy, implementing CBT for suicide attempt prevention presents unique clinical, systemic, and ethical challenges.

Therapist Anxiety and Countertransference

Working with individuals at high risk for suicide generates significant anxiety among clinicians. Fear of patient death, legal liability and emotional burnout can cause therapists to become overly cautious by leading to inappropriate hospitalization or a breakdown in therapeutic alliance. Effective CBT implementation requires:

Regular and structured clinical supervision and peer consultation.

Protocols emphasizing shared responsibility rather than defensive practice.

Training in objective risk assessment frameworks e.g., C-SSRS - Columbia-Suicide Severity Rating Scale.

Acute Inpatient vs. Outpatient Delivery

While full CBT protocols require outpatient stability over weeks or months, the highest risk period for a repeated suicide attempt occurs immediately following discharge from an inpatient psychiatric unit (Chung et al., 2017). Adapting CBT elements into brief inpatient protocols e.g., delivering 3-4 session modules focusing entirely on Safety Planning, Means Restriction, and core coping strategies prior to discharge, is critical for bridging this transition.

Digital and Telehealth Delivery Options

To expand access, digital adaptations of CBT for suicide prevention such as web-based platforms and mobile apps delivering Safety Planning and cognitive exercises etc. have proliferated. While preliminary research indicates digital CBT can reduce suicidal ideation, empirical evidence establishing that self-guided apps can lower actual suicide attempts without human clinical support remains limited. Hybrid models consisting blending clinician oversight with digital skill practice offer a viable path forward.

Future Study Areas

To further maximize the preventive impact of CBT on attempted suicide, several following emerging areas of research and clinical innovation require attention:

1. **Integration with Machine Learning:** Predictive algorithms analyzing Electronic Health Records (EHR) can identify high-risk individuals before a suicide attempt occurs by allowing for proactive and automated delivery of targeted CBT micro-interventions.
2. **Cultural Adaptation of CBT Protocols:** Modifying cognitive interventions to align with cultural variations in expressiveness, family structures and stigma regarding mental health is essential for reducing health disparities in vulnerable populations.
3. **Biomarker-Informed CBT:** Pairing real-time physiological monitoring e.g., wearable devices tracking heart-rate variability and sleep fragmentation with ecological momentary interventions (EMI) can prompt patients to organize CBT distress tolerance skills precisely when biological indicators signal escalating acute stress.

VI. CONCLUSION

The empirical literature confirms that Cognitive Behavioral Therapy, when delivered through protocols tailored specifically to suicidal behavior is an effective psychotherapeutic intervention for preventing repeated suicide attempts. By moving beyond traditional diagnostic assumptions and targeting the suicidal mode directly, CBT addresses the cognitive, emotional and behavioural mechanisms that drive individuals from severe distress to self-harm.

Through structured clinical mechanisms including cognitive restructuring of helplessness and burdensomeness, problem-solving training, distress tolerance, lethal means restriction and the Stanley-Brown Safety Planning Intervention, CBT replaces lethal escape behaviours with functional coping strategies. The widespread integration of suicide-focused CBT into psychiatric emergency services, outpatient clinics and community mental health systems represents an actionable and evidence-based strategy to save lives and mitigate the global burden of suicidal behaviour.



REFERENCES

1. World Health Organization. (2021). *Suicide worldwide in 2019: Global health estimates*. World Health Organization.
2. Joiner, T. E. (2005). *Why people die by suicide*. Harvard University Press.
3. Tarrier, N., Taylor, K., & Gooding, P. (2008). Cognitive-behavioral interventions to reduce suicide behavior: A systematic review and meta-analysis. *Behavior Modification*, 32(1), 77–108.
4. Beck, A. T. (1996). Beyond belief: A theory of modes, personality, and psychopathology. In P. M. Salkovskis (Ed.), *Frontiers of cognitive therapy* (pp. 1–25). Guilford Press.
5. Shneidman, E. S. (1993). *Suicide as psychache: A clinical approach to self-destructive behavior*. Jason Aronson.
6. Bostwick, J. M., Powney, C., & Lineberry, T. W. (2016). Suicide attempt as a risk factor for completed suicide: A systematic review and meta-analysis. *Journal of Clinical Psychiatry*, 77(8), 1092–1100.
7. Brown, G. K., Ten Have, T., Henriques, G. R., Xie, S. X., Hollander, J. E., & Beck, A. T. (2005). Cognitive therapy for the prevention of suicide attempts: A randomized controlled trial. *JAMA*, 294(5), 563–570.
8. Chung, D. T., Ryan, C. J., Hadzi-Pavlovic, D., Singh, S. P., Stanton, C., & Large, M. M. (2017). Suicide rates after discharge from psychiatric hospital: A systematic review and meta-analysis. *JAMA Psychiatry*, 74(7), 694–702.
9. Jobes, D. A. (2016). *Managing suicidal risk: A collaborative approach* (2nd ed.). Guilford Press.
10. Jollant, F., Lawrence, N. S., Olić, E., Guillaume, S., & Courtet, P. (2011). The suicidal mind and brain: A review of neuropsychological and neuroimaging studies. *World Journal of Biological Psychiatry*, 12(5), 319–339.
11. Linehan, M. M., Comtois, K. A., Murray, A. M., Brown, M. Z., Gallop, R. J., Heard, H. L., Korslund, K. E., Tutek, D. A., Reynolds, S. K., & Lindenboim, N. (2006). Two-year randomized controlled trial and follow-up of dialectical behavior therapy vs therapy by experts for suicidal behaviors and borderline personality disorder. *Archives of General Psychiatry*, 63(7), 757–766.
12. Mewton, L., & Andrews, G. (2016). Cognitive behavioral therapy for suicidal ideation and behaviors in young people: A meta-analysis. *Cognitive Behaviour Therapy*, 45(6), 462–477.
13. Neazua, E. G., & Asarnow, J. R. (2013). Cognitive-behavioral therapy for suicidal youth: State of the science. *Child and Adolescent Psychiatric Clinics of North America*, 22(2), 277–304.
14. Riblet, N. B., Shiner, B., Young-Xu, Y., & Watts, B. V. (2017). Strategies to decrease machine-identified suicide risk: A systematic review. *Journal of Clinical Psychiatry*, 78(9), e1300–e1308.
15. Rudd, M. D. (2006). *The Assessment and Management of Suicidality*. Hogrefe Publishing.
16. Rudd, M. D., Bryan, C. J., Wertenberger, E. G., Peterson, A. L., Young-McCaughan, S., Mintz, J., Williams, S. R., Arne, K. A., Breitbart, J., Delano, K., Wilkinson, E., & Bruce, T. O. (2015). Brief cognitive-behavioral therapy for suicidal soldiers: A randomized controlled trial. *American Journal of Psychiatry*, 172(5), 441–449.
17. Stanley, B., & Brown, G. K. (2012). Safety planning intervention: A brief intervention to mitigate suicide risk. *Cognitive and Behavioral Practice*, 19(2), 256–264.
18. Stanley, B., Brown, G. K., Brenner, L. A., Galfalvy, H. C., Currier, G. W., Knox, K. L., Chaudhury, S. R., Bush, A. L., & Green, K. L. (2018). Comparison of the safety planning intervention with follow-up vs standard care of suicidal patients treated in emergency departments. *JAMA Psychiatry*, 75(9), 894–900.
19. Yip, P. S., Caine, E. D., Yousuf, S., Chang, S. S., Wu, K. C., & Chen, Y. Y. (2012). Means restriction for suicide prevention. *The Lancet*, 379(9834), 2393–2399.

