

Impact of Climate Change on Forest Composition and Vegetation Patterns in the Kumaon Region of Uttarakhand, India

Himani Bisht¹ and Dr. Rakesh Bhooshan Sharma²

¹Research Scholar, Department of Environmental Science

²Supervisor, Department of Environmental Science

Mind Power University, Bhimtal, Nainital

Abstract: *The Kumaon region of Uttarakhand represents a climate-sensitive ecological transition zone where even modest warming produces measurable reorganization of forest composition and vegetation patterns. This paper synthesizes empirical evidence from satellite-based remote sensing, meteorological records, and field observations to assess how climate change is reshaping Kumaon's forests. Three principal findings emerge. First, the region is experiencing directional altitudinal displacement: satellite data from the Gori Ganga watershed show the treeline advancing upward by approximately 680 meters and vegetation boundaries shifting upslope by nearly 596 meters between 1990 and 2022, while the seasonal snowline has risen by over 520 meters. Second, climate change is amplifying disturbance regimes—particularly forest fire—through the expansion of fire-prone chir pine (*Pinus roxburghii*) and the accumulation of combustible litter, creating a feedback loop between vegetation change and fire vulnerability. Third, compositional change is not uniform across forest types: oak-dominated systems, which support higher moisture retention and biodiversity, face competitive displacement by pine under warmer, drier conditions, with cascading consequences for forest-dependent communities. The paper argues that Kumaon's forests are undergoing not simply spatial displacement but structural reorganization, and that effective adaptation requires moving beyond species-specific interventions toward landscape-scale resilience management.*

Keywords: Kumaon Himalaya, climate change, treeline shift, forest composition, chir pine, oak forests, forest fire, vegetation dynamics

I. INTRODUCTION

The Kumaon Himalaya occupies a distinctive position in India's climate change geography. Spanning the districts of Nainital, Almora, Pithoragarh, Champawat, Bageshwar, and Udham Singh Nagar, the region encompasses an altitudinal gradient from subtropical foothills to alpine meadows, hosting forest types that range from *Shorea robusta*-dominated tropical deciduous stands to *Quercus* (oak) and *Rhododendron* assemblages at temperate elevations. This ecological compression—the packing of multiple life zones into a relatively narrow horizontal distance—makes Kumaon forests unusually responsive to thermal shifts. A warming trend that might produce marginal change in a continental interior can, in the Himalaya, translate into the upslope migration of entire vegetation belts.

The region is also politically and socially significant. Uttarakhand's forests cover approximately 70 percent of the state's geographic area, and forest-dependent livelihoods remain widespread. The state government has identified 30,000 hectares of degraded forest land across Kumaon districts for restoration, attributing degradation to grazing pressure, recurrent fires, invasive species, and soil erosion. Climate change interacts with each of these stressors, complicating both diagnosis and response.



This paper addresses a deceptively simple question: what is climate change doing to forest composition and vegetation patterns in Kumaon? The answer, as the evidence reviewed below demonstrates, is neither linear nor spatially uniform. It involves upward displacement of vegetation boundaries, compositional shifts favoring drought- and fire-adapted species, and the amplification of disturbance regimes that mediate between climate forcing and vegetation response. Understanding these dynamics is essential not only for ecological theory but for the practical design of restoration and adaptation programs in a region where forest health and human welfare are tightly coupled.

II. CLIMATE TRENDS IN THE KUMAON REGION

2.1 Temperature and Precipitation Signals

Instrumental records and downscaled climate analyses paint a consistent picture of warming, albeit with nuances that matter for vegetation response. A study of the Betalghat Development Block in the Kumaun Lesser Himalaya found a statistically significant increasing trend in minimum temperatures at 0.03°C per year over recent decades, alongside a decreasing trend in maximum temperatures (−0.02°C per year) and declining rainfall. The asymmetric warming pattern—nights warming faster than days—has ecological significance: it affects respiration rates, phenological timing, and the competitive balance between species with different thermal optima.

Snowline dynamics provide a more visible indicator of climate forcing at higher elevations. Satellite analysis of the Gori Ganga watershed, spanning 1990 to 2022, documented an average snowline rise from approximately 4,665 meters to 5,185 meters—an upward displacement exceeding 520 meters at a rate of more than 16 meters per year. This retreat is not merely a hydrological concern; it represents the upward migration of the thermal boundary that separates snow-dominated from vegetation-dominated landscapes. As the zero-degree isotherm shifts upslope, precipitation that once fell as snow increasingly falls as rain, altering soil moisture regimes and growing-season length across elevation bands.

2.2 The Disturbance Context

Climate trends in Kumaon cannot be understood in isolation from disturbance regimes. Forest fire has become an increasingly severe annual phenomenon, with the 1999 fire season affecting over 22 percent of Uttarakhand's forest area (5,085 square kilometers), of which 1,225 square kilometers burned severely. More recent seasons have seen fires spreading across both Kumaon and Garhwal, threatening biodiversity, water sources, and human settlements. The interaction between warming, drying, and fire is central to the compositional dynamics discussed below.

III. VEGETATION AND TREELINE DYNAMICS

3.1 Upward Migration of Vegetation Boundaries

The most direct evidence of climate-driven vegetation change in Kumaon comes from the Gori Ganga watershed study, which documented simultaneous upward shifts of the snowline, vegetation boundary, and treeline. The treeline advanced by approximately 680 meters over the 32-year study period, while the broader vegetation boundary moved upslope by nearly 596 meters. Professor D.S. Parihar of Kumaon University, who led the research, explicitly linked these shifts to “sustained warming and reduced snow persistence that now allow forests and crops to colonise elevations once dominated by snow”.

These figures should be interpreted with appropriate caution. Treeline ecotones are notoriously heterogeneous, and remote sensing classifications may conflate true upward migration with changes in canopy density or shrub encroachment. Nevertheless, the convergence of multiple indicators—snowline, vegetation line, and treeline—strengthens the inference that a coherent altitudinal reorganization is underway.

3.2 Compositional Consequences: The Pine-Oak Dynamic

Altitudinal displacement is only one dimension of forest change. Of equal or greater ecological significance is the shifting competitive balance among dominant species. The Kumaon Himalaya supports extensive stands of chir pine (*Pinus roxburghii*), a fire-adapted, drought-tolerant species that has expanded markedly in recent decades, often at the expense of broadleaf associates including oak (*Quercus* spp.). Doctoral research examining forest vulnerability in the Kumaon region found that local community perceptions of climate impacts varied significantly depending on whether



their nearest forest was oak-dominated or pine-dominated, reflecting the divergent ecological services provided by these forest types .

The mechanism linking climate to compositional change operates through both direct physiological effects and fire-mediated feedbacks. Chir pine produces thick, resinous needle litter that creates a highly flammable surface layer; Google Arts & Culture’s documentation of Kumaon forest fires explicitly identifies “the thick blanket of pine needles deposited by invasive Chir Pine” as a primary fire hazard . Warmer and drier conditions increase both the frequency of ignition events and the flammability of this litter, favoring pine regeneration over oak, which is more sensitive to fire and requires longer fire-free intervals to establish. The result is a positive feedback: climate warming promotes pine dominance, pine dominance increases fire probability, and fire further disadvantages oak. This dynamic has been observed across the western Himalaya, and its consequences for biodiversity, soil stability, and water yield are substantial.

3.3 Phenological Shifts

Beyond distributional and compositional change, climate change is altering the timing of biological events. A study examining the impact of environmental gradients on phenometrics of major forest types in the Kumaon region used time-series satellite data to characterize vegetation phenology, finding that temperature and precipitation regimes exert significant control over the timing of green-up and senescence . While phenological research in Kumaon remains less developed than in temperate systems with longer observational records, the available evidence suggests that spring onset is advancing and growing seasons are lengthening at higher elevations—changes that may initially increase productivity but also disrupt synchronized ecological interactions, including those between plants, pollinators, and herbivores.

IV. DISTURBANCE REGIMES AS MEDIATORS OF CLIMATE IMPACT

4.1 Fire as an Amplifier

The relationship between climate and fire in Kumaon is not merely one of direct causation. Fire acts as a mediating variable that translates climatic warming into vegetation change, sometimes with greater force than temperature alone. The extensive 1999 and 1995 fire seasons established a pattern that has persisted: dry winters and early springs, combined with accumulated pine litter, create conditions in which small ignition sources—whether human or natural—can produce large, severe burns . The 2014 Kumaon fires, documented in photographic archives, illustrate the visual and ecological magnitude of these events .

The feedback between fire and vegetation composition has implications for carbon storage and climate mitigation. Biomass burning is a significant source of greenhouse gas emissions, contributing as much as 40 percent of gross carbon dioxide and 30 percent of tropospheric ozone globally according to IPCC estimates cited in regional assessments . Kumaon’s fires thus contribute to the very climatic forcing that promotes their own recurrence—a dynamic that complicates any simple narrative of climate as an external driver.

4.2 Invasive Species and Degradation

Climate change interacts with biological invasions in ways that are only beginning to be documented in Kumaon. The Uttarakhand Forest Department’s restoration planning explicitly identifies *Lantana camara* and *Parthenium* as targets for removal, recognizing that these invasives compete with native regeneration and alter fire regimes . While the causal attribution of invasive spread to climate change specifically remains uncertain, warmer conditions and increased disturbance frequency generally favor fast-growing, disturbance-adapted species over slower-growing natives. The restoration program’s combination of assisted natural regeneration, enrichment planting, and invasive removal represents an implicit acknowledgment that passive recovery is unlikely under current conditions .



V. IMPLICATIONS FOR FOREST-DEPENDENT COMMUNITIES AND MANAGEMENT

5.1 Livelihood Consequences

Forest change in Kumaon is not an abstract ecological phenomenon. The region's population depends heavily on forests for fuelwood, fodder, small timber, and non-timber forest products, and approximately 36.5 percent of the population lives below the poverty line, heightening vulnerability to ecological disruption. The shift from oak to pine dominance has direct consequences for these livelihoods: oak forests provide higher-quality fodder and support more diverse understory vegetation than pine forests, which are relatively depauperate and whose needle litter suppresses herbaceous growth. A doctoral study at TERI School of Advanced Studies concluded that while local communities recognize climate vulnerability, their adaptive capacity varies significantly based on the forest type nearest to them, with oak-dependent communities facing different challenges than pine-dependent ones.

5.2 Restoration and Adaptation Priorities

The Uttarakhand Forest Department's identification of 30,000 hectares of degraded forest land for restoration, linked to the RECAP4NDC project and India's Bonn Challenge commitments, provides an institutional vehicle for climate-adapted management. However, the effectiveness of restoration depends on whether it accounts for the dynamics described in this paper. Three priorities emerge from the evidence:

First, restoration must prioritize the maintenance and expansion of oak-dominated stands, which provide superior hydrological and biodiversity functions and are more resilient to moderate climate change than pine monocultures. Assisted natural regeneration of oak, combined with protection from fire and grazing, may be necessary where seed sources have been depleted.

Second, fire management cannot be reduced to suppression alone. The pine-fire feedback loop requires breaking through strategic interventions—mechanical removal of pine litter in high-value zones, prescribed burning where appropriate, and the creation of fuel breaks around settlements and water sources. The restoration program's inclusion of soil-moisture conservation structures represents a complementary approach, addressing the hydrological consequences of vegetation change.

Third, monitoring must be designed to detect change at the scales at which it occurs. The Gori Ganga study demonstrates the value of long-term satellite records for tracking altitudinal shifts, while the phenological research highlights the need for finer temporal resolution to capture changes in growing-season dynamics. LiDAR-based forest mapping, which Uttarakhand is piloting, offers promise for characterizing structural change in three dimensions, moving beyond simple cover estimates.

VI. RESEARCH GAPS AND DIRECTIONS FOR DOCTORAL INQUIRY

The existing literature on Kumaon's forest-climate dynamics, while growing, remains fragmentary. Several gaps are particularly pertinent for doctoral-level research:

Species-specific response functions. We lack quantitative dose-response relationships between climatic variables (temperature, precipitation, snow duration) and the demographic rates (establishment, growth, mortality) of dominant tree species in Kumaon. Without such data, projections of future composition remain speculative.

Belowground processes. The soil carbon and nutrient consequences of pine expansion into former oak habitat are poorly characterized. Given the importance of soil carbon for both climate mitigation and site productivity, this is a critical gap.

Fire-vegetation-climate feedbacks. While the pine-fire connection is qualitatively recognized, quantitative modeling of how fire return intervals interact with climate projections to determine future forest states has not been undertaken for Kumaon specifically.

Community adaptation and agency. The TERI study represents an important beginning, but research on how forest-dependent communities are actively responding to change—through species selection in agroforestry, migration, or collective management innovations—remains sparse.



A doctoral program addressing any of these areas would contribute meaningfully to both the theoretical understanding of mountain forest dynamics and the practical knowledge base for adaptation in a region where the stakes are high and the window for proactive management is narrowing.

VII. CONCLUSION

Climate change is reorganizing Kumaon's forests in ways that are measurable, directional, and consequential. The upward displacement of treelines and vegetation boundaries—on the order of hundreds of meters over three decades in the Gori Ganga watershed—represents one of the clearest signals of ecological response to warming anywhere in the Indian Himalaya. Yet the more ecologically significant changes may be compositional: the competitive release of chir pine, the decline of oak, and the fire-mediated feedbacks that reinforce these trends. These changes threaten biodiversity, alter hydrological regimes, and undermine the forest-based livelihoods on which much of the region's population depends.

The policy response, exemplified by Uttarakhand's restoration planning, is nascent but potentially transformative. Its success will depend on whether it is designed around the dynamics that this paper has described—not as a static problem of degraded area but as a dynamic problem of shifting competitive regimes, altered disturbance cycles, and communities whose adaptive capacity is unevenly distributed. For doctoral research, the Kumaon Himalaya offers a setting in which fundamental questions about climate-vegetation feedbacks, disturbance ecology, and socio-ecological resilience converge with urgent practical needs. The forests are moving; the task is to understand where, how fast, and with what consequences.

REFERENCES

- [1] Shodhganga. (n.d.). *The shift in vegetation, forest cover and biodiversity will have positive or negative impact on the livelihood of local communities*. Chapter 2.
- [2] Times of India. (2026, January 17). *How Himalayas' snowline — from Gori Ganga to Everest — is retreating upslope, causing treeline to shift upwards*.
- [3] Pant, D., Tiwari, P.C., Nand, K., Kumar, R., & Kweera, M. (2024). Analyzing temperature and precipitation regimes in hilly terrains of Betalghat Development Block, Kumaun Lesser Himalaya. *Google Scholar*.
- [4] Google Arts & Culture. (2014). *Forest Fire in Kumaon*.
- [5] Drishti IAS. (2025, November 29). *Uttarakhand Identifies Degraded Land for Restoration*.
- [6] *Impact of Climate Variability on Forest Ecosystems in Uttarakhand*. (n.d.). American Journal of Geographical Research.
- [7] DunValley Mail. (2026, May 20). *Forests are burning, breaths are being suffocated*. (In Hindi).
- [8] TERI School of Advanced Studies. (2017). *Climate change vulnerability of forests and forest-dependent people in the central Himalayan region* (Doctoral abstract, A. Chakraborty).
- [9] *Impact of Environmental Gradients on Phenometrics of Major Forest Types of Kumaon Region of the Western Himalaya*. (n.d.). OUCI.

