IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 7, October 2026
Impact Factor: 8.2

Breast Enlargement from A Cosmetic Perspective: Ayurvedic and Modern Medical Review

Dr. Madhuri Sanjay Bhalgat[1] and Dr. Sharayu Hanumandas Mantri[2]
[1] Professor and Head of Department
[2] PG Scholar, M.S. Prasuti Tantra Evam Stri Roga
[1] [2]Department of Prasuti Tantra Evam Stri Roga
SVNH Ayurved Mahavidyalaya, Rahuri Factory, Ahilyanagar, Maharashtra, India

Abstract: *Background: Breast size and contour vary considerably among individuals and are influenced by genetic, hormonal, developmental and environmental factors. Interest in cosmetic breast enlargement includes surgical procedures as well as non-surgical and traditional preparations. Ayurveda discusses stana, stanya and maternal nourishment within its broader account of female reproductive physiology; these concepts require careful interpretation when considered alongside contemporary cosmetic practice.*
Objective: To discuss the anatomical and physiological determinants of breast volume, relevant Ayurvedic concepts, available modern cosmetic interventions, their safety profile and the principles of responsible counselling.
Review approach: This article brings together established anatomical and physiological knowledge, contemporary professional and regulatory information on breast augmentation, and commonly taught Ayurvedic concepts relevant to stana and stanya. It is a focused educational review and does not report original patient data or a clinical intervention.
Discussion: Implant-based augmentation and autologous fat transfer can increase breast volume, but both involve limitations and potential complications. Exercise, massage, topical products and marketed supplements have not demonstrated reliable, durable breast enlargement. Ayurvedic concepts may support culturally sensitive discussion of health and nourishment, but they do not establish the efficacy of a specific breast-enlargement formulation.
Conclusion: Cosmetic breast enlargement requires individualised assessment, realistic expectations and informed consent. Traditional claims should be distinguished from clinically demonstrated outcomes, and patient safety should remain central.

Keywords: Stana; breast augmentation; Ayurveda; breast anatomy; cosmetic gynaecology; breast implants; autologous fat transfer; patient counselling.

I. INTRODUCTION

Breast size is a normal aspect of human anatomical variation. It is not, by itself, a measure of femininity, reproductive potential, general health or the ability to breastfeed. Some individuals seek a change in breast volume or contour because of perceived small breast size, asymmetry, developmental variation, changes after pregnancy and lactation, weight loss or previous surgery. The clinician's first responsibility is to understand the individual's concern and expectations without implying that one breast size is medically or socially preferable.

Cosmetic breast enlargement should be distinguished from the evaluation of a breast lump, sudden or unilateral enlargement, congenital breast or chest-wall abnormalities, endocrine disorders and post-mastectomy reconstruction. A new lump, nipple discharge, skin or nipple retraction, persistent focal pain, redness or other inflammatory changes require appropriate diagnostic assessment rather than cosmetic treatment.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 7, October 2026
Impact Factor: 8.2

II. AIM AND SCOPE

This review examines cosmetic breast enlargement from Ayurvedic and modern medical perspectives. It outlines breast structure and physiology, discusses the relevance and limits of Ayurvedic concepts, summarises established cosmetic procedures and non-surgical claims, and highlights safety, ethical counselling and areas requiring further research.

III. ANATOMY AND PHYSIOLOGY OF THE BREAST

The breast is a modified sweat gland situated in the superficial fascia of the anterior chest wall. It contains glandular tissue, fibrous connective tissue and adipose tissue. The glandular component is arranged into lobes and lobules, with ducts converging towards the nipple. Cooper's suspensory ligaments contribute to breast support and contour. The relative proportions of glandular and adipose tissue differ between individuals and change over time.

Oestrogen contributes to ductal development, while progesterone and other hormonal signals participate in lobuloalveolar development. Prolactin is central to milk synthesis and oxytocin to milk ejection. Puberty, menstrual cycling, pregnancy, lactation, menopause, changes in body composition and exposure to exogenous hormones may alter breast fullness and tissue characteristics. These physiological relationships do not justify hormonal manipulation solely for cosmetic enlargement.

Breast volume is multifactorial. Genetic background, age, body-fat distribution, nutritional status, endocrine function, pregnancy history and ageing all contribute. No single clinical parameter reliably predicts an individual's final breast size.

IV. AYURVEDIC PERSPECTIVE

4.1 Stana and its physiological relevance

Ayurvedic literature describes stana as an important feature of the female body and discusses the breasts in relation to reproductive physiology, maternal nourishment and feeding of the infant. References to breast development and function occur within broader discussions of sharira, garbha and stanya. The classical explanatory framework uses concepts such as dosha, dhatu, agni and srotas. These are traditional constructs and should not be equated directly with modern anatomical structures, hormones or physiological pathways.

4.2 Rasa dhatu and stanya

In commonly taught Ayurvedic accounts, stanya is described as an upadhatu associated with rasa dhatu. This relationship is relevant to the traditional understanding of maternal nourishment and lactation. It does not demonstrate that increasing a particular dhatu, administering a rasayana or improving general nourishment will selectively increase breast volume.

4.3 Interpretation of stana-vriddhi claims

The term stana-vriddhi may appear in contemporary discussions of breast enlargement. Any claim attributed to a classical text should be supported by the exact passage, edition, chapter and commentary. General references to nourishment, tissue support, massage or rasayana cannot be interpreted as clinical proof of predictable breast enlargement. Classical descriptions were not designed as controlled clinical trials and cannot independently establish efficacy, dose, contraindications or long-term safety.

4.4 Herbal preparations, oils and topical applications

Various herbal preparations, oils, creams and massage practices are promoted for breast enlargement. Robust human evidence showing a reproducible and durable increase in breast volume from these products is lacking. Topical products may cause irritation or allergic contact dermatitis. Oral products may vary in composition, interact with medicines or contain undeclared pharmacologically active substances. Products with hormonal or hormone-like activity require particular caution, especially during pregnancy, lactation and in people with hormone-sensitive conditions. Traditional use alone does not establish safety or efficacy.

Any Ayurvedic medicine should be considered only after individual assessment by a qualified practitioner. Product identity, ingredients, manufacturing quality, dose, contraindications, possible interactions and adverse effects require

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 7, October 2026
Impact Factor: 8.2

attention. No formulation should be described as a proven breast-enlargement treatment without suitable product-specific clinical evidence.

V. MODERN MEDICAL PERSPECTIVE

5.1 Assessment before cosmetic intervention

Assessment should explore the reason for seeking enlargement, duration of concern, developmental and pregnancy history, medication and hormone exposure, smoking status, relevant medical and surgical history, breast symptoms, family history and expectations. Examination may document breast base width, skin envelope, nipple-areola position, asymmetry, ptosis, chest-wall characteristics and available soft-tissue coverage. Imaging should be guided by age, symptoms, risk profile and applicable screening recommendations. Cosmetic intent does not replace indicated diagnostic evaluation.

5.2 Implant-based breast augmentation

Implant-based augmentation is an established surgical option for increasing breast volume. Implants differ in fill material, shape, surface and dimensions. Selection is individualised according to anatomy, desired outcome and the surgeon's assessment. Potential complications include bleeding, infection, pain, scarring, altered breast or nipple sensation, asymmetry, implant malposition, capsular contracture, rupture or leakage, and anaesthetic complications. Further surgery may be required.

Breast implants are not lifetime devices. The likelihood of complications and additional surgery increases with the duration of implantation. Patients should receive device-specific information and understand the need for follow-up. The US Food and Drug Administration identifies breast implant-associated anaplastic large cell lymphoma (BIA-ALCL), a lymphoma of the immune system rather than breast carcinoma, as an implant-associated risk. It has been reported more frequently with textured than smooth implant surfaces. The FDA has also communicated reports of squamous cell carcinoma and other lymphomas in the capsule around implants; these reports are rare and remain under evaluation [1,2].

5.3 Autologous fat transfer

Autologous fat transfer involves harvesting adipose tissue, usually by liposuction, processing it and injecting it into the breast. It avoids a permanent implant, but the final volume is variable because some transferred fat is resorbed. More than one session may be required. Reported complications include fat necrosis, oil cysts, calcifications, infection, contour irregularity and morbidity at the donor site. Post-procedure imaging findings should be assessed by clinicians familiar with fat grafting [3].

5.4 Non-surgical approaches

Exercise can strengthen the pectoral muscles and improve posture, but it does not enlarge breast glandular tissue. Weight gain may increase breast adipose tissue in some individuals, but the distribution and degree of change are unpredictable; weight gain is not a targeted treatment for isolated breast enlargement. Massage, creams, vacuum devices and supplements have not been shown to produce reliable, durable enlargement. Available evidence does not support the use of hormones solely to increase breast size because systemic risks may outweigh an unproven cosmetic benefit [4,5].

5.5 Psychological and ethical considerations

Consultation should explore whether expectations are realistic and whether body-image distress is disproportionate or associated with significant functional impairment. If body dysmorphic disorder, coercion or external pressure is suspected, elective cosmetic intervention should be deferred until appropriate assessment. Shared decision-making should include the option of no intervention. Consent must be voluntary, informed and free from pressure.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 7, October 2026
Impact Factor: 8.2

VI. COMPARATIVE OVERVIEW

Domain	Ayurvedic perspective	Modern medical perspective
Conceptual basis	Stana, stanya, dhatu, dosha and nourishment concepts.	Anatomy, physiology, clinical examination and outcomes research.
Evidence for cosmetic enlargement	No established high-quality evidence for a specific Ayurvedic product as a reliable breast-volume enhancer.	Implants and fat transfer can increase volume; outcomes and risks are described in clinical practice.
Counselling focus	Individual assessment, traditional context, product identity and safety.	Anatomy, goals, alternatives, complications and follow-up.
Safety priorities	Ingredient verification, interactions, contamination and adverse reactions.	Surgical and anaesthetic risks, implant-related risks, imaging and possible reoperation.

VII. DISCUSSION

The principal distinction between Ayurvedic concepts and contemporary cosmetic interventions concerns the nature and strength of evidence available for measurable breast-volume change. Implant surgery and autologous fat transfer have defined procedural techniques, measurable outcomes and recognised complications, although neither is risk-free. In contrast, claims that an oral or topical Ayurvedic preparation selectively enlarges breast tissue have not been established by convincing, reproducible clinical trials.

An integrative approach does not require treating all interventions as equally effective. Ayurvedic knowledge may contribute to culturally responsive communication and discussion of traditional health practices. However, any medicine or procedure proposed specifically for cosmetic enlargement should be evaluated independently through pharmacological characterisation, quality assurance, safety monitoring and appropriately designed clinical studies.

For postgraduate clinical practice, the clinician should identify concerns reflecting normal anatomical variation, explain what can and cannot be changed, recognise symptoms requiring diagnostic work-up and discuss evidence-based options without promoting unrealistic expectations. Clear communication that no non-surgical method has reliably demonstrated durable breast enlargement is an important part of counselling.

VIII. RESEARCH PRIORITIES

Future research should address the following priorities:

Clearly define the study population, intervention, comparator, validated outcome measures and follow-up duration.

Authenticate botanical ingredients and report chemical characterisation, contaminant testing and batch-to-batch quality control for herbal products.

Measure objective breast volume using validated methods, alongside patient-reported satisfaction, durability and adverse events. Distinguish breast-volume change from changes related to body weight, hydration, menstrual phase, pregnancy or lactation.

Obtain appropriate ethics approval, register prospective clinical trials where applicable, and report methods and outcomes transparently.

IX. CONCLUSION

Cosmetic breast enlargement requires individualised, ethical and evidence-informed counselling. Implant-based augmentation and autologous fat transfer can increase breast volume, but both carry meaningful risks and may require

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 7, October 2026
Impact Factor: 8.2

further procedures. Exercise, massage, creams and marketed supplements do not have reliable evidence for durable breast enlargement. Ayurvedic concepts concerning stana, stanya and tissue nourishment provide a traditional framework for understanding female physiology, but they do not establish the efficacy of a particular enlargement treatment. Claims should be communicated cautiously, and patient safety must remain paramount.

REFERENCES

1. Standring S, editor. Gray's Anatomy: The Anatomical Basis of Clinical Practice. 42nd ed. London: Elsevier; 2020.
2. U.S. Food and Drug Administration. Risks and Complications of Breast Implants [Internet]. Silver Spring (MD): FDA;updated 2023 Dec 14 [cited 2026 Oct 31]. Available from: https://www.fda.gov/medical-devices/breast-implants/risks-and-complications-breast-implants
3. American Society of Plastic Surgeons. Fat Transfer Breast Augmentation: Risks and Safety [Internet]. ArlingtonHeights (IL): ASPS [cited 2026 Oct 31]. Available from: https://www.plasticsurgery.org/cosmetic-procedures/fat-transfer-breast-augmentation/safety
4. Mayo Clinic. Natural breast enhancement: Do supplements work? [Internet]. Rochester (MN): Mayo Foundation forMedical Education and Research; 2024 Jul 9 [cited 2026 Oct 31]. Available from: https://www.mayoclinic.org/healthy -lifestyle/womens-health/expert-answers/natural-breast-enhancement/faq-20057834
5. Fugh-Berman A. 'Bust enhancing' herbal products. Obstet Gynecol. 2003;101(6):1345-1349. doi:10.1016/S0029-7844(03)00362-4.
6. U.S. Food and Drug Administration. Questions and Answers about Breast Implant-Associated Anaplastic Large CellLymphoma (BIA-ALCL) [Internet]. Silver Spring (MD): FDA [cited 2026 Oct 31]. Available from: https://www.fda.gov/ medical-devices/breast-implants/questions-and-answers-about-breast-implant-associated-anaplastic-large-cell-lymp homa-bia-alcl
7. American Society of Plastic Surgeons. Breast Augmentation: Risks and Safety [Internet]. Arlington Heights (IL): ASPS[cited 2026 Oct 31]. Available from: https://www.plasticsurgery.org/cosmetic-procedures/breast-augmentation/safety
8. Sharma PV, translator. Caraka-Samhita. Varanasi: Chaukhambha Orientalia. Classical concepts of stana, stanya anddhatu should be cited to the exact edition, sthana, chapter and verse consulted.
9. Shastri A, editor. Sushruta-Samhita. Varanasi: Chaukhambha Sanskrit Sansthan. Classical descriptions should becited to the exact edition, sthana, chapter and verse consulted.
10. Vagbhata. Ashtanga Hridaya. Classical references should be cited to the exact edition, translator, chapter and verseconsulted.