IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 1, November 2026
Impact Factor: 8.2

A Review On Potential Role of Jamun Leaves in Herbal Handwash Paper Soap

Miss. Bhosale Shraddha J*[1] and Ms. Anap Vidya R*[2]
Student, Department of Pharmacy[1]
Guide, Department of Pharmacy[2]
Mrs. Saraswati Wani of Pharmacy, Ganegaon, Maharashtra
Affiliated to Dr.Babasaheb Ambedkar Technology University, Lonere, Raigad
ayushbhosale952@gmail.com

Abstract: *Hand hygiene is an important daily practice for reducing the spread of many microorganisms. Conventional liquid handwashes are useful, but they are usually supplied in plastic containers and may be inconvenient to carry in small bags or while travelling. Paper soap is a thin, lightweight sheet containing cleansing ingredients. When placed in wet hands, the sheet dissolves or disperses and produces a lather that can be rinsed away. Herbal ingredients are being explored as additional components in personal-care products. Jamun, commonly known as Indian blackberry, is a medicinal tree whose leaves contain several classes of plant compounds, including phenolics and flavonoids. Published laboratory studies have reported antioxidant and antibacterial activity in certain Jamun leaf extracts and essential oils. These findings provide a scientific reason to investigate Jamun leaves as a botanical ingredient in handwash paper soap. However, evidence for the leaf extract itself should not be confused with proof that a finished paper-soap product prevents infection. The final product must be formulated and tested for cleansing performance, physical quality, microbial quality, skin compatibility and stability. This review summarises the botanical background, reported phytochemicals, relevant biological findings, possible formulation role, preparation approach, evaluation tests and research gaps associated with Jamun-leaf-based paper soap.*

Keywords: Jamun leaves, *Syzygium cumini*, herbal handwash, paper soap, antibacterial activity, phytochemicals, cosmetic formulation

I. INTRODUCTION

Handwashing with soap and clean water is a basic hygiene measure. Hands frequently touch phones, door handles, money, public surfaces and other objects. Washing with soap helps loosen dirt, oils and microorganisms so that they can be removed during rinsing. A handwash product is therefore expected to clean effectively, be convenient to use and remain acceptable to the skin.

Paper soap is a portable form of cleansing product. It is generally prepared by applying a suitable soap or surfactant solution to a thin, water-compatible sheet, followed by drying. A small sheet can be carried without a bottle. On contact with water, it should soften, dissolve or break apart sufficiently to release the cleansing material. Its performance depends on the type of sheet, surfactant, plasticiser, extract, drying method and storage conditions.

The interest in herbal cosmetics has encouraged the study of plant extracts as possible supporting ingredients. Jamun leaves are of interest because scientific literature describes a range of phytochemicals and biological activities. This review considers their possible use in paper soap as an ingredient for further investigation, not as an established replacement for validated cleansing agents.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 1, November 2026
Impact Factor: 8.2



II. BOTANICAL PROFILE OF JAMUN

Jamun is the common Indian name for *Syzygium cumini* (L.) Skeels, a tree belonging to the family Myrtaceae. It is also referred to as Java plum, black plum or jambolan in different regions. The tree is cultivated in many parts of India and other tropical and subtropical areas. Its fruits are well known, while the leaves have also been used traditionally and investigated in pharmacological research.

2.1 Description of the plant

The Jamun tree is generally evergreen and may grow to a considerable height. Its leaves are simple, opposite, smooth and leathery, with a characteristic aromatic smell when crushed. The flowers are small and pale, and the fruit changes colour as it matures, commonly becoming deep purple. For a cosmetic formulation, correctly identifying the plant is essential because different species may have different chemical compositions and biological properties.

2.2 Selection and collection of leaves

Healthy, mature leaves should be collected from a correctly identified plant, away from roadsides, industrial emissions and pesticide-treated areas. Damaged, diseased or visibly contaminated leaves should be rejected. Leaves should be washed with clean water to remove dust, drained and dried under shade with good air circulation. Direct intense sunlight may affect some plant constituents. Dried leaves can be coarsely powdered and stored in a clean, dry, closed container until extraction.

2.3 Why consider Jamun leaves?

Jamun leaves are a locally available plant material in many Indian regions. Published reviews describe phenolic acids, flavonoids, tannins and other constituents in different parts of the plant. Some laboratory investigations of leaf extracts and leaf essential oil have reported antibacterial effects against selected test organisms. This makes the leaves a reasonable candidate for preliminary cosmetic research. Availability alone, however, does not establish safety, effectiveness or suitability for use on skin.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 1, November 2026
Impact Factor: 8.2

III. PHYTOCHEMICAL CONSTITUENTS

Plant extracts contain mixtures of compounds rather than one single active chemical. The amount and type of compounds can vary with the plant variety, location, season, maturity of leaves, drying conditions and extraction solvent. Reviews of *S. cumini* describe phenolic compounds such as gallic acid and ellagic acid, flavonoids such as quercetin and myricetin, tannins and volatile constituents. Not every compound will be present in every extract, and a particular formulation should be analysed rather than assuming a fixed composition.

Phenolic compounds: These plant constituents are widely studied for their ability to participate in antioxidant reactions. Their presence may contribute to the measured antioxidant activity of an extract.

Flavonoids: Flavonoids are a broad group of plant polyphenols. Some have shown biological activity in laboratory models, including antioxidant and antimicrobial effects. Their activity depends on the specific molecule and test conditions.

Tannins: Tannins are polyphenolic substances that can interact with proteins. They may contribute to astringent properties and have been studied for antimicrobial effects. Their concentration should be controlled because excessive tannin content may affect colour, feel or compatibility.

Volatile constituents: Essential oil obtained from Jamun leaves contains volatile compounds. The reported profile differs among studies and extraction methods. Essential oil is not interchangeable with an aqueous or hydroalcoholic leaf extract.

IV. REPORTED BIOLOGICAL ACTIVITIES RELEVANT TO HAND HYGIENE

4.1 Antibacterial activity

Antibacterial activity means the ability of a material to inhibit bacterial growth or reduce bacterial survival under defined test conditions. Mohamed and colleagues investigated crude extracts and essential oils from Jamun leaves and reported antibacterial activity, with the methanolic extract showing stronger activity than some of the other tested preparations in their experimental system. The study also measured antioxidant properties. These findings support further investigation of Jamun leaf extracts, but they do not establish the effectiveness of a handwash paper soap.

A later study of *S. cumini* leaf essential oil analysed its chemical composition and tested activity against selected standard bacterial strains and multidrug-resistant clinical isolates. The oil contained multiple volatile constituents, with alpha-pinene reported as a major component in that sample. The results demonstrate that leaf essential oil can have measurable antibacterial activity in laboratory assays. Essential oil composition and activity may vary according to geographical origin and extraction conditions.

Antibacterial tests are commonly performed using agar diffusion methods or broth dilution methods. In an agar diffusion test, a sample is placed on an inoculated agar plate and the clear zone around it is measured. In broth dilution, the sample is tested at different concentrations to estimate the minimum inhibitory concentration (MIC). A zone of inhibition alone does not predict how well a soap sheet will clean hands. Surfactants, extract solubility, contact time, dilution during washing and rinsing can all change performance.

4.2 Antioxidant activity

Antioxidants are substances that can reduce or delay oxidation reactions in suitable test systems. Jamun leaf extracts have been evaluated using assays such as DPPH radical scavenging and ferric reducing antioxidant power (FRAP). Mohamed et al. reported antioxidant activity in the extracts they studied, with the methanolic extract showing stronger activity than some comparison samples in their tests. Antioxidant activity may be relevant to the chemical characterisation of the extract and its stability, but it is not the same as antibacterial action or a direct measure of hand-cleansing ability.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 1, November 2026
Impact Factor: 8.2

4.3 Other reported properties

Reviews describe several biological activities associated with different parts of Jamun, including anti-inflammatory and other pharmacological effects. These broader findings should be interpreted carefully. Evidence obtained from fruit, seeds, bark or laboratory animal studies cannot automatically be applied to a leaf extract in a cosmetic product. For this paper-soap concept, the most relevant evidence is the published work specifically examining Jamun leaves and their extracts.

4.4 Important distinction

A herbal ingredient may show activity when concentrated in a laboratory test, yet show little or no measurable effect after it is mixed into a finished product. A paper soap is diluted with water and used for a short period. Therefore, the complete formulation—not only the raw extract—must be tested. The product should be described as a cleansing paper soap unless appropriate product-level tests support stronger claims.

V. POSSIBLE ROLE OF JAMUN LEAF EXTRACT IN PAPER SOAP

Jamun leaf extract may be explored as a botanical additive in a paper-soap formulation. The main cleansing function should still be provided by a suitable, skin-compatible surfactant or soap base. The extract is not a substitute for the cleansing base, and its addition should not make the sheet brittle, sticky, strongly coloured, difficult to dissolve or irritating.

5.1 Potential formulation contribution

Botanical value: Including a characterised plant extract can provide a clear herbal research theme and allow the formulation to be compared with a base paper soap without extract.

Potential antibacterial contribution: Since selected Jamun leaf extracts have shown antibacterial activity in laboratory studies, it is reasonable to test whether the extract retains measurable activity after incorporation into the paper-soap matrix. This is a research question, not a confirmed product benefit.

Phenolic and antioxidant profile: The extract can be assessed for total phenolic content and antioxidant activity before and after formulation. Such tests may help understand whether processing and storage affect the extract.

Local plant resource: Jamun leaves may be accessible in many Indian communities. Responsible sourcing and correct botanical identification are important if the ingredient is to be developed further.

5.2 Choice of extraction method

Water extraction is relatively simple and avoids using alcohol, but it may extract a different range of compounds and can be more prone to microbial spoilage. Hydroalcoholic extraction can recover a broader range of polar and moderately polar constituents, but the solvent must be removed appropriately and residual solvent controlled. The choice should be based on the intended product, extract profile, safety and available laboratory facilities. Essential oil extraction produces a different material and should not be described as equivalent to leaf extract.

5.3 Formulation considerations

The paper substrate should be thin, flexible, absorbent and compatible with the chosen cleansing base. The surfactant should produce adequate lather and rinse cleanly. A plasticiser or humectant may be used in a controlled amount to reduce brittleness, while too much may make the sheet tacky or slow to dry. The Jamun extract should be added at a low, experimentally selected concentration after preliminary compatibility and safety checks. Its colour, odour, pH effect and solubility should be monitored.

The extract may darken the sheet or change its smell. These changes are not necessarily signs of failure, but they affect consumer acceptability and batch consistency. A suitable preservative system may be needed depending on water

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 1, November 2026
Impact Factor: 8.2

content, raw materials and packaging. Preservative effectiveness and microbial quality should be assessed using appropriate methods.

5.4 Safety and skin compatibility

Natural origin does not guarantee that an ingredient is non-irritant or non-allergenic. Plant extracts can contain substances that cause sensitivity in some people. The final product should undergo an appropriate safety assessment, including review of ingredient quality, pH, irritation potential and microbiological quality. Human patch testing, if considered, should be conducted only under suitable ethical and professional supervision.

VI. SUGGESTED LABORATORY PREPARATION OF JAMUN HERBAL PAPER SOAP

The following is a research-development outline for a student laboratory. It is not a validated commercial formula. The exact quantities should be selected through small pilot batches and approved by the project guide. Use cosmetic-grade materials and follow institutional laboratory safety procedures.

6.1 Materials

Possible materials include authenticated dried Jamun leaves, purified water, a suitable water-compatible surfactant or mild liquid soap base, a flexible water-soluble paper or thin sheet substrate, a compatible plasticiser such as glycerin, and a suitable preservative if required. Equipment may include a weighing balance, grinder, sieve, beakers, measuring cylinder, glass rod or magnetic stirrer, filtration cloth or filter paper, heating arrangement where appropriate, drying tray or hot-air oven at a controlled low temperature, pH meter and sealed packaging.

6.2 Preparation of leaf extract

First, wash and shade-dry the leaves. Grind the dried material to a coarse powder. Weigh a measured quantity and add a selected extraction solvent, such as purified water or an appropriate hydroalcoholic mixture, using a documented plant-to-solvent ratio. Allow extraction with occasional mixing for a defined period. Filter the mixture and, if required, repeat extraction. Combine the filtrates and concentrate using a suitable low-temperature method. Record the starting weight, solvent, extraction time, final extract weight or volume and percentage yield. Store the extract in a clean, labelled container under suitable conditions.

6.3 Preparation of paper soap sheet

Prepare the cleansing solution by mixing the selected surfactant base with purified water. Add the measured plasticiser and mix gently to minimise excess foam. Add a small, preselected amount of Jamun leaf extract and mix until uniform. Check the pH and appearance. Spread a measured quantity of the mixture evenly over the selected sheet or substrate using a clean applicator. Dry under controlled conditions until the sheet is sufficiently dry and can be handled without sticking. Cut into equal-sized pieces and pack immediately in moisture-resistant sachets or a suitable airtight container.

6.4 Suggested development groups

For meaningful comparison, prepare at least three groups: (A) base paper soap without Jamun extract; (B) paper soap with a lower selected extract concentration; and (C) paper soap with a higher selected extract concentration. Keep the sheet size, surfactant level, plasticiser level, drying conditions and packaging as consistent as possible. Record all batch details. Do not assume that the higher extract concentration will perform better; it may affect sheet quality, pH, colour, odour or skin feel.

VII. EVALUATION PARAMETERS FOR HERBAL PAPER SOAP

Evaluation should examine the raw extract, the paper-soap sheet and the performance of the finished product. Tests should be performed using written procedures and suitable controls.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429 Volume 6, Issue 1, November 2026 Impact Factor: 8.2

7.1 Physical and organoleptic evaluation

Appearance and colour: Observe the sheet for uniformity, spots, cracks and colour variation. Record the colour of the extract and final sheet.

Odour: Record the characteristic odour and any unpleasant or changed smell during storage.

Weight and dimensions: Weigh individual sheets and measure length, width and thickness where possible. This helps assess batch uniformity and dose consistency.

Flexibility and handling: Observe whether the sheet bends without cracking, tears during handling or becomes sticky.

Disintegration or dissolution time: Place a sheet in a defined volume and temperature of water using a standardised agitation method. Record the time required for the sheet to break up or dissolve sufficiently. Use the same method for all batches.

7.2 Physicochemical evaluation

pH: Prepare a defined aqueous dispersion of the sheet and measure with a calibrated pH meter. Report the sample concentration and temperature. pH should be appropriate for the intended rinse-off product and should be assessed by a qualified formulator.

Foam characteristics: Measure foam formation and, if relevant, foam persistence using a consistent method. Foam volume alone does not prove better cleansing.

Moisture content: Determine moisture using a suitable validated or standard laboratory method. Excess moisture can affect sheet stability and microbial quality.

Extract content / total phenolics: If facilities are available, determine extract loading or total phenolic content using a suitable analytical method. This helps compare batches and monitor processing.

Stability: Store packed samples under defined conditions and inspect changes in colour, odour, flexibility, dissolution, pH and microbial quality at scheduled intervals.

7.3 Microbiological evaluation

The microbial quality of the finished product should be checked using suitable pharmacopoeial or cosmetic microbiology procedures and relevant limits. If antibacterial activity is being studied, use a recognised method with appropriate positive and negative controls, standardised inoculum, defined contact time and replicate testing. Test the complete formulated product, not just the concentrated extract. A neutralisation step may be needed to prevent residual surfactant from continuing to act during the assay. Such testing should be conducted in an equipped laboratory under trained supervision.

7.4 Skin compatibility and user acceptability

Assess ingredient safety before any use on volunteers. A qualified supervisor should determine whether a controlled skin compatibility test is appropriate and obtain required ethical approval and informed consent. User acceptability may include ease of carrying, ease of handling, dissolution, fragrance, rinsing and after-feel. Avoid testing on broken or diseased skin.

7.5 Packaging

Paper soap can absorb moisture from humid air. Packaging should protect sheets from moisture, dust and contamination while allowing convenient single-sheet removal. Compare packaging options for moisture protection and ease of use.

Record any sticking, softening or loss of flexibility during storage.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 1, November 2026
Impact Factor: 8.2

VIII. DISCUSSION, LIMITATIONS AND FUTURE SCOPE

Published research gives a scientific basis for investigating Jamun leaves as a botanical ingredient. Reviews describe a diverse phytochemical profile, while laboratory studies have reported antioxidant and antibacterial activity for selected leaf extracts or essential oils. These findings make Jamun a relevant candidate for exploratory cosmetic formulation. Nevertheless, the evidence has important limitations. Extracts prepared with different solvents are chemically different; the composition can vary between samples; and laboratory antimicrobial tests do not reproduce the short, diluted contact that occurs during handwashing.

A key research gap is the limited direct evidence for a finished Jamun-leaf handwash paper-soap product. A student project can address this gap by preparing a base sheet and extract-containing sheets, then comparing their physical quality, dissolution, pH, stability and laboratory antimicrobial performance. A good project should report negative or inconclusive findings honestly. It should also avoid claiming that the product prevents disease or replaces proper handwashing.

IX. CONCLUSION

Jamun (*Syzygium cumini*) leaves contain several groups of phytochemicals and have been investigated for antioxidant and antibacterial activities. These published findings support studying the leaf extract as a possible herbal additive in handwash paper soap. The main cleansing action must come from a suitable surfactant system, while the extract should be treated as an ingredient whose compatibility, safety and contribution need to be demonstrated. Further work should standardise the extract, optimise the sheet composition, assess stability and skin compatibility, and test the complete product using appropriate microbiological methods. At present, Jamun-leaf paper soap should be considered a promising research concept rather than a clinically proven antibacterial product.

REFERENCES

1. Kumari N, Kumar M, Chaudhary N, et al. Exploring the Chemical and Biological Potential of Jamun (*Syzygium cumini* (L.) Skeels) Leaves: A Comprehensive Review. *Chemistry & Biodiversity*. 2023;20(9):e202300479. DOI: 10.1002/cbdv.202300479.
2. Mohamed AA, Ali SI, El-Baz FK. Antioxidant and antibacterial activities of crude extracts and essential oils of *Syzygium cumini* leaves. *PL0olE*. 2013;8(4):e60269. DOI: 10.1371/journal.pone.0060269.
3. Phytochemical Analysis, Antibacterial Activity and Modulating Effect of Essential Oil from *Syzygium cumini* (L.) Skeels. *Molecules*. 2022;27(10):3281. DOI: 10.3390/molecules27103281.
4. Ayyanwar M, Subash-Babu P. *Syzygium cumini* (L.) Skeels: A review of its phytochemical constituents and traditional uses. *Asian Pacific Journal of Tropical Biomedicine*. 2012;2(3):240–246. DOI: 10.1016/S2221-1691(12)60050-1.
5. Chikara N, Kaur R, Jag lan S, Sharma P, Gat Y, Panghal A. Bioactive compounds and pharmacological and food applications of *Syzygium cumini* – a review. *Food & Function*. 2018;9:6096–6115. DOI: 10.1039/C8FO00654G.
6. Yadav D, Kapoor R. *Syzygium cumini* L. leaves: A comprehensive review on phytochemical composition, antidiabetic mechanisms, and therapeutic prospects. *Journal of Pharmacognosy and Phytochemistry*. 2025;14(6):203–210. DOI:
7. 10.22271/phyto.2025.v14.i6c.15659.
8. Tambe BD, Pedhekar P, Pardeshi H. Phytochemical Screening and Antibacterial Activity of *Syzygium cumini* (L.) (Myrtaceae) Leaves Extracts. *Asian Journal of Pharmaceutical Research and Development*. 2021;9(5):50–54. DOI: 10.22270/ajprd.v9i5.1023.
9. Kumari N, Kumar M, Chaudhary N, et al. Exploring the Chemical and Biological Potential of Jamun (*Syzygium cumini* (L.) Skeels) Leaves: A Comprehensive Review. *Chemistry & Biodiversity*. 2023;20(9):e202300479. DOI: 10.1002/cbdv.202300479.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 1, November 2026
Impact Factor: 8.2

10. Mohamed AA, Ali SI, El-Baz FK. Antioxidant and antibacterial activities of crude extracts and essential oils of *Syzygium cumini* leaves. *PL0olE*. 2013;8(4):e60269. DOI: 10.1371/journal.pone.0060269.
11. Phytochemical Analysis, Antibacterial Activity and Modulating Effect of Essential Oil from *Syzygium cumini* (L.) Skeels. *Molecules*. 2022;27(10):3281. DOI: 10.3390/molecules27103281.
12. Ayyanwar M, Subash-Babu P. *Syzygium cumini* (L.) Skeels: A review of its phytochemical constituents and traditional uses. *Asian Pacific Journal of Tropical Biomedicine*. 2012;2(3):240–246. DOI: 10.1016/S2221-1691(12)60050-1.
13. Chikara N, Kaur R, Jag lan S, Sharma P, Gat Y, Panghal A. Bioactive compounds and pharmacological and food applications of *Syzygium cumini* – a review. *Food & Function*. 2018;9:6096–6115. DOI: 10.1039/C8FO00654G.
14. Yadav D, Kapoor R. *Syzygium cumini* L. leaves: A comprehensive review on phytochemical composition, antidiabetic mechanisms, and therapeutic prospects. *Journal of Pharmacognosy and Phytochemistry*. 2025;14(6):203–210. DOI: 10.22271/phyto.2025.v14.i6c.15659.
15. Tambe BD, Pedhekar P, Pardeshi H. Phytochemical Screening and Antibacterial Activity of *Syzygium cumini* (L.) (Myrtaceae) Leaves Extracts. *Asian Journal of Pharmaceutical Research and Development*. 2021;9(5):50–54. DOI: 10.22270/ajprd.v9i5.1023.