IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 1, November 2026
Impact Factor: 8.2

An Integrated Ayurvedic and Modern Clinical Evaluation of Acne Vulgaris and Hirsutism in Women

Correlation of Tvak, Rakta and RaktavahaSrotas Dushti with Hormonal, Inflammatory and Advanced Diagnostic Biomarkers

Dr. Bhalgat Madhuri[1], Dr. Kamlu Aishwarya[2], Dr Nilkanth Ankita[3]
HOD Streerog Prasutitantra Department[1]
PG Diploma student Streerog Prasutitantra Department[2]
SVNH Ayurved Mahavidyala Rahuri Factory Ahilyanagar[1,2]
Assistant Professor Matoshri Asarabai Darade Ayurved College, Babhulgaon, Yeola[3]

Abstract: *Acne vulgaris and hirsutism represent hyperandrogenic and inflammatory dermatological manifestations that cause substantial psychosocial morbidity in women. Conventional management targets distal pathways (androgen blockade, follicular desquamation, microbial suppression) but often suffers from relapse post-discontinuation and systemic side effects. Classical Ayurveda categorizes these presentations under Mukhadushika (Yuvanapidika), TvakDushti, and RaktavahaSrotas Dushti.*
Objective: To synthesize classical Ayurvedic etiology (Nidana), pathogenesis (Samprapti), and therapeutics (Chikitsa) with modern endocrine-inflammatory pathophysiology, establishing an evidencemapped diagnostic and clinical research protocol.
Methods: A systematic cross-mapping of Charaka Samhita, Sushruta Samhita, and Ashtanga Hridaya against biomedical markers (free testosterone, DHEAS, SHBG, IL-1α, TNF-α, dermal sebum metrics) was executed, paired with a prospective RCT protocol (N=120).
Conclusion: Bridging Rakta Prasadana/Shodhana therapies with precise hormonal and dermatoscopic profiling offers a holistic, disease-modifying strategy for female hyperandrogenism.

85% 10%	*70-80%*	*Pitta-Kapha*	
ACNE PREVALENCE IN YOUNG FEMALES	*HIRSUTISM GLOBAL PREVALENCE*	*PCOS CO-MORBIDITY RATE*	*DOMINANT DUSHTI TRAIT*

Key Words: *Acne vulgaris*

I. INTRODUCTION

Acne vulgaris and hirsutism are among the most frequent clinical presentations in female dermatology and endocrinology clinics worldwide. Epidemiological surveys demonstrate that acne affects up to 85% of adolescent and adult women, with persistent adult acne increasing in prevalence over the past two decades. Hirsutism affects approximately 5–10% of women of reproductive age and serves as a key clinical marker of cutaneous hyperandrogenism Escobar-Morreale (2012).

The pathophysiological intersection of acne and hirsutism frequently points to underlying systemic disorders, most prominently Polycystic Ovary Syndrome (PCOS), which accounts for 70–80% of hyperandrogenic presentations, non-classic congenital adrenal hyperplasia (NCCAH), hyperinsulinemia, and rare androgensecreting neoplasms. Current conventional therapies—including oral contraceptives, spironolactone, systemic isotretinoin, and topical retinoids—are effective but limited by potential adverse events, teratogenicity, posttreatment recurrence, and failure to target holistic metabolic-lifestyle drivers Thiboutot et al. (2018).

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429 Volume 6, Issue 1, November 2026 Impact Factor: 8.2

Ayurvedic Medicine offers a comprehensive framework that views localized cutaneous pathology as the external manifestation of systemic Doshic imbalance, Agni impairment, and systemic tissue toxicity (Ama) circulating within the RaktavahaSrotas (circulatory channels). This investigation bridges ancient Ayurvedic wisdom with modern clinical science to establish a diagnostic protocol and research design.

II. AYURVEDIC CONCEPTUAL FRAMEWORK

A. Classical Pathophysiology of Tvak, Rakta, and RaktavahaSrotas

According to classical text Sushruta Samhita (Sharira Sthana, Adhyaya 4), the skin (Tvak) consists of seven distinct layers, each nourished by and anatomically linked to specific Dhatus (tissues), particularly Rasa and Rakta Dhatu. The outer layers (Avabhasini and Lohita) reflect the systemic health of Rakta Dhatu and Pitta Dosha.

तासांपथमाऽवभासिनीनाम, यासर्ववर्णान्अवभासयतिपन्चविधाम्चछायाप्रकाशयती | साि◌सध्म-मुखदू◌ि◌षकाि◌धष्ठाना ||

Tāsāṁprathamā'vabhāsinīnāma, yāsarvavarṇānavabhāsayatipañcavidhāṁ ca chāyāṁprakāśayati | Sāsidhma-mukhadūṣikādhiṣṭhānā ||

Sushruta Samhita, ShariraSthana 4/4: "The first layer of skin is named Avabhasini; it reflects all complexions and illuminates the five types of shade. It is the primary site/seat of Sidhma (tinea/macules) and Mukhadushika (acne)."

Ayurvedic Concept	Classical Etiology (Nidana)	Symptomatology (Lakshana)	Pathophysiological Correlation
TvakDushti	Ahitakara Ahara, Viruddhahara, Excessive Katu/Lavana/Amla, Vega Dharana.	Vaivarnya (discoloration), Pidika (papules/pustules), Kandu (itching), Rukshata/Snigdhataa.	Epidermal barrier disruption, altered desquamation, follicular hyperkeratinization.
Rakta Dushti	Pitta-vitiating foods, alcohol, excessive anger, exposure to heat/sun, improper Shodhana.	Rakta-pidika, Daha (burning sensation), Paka (suppuration), Visarpa-like spread.	Systemic pro-inflammatory cytokine release (IL-1, TNF-α), oxidative stress.
Raktavaha Srotas Dushti	Snaigdha-Ushna-Drava Ahara, Vidahi foods, excessive sun, suppressed vomiting.	Kustha, Visarpa, Pidika, Asradara, Upakusha, Mukhapaka, Plihavriddhi.	Vascular engorgement, dermal microvascular alteration, chronic low-grade systemic inflammation.

II. AYURVEDIC CONCEPTUAL FRAMEWORK (CONTINUED)

B. Mukhadushika (Yuvanapidika) &Lomaharsha / Excessive Hair Growth

In classical literature, facial eruptions occurring in youth are termed Mukhadushika or Yuvanapidika. The pathology is primarily attributed to the simultaneous vitiation of Kapha, Pitta, and Rakta, which obstruct the hair follicles (Romakupa) on the face Vagbhata (Astanga Hridaya, Uttaratantra 31/5).

शाल्मलीकण्टकाकाराःि◌पडकाःसरुचोघनाः | कफवातरक्तजाः◌ेयामुखदू◌ि◌षकाः ||

Śālmalīkaṇṭakākārāḥpiḍakāḥsarucoghanāḥ | Kaphavātaraktajāḥjñeyāmukhadūṣikāḥ ||

Ashtanga Hridaya, Uttaratantra 31/5: "Eruptions resembling the thorns of the Shalmali tree (Salmaliamalabarica), accompanied by pain, hardness, and caused by Kapha, Vata, and Rakta are known as Mukhadushika."

While classical texts do not have a single direct term equivalent to androgenic hirsutism, features of excessive body and facial hair in females are understood under the umbrella of Atiloma or Asadharna Roma Vriddhi, associated with ArtavavahaSrotas Dushti (menstrual channel dysfunction), MedovahaSrotasDushti (lipid metabolism dysfunction), and Vata-Kapha dominance affecting hair follicles (Romakupa).

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429 Volume 6, Issue 1, November 2026 Impact Factor: 8.2

C. Diagnostic Equivalence and Conceptual Boundaries

It is vital in modern integrative medicine to avoid simplistic one-to-one translations between Ayurvedic traditional concepts and biomedical diseases:

Classical Ayurvedic Concept

Focuses on systemic energetics (Pitta-Kapha imbalance), Agni (digestive/metabolic energy), Ama (circulating Metabolic toxins), and tissue-level impairment in Rasa, Rakta, and Meda Dhatus within Srotas channels.

Modern Biomedical Science

Focuses on 5α-reductase activity, hyperandrogenemia (testosterone, DHEAS), insulin resistance, CutibacteriumAcnes colonization, and upregulation of inflammatory Pathways (NLRP3 inflammasome, IL-1α).

Integrative Hypothesis 1 Rakta Dushti correlated clinical features (erythema, burning, suppuration) correspond with elevated systemic and tissue inflammatory cytokines (TNF-α, high-sensitivityCRP, IL-6).

Integrative Hypothesis 2


Artavavaha / MedovahaSrotasDushti with Kapha dominance corresponds with hyperinsulinemia, altered LH:FSH ratio, and enlarged polycystic ovarian morphology.

III. PATHOPHYSIOLOGY OF ACNE VULGARIS & HIRSUTISM

Acne vulgaris is a chronic inflammatory disorder of the pilosebaceous unit driven by four primary pathophysiological factors: (1) androgen-mediated hyperseborrhea, (2) follicular hyperkeratinization leading to infundibular microcomedone formation, (3) proliferation of Cutibacterium acnes (formerly Propionibacterium acnes) within the clogged pore, and (4) secondary inflammatory cascades involving Toll-like receptor 2 (TLR-2) activation, IL-1α release, and neutrophil recruitment Zouboulis et al. (2014).

B. Pathophysiology of Hirsutism & Cutaneous Androgen Action

Hirsutism is defined as the presence of excessive terminal hair in females in androgen-dependent anatomical sites (upper lip, chin, chest, upper/lower abdomen, back, upper arms, thighs). It must be distinguished from hypertrichosis, which is androgen-independent generalized vellus hair growth. Circulating free testosterone and dihydrotestosterone (DHT)—converted locally by the enzyme 5α-reductase type 2 within the hair follicle—bind to androgen receptors in the dermal papilla cells, transforming fine vellus hairs into thick, pigmented terminal hairs Rosenfield & Ehrmann (2015).



IV. DIFFERENTIAL DIAGNOSIS & RED FLAGS

Accurate clinical differentiation is essential to separate isolated benign dermatological conditions from serious metabolic or neoplastic etiologies. Red flags demanding urgent endocrine or surgical referral include rapid onset (<6 months), severe virilization (clitoromegaly, voice deepening, male-pattern alopecia), and marked serum testosterone elevation (>150–200 ng/dL).

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429 Volume 6, Issue 1, November 2026 Impact Factor: 8.2

Condition	Distinguishing Clinical Features	Diagnostic Investigations	Red Flags & Indications for Referral
PCOS	Oligomenorrhea/amenorrhea, acne, hirsutism, central obesity, acanthosis nigricans.	Pelvic ultrasound (polycystic morphology), LH:FSH ratio > 2, free T elevated.	Severe metabolic dysregulation, endometrial hyperplasia risk.
Non-Classic CAH	Post-pubertal onset acne, hirsutism, irregular menses, family history.	Early morning baseline 17OHP > 200 ng/dL, ACTH stimulation test.	Adrenal crisis (rare in nonclassic), marked hyperandrogenemia.
Androgen-Secreting Tumors	Rapidly progressive hirsutism, severe virilization, pelvic or abdominal mass.	Total testosterone > 200 ng/dL, DHEAS > 700 µg/dL, CT/MRI imaging.	RED FLAG: Rapid onset (<6 mos), virilization; immediate referral required.
Cushing Syndrome	Moon facies, buffalo hump, purple abdominal striae, hypertension, hirsutism.	24-hr urinary free cortisol, overnight 1 mg dexamethasone suppression.	Severe hypertension, hypokalemia, progressive proximal muscle weakness.
Acneiform Eruption	Monomorphic papulopustules, absence of comedones, history of drug exposure.	Detailed medication history (corticosteroids, lithium, anticonvulsants, B12).	Sudden widespread onset postnew drug initiation.

V. INTEGRATED DIAGNOSTIC EVALUATION & ADVANCED TECHNOLOGIES

A rigorous evaluation requires integrating classical Ayurvedic constitutional assessment with quantitative biomedical assays and digital diagnostic technology to capture both systemic imbalance and local tissue pathology.

A. Standardized Ayurvedic Clinical Proforma

Integrated Proforma Domains

- Prakriti & Vikriti Assessment: Standardized score for Vata, Pitta, and Kapha phenotypic traits and current pathological state.
- Agni & Ama Evaluation: Assessment of digestive capacity (Manda/Tikshna/Vishama/Sama) and signs of systemic toxins (tongue coating, sluggishness, irregular bowel).
- Srotas Dushti Examination: Specific inquiry into Raktavaha (burning, skin flushes), Artavavaha (menstrual volume, length, pain), and Medovaha (weight changes, adiposity) Srotas.
- Tvak Lakshana Scoring: Quantitative grading of facial redness (Raga), pustular suppuration (Paka), hardness/induration (Khara), and oiliness (Snigdhataa).

B. Biomedical Laboratory Protocol

Endocrine screening must be performed during the early follicular phase (days 2–5 of spontaneous or withdrawal menses):

- **Hormonal Panel:** Total Testosterone (LC-MS/MS preferred), Sex Hormone-Binding Globulin (SHBG), Calculated Free Testosterone, DHEAS, early-morning 17-Hydroxyprogesterone, Serum Prolactin, Fasting TSH.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429 Volume 6, Issue 1, November 2026 Impact Factor: 8.2

- **Metabolic & Inflammatory Panel:** Fasting Insulin, Fasting Glucose (HOMA-IR calculation), Lipid Profile, high-sensitivity C-Reactive Protein (hs-CRP), IL-6.

C. Advanced Technology-Assisted Assessments

Modern non-invasive dermatological technologies provide objective baseline measurements and track treatment efficacy over time:

- Computer-Assisted Skin Image Analysis (VISIA): Quantitative measurement of porphyrins (C. acnes activity), sebum output, erythema index, and red areas.
- Digital Dermoscopy&Trichoscopy: High-magnification examination of follicular plug density, vascular patterns, and hair shaft diameter in hirsute zones.
- Phototrichogram& Digital Hair Density Analysis: Precise tracking of hair density (hairs/cm²) and anagen:telogen ratio to evaluate response to antiandrogen/Ayurvedic therapy.

VI. AYURVEDIC THERAPEUTIC EVALUATION

A. Topically Applied Lepa Formulations for Mukhadushika

Herbal pastes (Lepa) applied topically provide localized anti-inflammatory, antimicrobial, and desquamative actions without systemic toxicity. Below is a detailed pharmacological and classical analysis of key ingredients.

लेपःशाल्मलीकण्टकैःपयसाि◌ंपष्टैःसम्यक्मुखदू◌ंषकाघ्नः | लोधधान्यवचालेपःतथावक्त्रप्रसादनः ||

Lepaḥśālmalīkaṇṭakaiḥpayasāpiṣṭaiḥsamyakmukhadūṣikāghnaḥ |
Lodhradhānyavacālepaḥtathāvaktraprasādanaḥ ||
Sarangadhara Samhita, Madhyama Khanda 11/11:

"Application of paste made from Shalmali thorns ground with milk destroys Mukhadushika. Similarly, a paste of Lodhra, Dhanyaka, and Vacha enhances facial clarity and clears eruptions."

Botanical / Sanskrit Name	Classical Indication	Modern Bioactive & Pharmacological Evidence	Safety & Skin Tolerability
Lodhra Symplocos racemosa (Symplocaceae)	Mukhadushika, TvakDushti, PittaRakta Hara	Contains symplocososide. Demonstrates 5α-reductase inhibition, anti-inflammatory, and astringent properties Kumar et al. (2016).	Non-irritating; low risk of allergic contact dermatitis. Excellent for oily skin.
Vacha Acorus calamus (Acoraceae)	Kapha-Vatahara, Lekhana, Vedanasthapana	Rich in α- and β-asarone. Antibacterial activity against C. acnes and S. epidermidis; antiinflammatory.	Mild potential for localized skin sensitization; restrict to low concentrations (<2%).
Haridra Curcuma longa (Zingiberaceae)	Kusthaghna, Varnya, Vishaghna, Lekhana	Curcuminoids suppress NF-κB activation, downregulate TNF-α and IL-1β, inhibit C. acnes lipase.	Transient yellow skin staining; rare contact hypersensitivity. Highly effective.
Chandana Santalum album (Santalaceae)	Pittaghna, Dahaharana, Prasadana	α-santalol demonstrates powerful anti-inflammatory and tyrosinase-inhibiting (antipigmentary) activity.	Safe and non-irritating; ideal for post-inflammatory hyperpigmentation.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 1, November 2026
Impact Factor: 8.2

VI. AYURVEDIC THERAPEUTIC EVALUATION

B. Rakta Prasadana& Rakta Shodhana Dravya (Systemic Interventions)

In Ayurvedic pathology, systemic clearance of Rakta Dushti is required to prevent recurrent skin eruptions. While classical literature describes these herbs as "blood purifiers" (Rakta Shodhaka), biomedical research translates these effects into antioxidant upregulation, hepatic clearance enhancement, systemic antiinflammatory pathways, and modulation of endocrine sensitivity.

Drug (Botanical Name)	Ayurvedic Properties	Traditional Rationale	Preclinical & Clinical Evidence	Reproductive & Safety Profile
Manjistha Rubia cordifolia	TiktaKashayaMadhura; Ushna; Pitta-Kapha Hara	Rakta Prasadana, Raktashodhaka, Kusthaghna	Purpurin and alizarin downregulate cyclooxygenase-2 (COX-2) and 5lipoxygenase (5-LOX), inhibiting acne-related inflammation Pawar et al. (2011).	Avoid in high doses during 1st trimester of pregnancy. No significant hepatotoxicity.
GuduchiTinospora cordifolia	TiktaKashaya; Ushna; Tridoshahara	Rasayana, Dahaprashamana, Vayasthapana	Cordifolioside A enhances insulin sensitivity, reduces TNF-α, and modulates macrophage activity. Useful in hyperinsulinemic PCOS.	Extremely safe profile; well-tolerated in longterm human clinical studies.
Nimba Azadirachta indica	Tikta-Katu; Sheeta; Kapha-Pitta Hara	Krimighna, Kusthaghna, Raktashodhaka	Nimbolide and nimbidin exert potent antibacterial activity against resistant C. acnes strains; immunomodulatory.	Avoid seed oil internally during pregnancy (abortifacient potential). Leaf extracts safe at standard doses.
SarivaHemidesmus indicus	Madhura-Tikta; Sheeta; Tridoshahara	Daha-prashamana, Anulomana, Raktashodhaka	2-hydroxy-4-methoxybenzaldehyde provides significant free-radical scavenging and anti-lipid peroxidation activity.	High margin of safety; excellent tolerability in adolescent and adult females.

C. Safety, Quality Control, and Heavy Metal Testing

All oral Ayurvedic formulations utilized in clinical practice must meet strict pharmacopoeial standards (API). Mandatory testing includes Atomic Absorption Spectroscopy (AAS) to rule out lead, mercury, arsenic, and cadmium contamination, pesticide residue screening, and microbial load limits per WHO guidelines WHO (2011).

VII. CONVENTIONAL MANAGEMENT & INTEGRATIVE FRAMEWORK

A. Guideline-Based Conventional Therapy

Contemporary management of acne and hirsutism relies on validated international clinical guidelines Zaenglein et al. (2016):

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 1, November 2026
Impact Factor: 8.2

- Topical Acne Therapies: Benzoyl peroxide (2.5–5%), topical retinoid (adapalene 0.1%, tretinoin 0.025–0.05%), topical azelaic acid (15–20%), and clindamycin phosphate combined with BPO to prevent resistance.
- Systemic Acne Therapies: Oral tetracyclines (doxycycline 100 mg/day, minocycline 100 mg/day) restricted to 3–4 month courses; oral isotretinoin (0.5–1.0 mg/kg/day) for severe nodulocystic or treatmentresistant acne under strict REMS/pregnancy prevention programs.
- Hirsutism Therapies: Combined Oral Contraceptives (COCs) containing non-androgenic progestins (drospirenone, cyproterone acetate) as first-line pharmacotherapy. Oral Spironolactone (50–200 mg/day) as antiandrogen add-on after 6 months of COC therapy. Direct mechanical removal: Laser Hair Reduction (755 nm Alexandrite or 1064 nm Nd:YAG) and electrolysis.

VIII. CLINICAL TRIAL PROTOCOL (CONTINUED)

C. Trial Visit Schedule & Outcome Assessment

Visit / Timeline	Screening (V0) Day -7 to 0	Baseline (V1) Day 1	Week 4 (V2) ± 3 Days	Week 8 (V3) ± 3 Days	Week 16 (V4) End of Study
Informed Consent & Eligibility	✓				
GAGS &mFG Clinical Scoring	✓	✓	✓	✓	✓
Ayurvedic Proforma (Dosha/Srotas)	✓	✓	✓	✓	✓
VISIA Imaging &Dermoscopy		✓		✓	✓
Hormonal & Inflammatory Panel		✓			✓
Safety Labs (Liver/Kidney Function)	✓		✓		✓

IX. DISCUSSION & RESEARCH PRIORITIES

This study bridges ancient Ayurvedic clinical observational frameworks with contemporary endocrinedermatological science. By correlating Rakta Dushti symptoms with concrete biomarkers (hs-CRP, IL-6, free testosterone) and utilizing digital facial analysis (VISIA), the protocol establishes a rigorous foundation for integrative research. Key advantages of this approach include targeting systemic metabolic-inflammatory drivers while delivering localized symptom relief, potentially mitigating the high recurrence rates seen with isolated topical or hormonal therapies.

REFERENCES

1. Tan J, Bhate K. A global perspective on the epidemiology of acne. Br J Dermatol. 2015;172 Suppl 1:3-12.
2. Escobar-Morreale HF. Polycystic ovary syndrome: definition, aetiology, diagnosis and treatment. Nat Rev Endocrinol. 2018;14(5):270284.
3. Thiboutot D, Dreno B, Abanmi A, et al. Practical management of acne for clinicians: An international consensus from the Global Alliance to Improve Outcomes in Acne. J Am Acad Dermatol. 2018;78(2S1):S1-S23.
4. Vagbhata. Astanga Hridaya. Expressed with Vidyotini Hindi Commentary by Kaviraj Atrideva Gupta. Varanasi: Chaukhambha Prakashan; 2019.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 1, November 2026
Impact Factor: 8.2

5. Zouboulis CC, Picardo M, Ju Q, et al. Beyond acne: Current aspects of sebaceous gland biology and function. Rev EndocrMetabDisord. 2016;17(3):319-334.
6. Rosenfield RL, Ehrmann DA. The Pathogenesis of Polycystic Ovary Syndrome (PCOS): The Hypothesis of PCOS as Functional Ovarian Hyperandrogenism Revisited. Endocr Rev. 2016;37(5):467-520.
7. Kumar S, Lad S, et al. In vitro 5alpha-reductase inhibitory activity of selected Ayurvedic medicinal plants. J Ethnopharmacol. 2016;188:102-108.
8. Pawar AT, Shivakumar FR. Anti-inflammatory and anti-acne activities of Rubia cordifolia L. stem extracts. Oriental Pharm Exp Med. 2011;11(3):189-197.
9. World Health Organization. Quality Control Methods for Herbal Materials. Geneva: World Health Organization; 2011.
10. Zaenglein AL, Pathy AL, Schlosser BJ, et al. Guidelines of care for the management of acne vulgaris. J Am Acad Dermatol. 2016;74(5):945-973.