IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 5, September 2026
Impact Factor: 8.2

Benzimidazole: A Versatile Pharmacophore for the Development of Bioactive Compounds—An Updated Review.

Anurag Singh[1], Akash Singh[2], Gurumeet Wadhawa[3], Sajid Shaikh[4]
[1,2]Student P.G. Department of Chemistry
[3]Assistant Professor Department of chemistry
[4]I/C Principal and Head Department of Chemistry
[1,2,3]Veer Wajekar College, Phunde
[4]Anjuman Islam Degree College, Murud, Janjira

Abstract : *Benzimidazole is an important aromatic nitrogen-containing heterocyclic compound characterized by the fusion of a benzene ring with an imidazole moiety. This unique bicyclic framework represents a privileged structural scaffold in medicinal chemistry and has attracted considerable attention in the development of biologically active molecules. Modification of the benzimidazole nucleus through the introduction of diverse substituents, particularly at the 2- and 5-positions, provides considerable opportunities for structural optimization and the generation of therapeutically promising derivatives. Owing to its versatile chemical properties and favorable biological profile, benzimidazole serves as an important building block in the design and synthesis of several pharmacologically active compounds.*

Over the past decades, numerous synthetic approaches have been developed for the preparation of benzimidazole derivatives, reflecting their significance in contemporary organic and medicinal chemistry. A broad spectrum of biological activities has been reported for these compounds, including antimicrobial, antiviral, anticancer, anti-inflammatory, analgesic, anti-ulcer, antidiabetic, and other pharmacological effects. The biological potential of benzimidazole derivatives is strongly influenced by the nature, position, and electronic properties of substituents attached to the heterocyclic nucleus.

The present review provides a comprehensive overview of the synthetic strategies, structural modifications, and pharmacological properties of substituted benzimidazole derivatives reported in the literature. Particular emphasis is placed on the relationship between structural features and biological activity, with the objective of providing useful insights into structure–activity relationship (SAR) studies. This review may serve as a valuable reference for researchers involved in synthetic chemistry, medicinal chemistry, drug discovery, and the development of novel benzimidazole-based therapeutic agents.

Key Words: Benzimidazole; Nitrogen-containing heterocycles; Heterocyclic chemistry; Synthetic strategies; Structure–activity relationship (SAR); Antimicrobial activity; Antiviral activity; Anticancer activity; Anti-inflammatory activity; Analgesic activity; Anti-ulcer activity; Antidiabetic activity; Medicinal chemistry; Drug discovery.

I. INTRODUCTION

Benzimidazole is an important aromatic, nitrogen-containing heterocyclic compound that has attracted considerable attention in organic chemistry, medicinal chemistry, pharmaceutical sciences, and drug discovery. Structurally, benzimidazole is a bicyclic heterocyclic system formed by the fusion of a benzene ring with an imidazole ring. This

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429 Volume 6, Issue 5, September 2026 Impact Factor: 8.2

fused architecture provides an extended π-electron system and two chemically distinct nitrogen atoms, which contribute to its aromaticity, hydrogen-bonding ability, coordination behavior, and broad range of biological interactions. Because of these structural and electronic characteristics, the benzimidazole nucleus is widely recognized as a privileged pharmacophore for the design and development of biologically active molecules.

The discovery and development of benzimidazole chemistry have evolved considerably over the past several decades. Early investigations were primarily focused on understanding the synthesis and fundamental chemical properties of the nucleus, whereas modern research emphasizes the rational modification of the benzimidazole scaffold to obtain compounds with improved potency, selectivity, pharmacokinetic properties, and therapeutic efficacy. The incorporation of different functional groups into the benzimidazole framework can significantly influence its biological profile. Consequently, benzimidazole has become an attractive scaffold for generating structurally diverse compounds through molecular hybridization, ring substitution, annulation, and other synthetic strategies.

The chemical structure of benzimidazole is particularly suitable for structural diversification. The N-1 position can undergo alkylation, arylation, acylation, and other modifications, while the C-2 position is highly valuable for introducing aryl, heteroaryl, alkyl, carbonyl, nitrile, and other pharmacologically relevant substituents. Substitution on the benzene portion, especially at the 5- and 6-positions, can further modify electronic, steric, lipophilic, and hydrogen-bonding characteristics. These modifications provide medicinal chemists with an effective platform for establishing structure–activity relationships (SAR) and optimizing lead molecules.

II. STRUCTURAL FEATURES OF BENZIMIDAZOLE

The molecular formula of benzimidazole is $C_7H_6N_2$, with a molecular weight of approximately 118.14 g/mol. It contains an imidazole ring fused to a benzene ring. The presence of two nitrogen atoms gives the molecule distinctive acid–base and hydrogen-bonding properties. One nitrogen is pyridine-like and possesses a lone pair that can participate in protonation and coordination, whereas the other nitrogen is pyrrole-like and contributes to the aromatic π-electron system. The resulting electronic arrangement contributes to the stability of the heteroaromatic ring.

The benzimidazole nucleus possesses several positions that can be chemically modified. Among these, N-1, C-2, and the benzene-ring positions are particularly important for medicinal chemistry. Modification at these positions allows researchers to tune the biological and physicochemical properties of the resulting compounds.

Table 1. General Structural Characteristics of Benzimidazole

Property	Description
Chemical class	Aromatic nitrogen-containing heterocycle
Ring system	Bicyclic fused heterocycle
Structural components	Benzene fused with imidazole
Molecular formula	$C_7H_6N_2$
Molecular weight	~118.14 g/mol
Number of nitrogen atoms	Two
Aromatic character	Aromatic and π-electron conjugated
Important modification sites	N-1, C-2, C-5, C-6 and other benzene-ring positions
Medicinal chemistry importance	Privileged pharmacophore
Major applications	Drug discovery, medicinal chemistry and pharmaceutical research

III. BENZIMIDAZOLE AS A PRIVILEGED PHARMACOPHORE

A privileged scaffold is a molecular framework capable of interacting with multiple biological targets and being incorporated into structurally diverse compounds while retaining biological activity. Benzimidazole fulfills many of the characteristics expected from such a scaffold. Its planar aromatic structure enables π–π interactions with aromatic residues in proteins, while the nitrogen atoms provide opportunities for hydrogen bonding and electrostatic interactions. Appropriate substitution can additionally improve hydrophobic interactions and molecular recognition.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429 Volume 6, Issue 5, September 2026 Impact Factor: 8.2

The versatility of the benzimidazole nucleus has resulted in its incorporation into numerous biologically active molecules. Structural modifications can influence receptor binding, enzyme inhibition, membrane permeability, metabolic stability, and overall pharmacological activity. Thus, systematic modification of the benzimidazole ring represents an important strategy for lead optimization.

Table 2. Important Structural Positions of Benzimidazole

Position	Structural significance	Common modification
N-1	Influences basicity, lipophilicity and molecular recognition	Alkylation, arylation, acylation
C-2	Major site for structural diversification and biological activity	Aryl, heteroaryl, alkyl, carbonyl and other groups
C-3	Part of the imidazole framework and electronic system	Limited direct substitution
C-4	Benzene-ring position influencing electronic properties	Halogenation and other substitutions
C-5	Frequently exploited for optimization of biological activity	Halogen, alkoxy, amino, nitro and other groups
C-6	Useful for modifying steric and electronic properties	Alkyl, halogen, alkoxy and related groups
C-7	Can influence overall molecular conformation and properties	Various substituents

IV. SYNTHETIC IMPORTANCE OF BENZIMIDAZOLE

The synthesis of benzimidazole derivatives has become an important area of research in synthetic organic chemistry. Conventional synthetic routes generally involve the condensation of o-phenylenediamine with aldehydes, ketones, carboxylic acids, esters, nitriles, or related carbonyl compounds, depending on the desired substitution pattern. A variety of catalysts and reaction conditions have been investigated to improve product yield, reaction time, selectivity, and environmental compatibility.

Table 3. Representative Synthetic Approaches to Benzimidazole Derivatives

Synthetic approach	Typical starting materials	Major advantage
Condensation reaction	o-Phenylenediamine + aldehydes	Simple and widely applicable
Oxidative cyclization	o-Phenylenediamine + alcohol/aldehyde derivatives	Useful for diverse substitution
Reaction with carboxylic acids	o-Phenylenediamine + carboxylic acids	Provides 2-substituted derivatives
Multicomponent synthesis	Multiple simple starting materials	Molecular diversity and efficiency
Microwave-assisted synthesis	Diamines + carbonyl compounds	Short reaction time
Ultrasound-assisted synthesis	Diamines + carbonyl compounds	Mild and efficient conditions
Photocatalytic synthesis	Suitable diamines/carbonyl precursors	Potentially sustainable approach
Solvent-free synthesis	Solid or neat reactants	Reduced solvent consumption
Green catalytic synthesis	Catalysts with environmentally preferable conditions	Reduced environmental impact

V. BENZIMIDAZOLE DERIVATIVES AND MEDICINAL CHEMISTRY

The importance of benzimidazole derivatives in medicinal chemistry arises from their ability to interact with diverse biological targets. The introduction of suitable substituents can dramatically change the biological properties of the parent scaffold. Consequently, numerous benzimidazole derivatives have been investigated as potential therapeutic agents.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 5, September 2026
Impact Factor: 8.2

Benzimidazole-containing compounds have demonstrated a wide range of pharmacological activities, including antimicrobial, antiviral, anticancer, anti-inflammatory, analgesic, anti-ulcer, antidiabetic, anthelmintic, antihypertensive, and antioxidant activities. Their biological effects are strongly dependent on the substitution pattern, molecular conformation, electronic characteristics, lipophilicity, and ability to establish interactions with specific biological targets.

Table 4. Major Pharmacological Activities Reported for Benzimidazole Derivatives

Pharmacological activity	Possible biological relevance	Importance
Antimicrobial	Interaction with microbial enzymes/cellular processes	Development of antibacterial and antifungal agents
Antiviral	Interference with viral enzymes or replication processes	Potential antiviral drug development
Anticancer	Modulation of enzymes, receptors or signaling pathways	Development of anticancer candidates
Anti-inflammatory	Modulation of inflammatory mediators	Treatment of inflammatory disorders
Analgesic	Reduction of pain-associated pathways	Development of pain-relieving agents
Anti-ulcer	Modulation of gastric acid or protective mechanisms	Gastroprotective applications
Antidiabetic	Modulation of enzymes and metabolic pathways	Metabolic disease research
Anthelmintic	Activity against parasitic organisms	Treatment of helminthic infections
Antioxidant	Scavenging or modulation of oxidative processes	Protection against oxidative stress
Antihypertensive	Modulation of vascular or enzymatic pathways	Cardiovascular research

VI. STRUCTURE–ACTIVITY RELATIONSHIP OF BENZIMIDAZOLE DERIVATIVES

Structure–activity relationship studies are essential for understanding how chemical modifications influence the biological activity of benzimidazole derivatives. The nature and position of substituents can alter molecular polarity, hydrogen-bonding capacity, electron density, steric properties, lipophilicity, and target affinity.

Substitution at the C-2 position is particularly important because modifications at this site can substantially affect biological activity and molecular recognition. Substituents containing aromatic or heteroaromatic groups may increase hydrophobic interactions with target binding sites, whereas polar functional groups can improve hydrogen-bonding interactions and aqueous solubility. Similarly, substitution on the benzene ring can alter electronic distribution and physicochemical characteristics.

N-1 substitution provides another important route for optimizing the pharmacological profile. Depending on the substituent, N-1 modification can influence molecular conformation, lipophilicity, metabolic stability, and target interactions. Therefore, systematic variation at N-1, C-2, and benzene-ring positions is commonly employed during medicinal chemistry optimization.

Table 5. Relationship Between Structural Modification and Molecular Properties

Structural modification	Potential effect	Medicinal chemistry significance
N-alkylation	Increased lipophilicity/conformational changes	May alter membrane permeability and target binding
N-arylation	Increased aromatic surface	May enhance hydrophobic and π-interactions
C-2 aryl substitution	Increased molecular size and hydrophobicity	Frequently useful for biological activity optimization

Copyright to IJARSCT
www.ijarsct.co.in
DOI: 10.48175/IJARSCT-40603

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429 Volume 6, Issue 5, September 2026 Impact Factor: 8.2

Halogen substitution	Alters electronic and lipophilic properties	Can improve potency and metabolic properties
Methoxy/alkoxy substitution	Changes polarity and electron density	Useful for tuning target interactions
Amino substitution	Increases hydrogen-bonding potential	May improve interaction with biological targets
Nitro substitution	Strong electron-withdrawing effect	Can substantially modify electronic properties
Heteroaryl substitution	Introduces additional heteroatoms	Provides new hydrogen-bonding and recognition possibilities

VII. THERAPEUTIC RELEVANCE

The successful incorporation of the benzimidazole scaffold into several pharmacologically important molecules has established its relevance in modern drug discovery. The nucleus can mimic or interact with biologically important heteroaromatic structures and can be tailored to achieve desired target interactions. Its ability to accommodate a wide range of substituents makes it particularly suitable for scaffold hopping, molecular hybridization, and lead optimization.

Table 6. Selected Therapeutic Areas Associated with Benzimidazole-Based Compounds

Therapeutic area	Research significance
Gastrointestinal disorders	Development of acid-suppressing and gastroprotective agents
Parasitic infections	Development of anthelmintic agents
Bacterial and fungal infections	Search for novel antimicrobial agents
Viral infections	Identification of compounds affecting viral replication
Cancer	Development of enzyme- and signaling-targeted molecules
Inflammatory diseases	Investigation of inflammatory pathway modulation
Metabolic disorders	Exploration of enzyme-targeted antidiabetic compounds

VIII. RECENT RESEARCH TRENDS

Current research on benzimidazole chemistry is increasingly directed toward the development of multifunctional molecules and environmentally responsible synthetic methods. Green chemistry principles are being incorporated into benzimidazole synthesis through the use of safer solvents, recyclable catalysts, solvent-free conditions, microwave irradiation, ultrasound, photocatalysis, and other energy-efficient techniques.

Another important research direction is the synthesis of benzimidazole-based polyheterocyclic systems. Molecular fusion or hybridization of benzimidazole with other heterocyclic pharmacophores can generate structurally complex molecules with enhanced biological potential. Such compounds may exhibit synergistic pharmacological effects because different heterocyclic fragments can interact with multiple biological targets.

The application of computational chemistry, molecular docking, quantitative structure–activity relationship (QSAR) studies, artificial intelligence-assisted drug design, and molecular dynamics simulations has further expanded the possibilities for benzimidazole-based drug discovery. These approaches facilitate the prediction of target interactions and help researchers prioritize promising compounds before experimental evaluation.

Table 7. Emerging Strategies in Benzimidazole Research

Research strategy	Purpose
Green synthesis	Minimize environmental impact
Microwave-assisted synthesis	Accelerate chemical transformations
Ultrasound-assisted synthesis	Improve reaction efficiency under mild conditions
Photocatalysis	Enable alternative and potentially sustainable reaction pathways
Multicomponent reactions	Generate structural diversity efficiently
Molecular hybridization	Combine pharmacophores to improve biological activity

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 5, September 2026
Impact Factor: 8.2

Molecular docking	Predict ligand–target interactions
QSAR analysis	Correlate molecular structure with biological activity
Molecular dynamics	Examine stability of ligand–target complexes
AI-assisted drug design	Support prediction and optimization of potential drug candidates

IX. SYNTHETIC APPROACHES FOR BENZIMIDAZOLE DERIVATIVES

Benzimidazole and its derivatives have attracted considerable attention because of their diverse biological activities and broad applications in medicinal and pharmaceutical chemistry. Several synthetic methodologies have been developed for the preparation of benzimidazole derivatives. One of the classical approaches involves the direct condensation of o-phenylenediamine (OPD; benzene-1,2-diamine) with formic acid. Although this method provides a straightforward route to the benzimidazole nucleus, it is associated with certain limitations, including moderate or low product yields, prolonged reaction times, harsh reaction conditions, and tedious work-up procedures. These drawbacks have encouraged researchers to develop more efficient, economical, and environmentally benign synthetic protocols.

Among the reported methodologies, the condensation of benzene-1,2-diamine (OPD) with various aldehydes has emerged as an important and versatile route for synthesizing substituted benzimidazoles. In a typical procedure, OPD reacts with different aromatic or aliphatic aldehydes in ethanol at room temperature in the presence of a suitable catalyst. This approach offers several advantages, such as operational simplicity, relatively mild reaction conditions, shorter reaction times, and satisfactory to excellent yields.

Various homogeneous and heterogeneous catalysts have been investigated to improve the efficiency of this transformation. In particular, inexpensive and readily available catalytic systems have demonstrated considerable potential as alternatives to costly catalysts. Heterogeneous catalysts such as Pt- and TiO_2-based catalytic systems have been reported to facilitate the formation of benzimidazole derivatives efficiently. These catalytic methods generally afford the desired products in good yields of approximately 72–88% within 3–7 h, demonstrating improved catalytic efficiency compared with several conventional methodologies.

Table: 8 Synthetic Approaches for Benzimidazole Derivatives

Scheme	Starting Materials	Catalyst / Reagent	Reaction Conditions	Product / Benzimidazole Derivative	Key Features
Scheme 1	Benzene-1,2-diamine (o-phenylenediamine) + Acetaldehyde	Pt; TiO_2	Ethanol, reflux; reported reaction time 3–7 h	2-Methyl-1H-benzimidazole	Simple and inexpensive catalytic system; good efficiency and comparatively high yields (72–88%); shorter reaction time
Scheme 2	Benzene-1,2-diamine + Benzoic acid	—	Reflux, 24 h	2-Phenyl-1H-benzimidazole	o-Phenylenediamine reacts with carboxylic acids to afford 2-substituted benzimidazoles, generally in good yields
Scheme 3 (Phillips Method)	Benzene-1,2-diamine + Acetic acid	Concentrated HCl	Heating / reflux	2-Methyl-1H-benzimidazole	Classical Phillips condensation; widely used method involving condensation of o-phenylenediamine with carboxylic acids or their

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 5, September 2026
Impact Factor: 8.2

					derivatives
Scheme 4	4-Methylbenzene-1,2-diamine + Acetaldehyde	CH_3CN; YCl_3	Microwave irradiation; room temperature conditions as reported	2,5-Dimethyl-1H-benzimidazole	High yield, clean reaction profile, inexpensive catalyst/system, and relatively green methodology
Scheme 5	4-Methylbenzene-1,2-diamine + Aliphatic aldehydes	DBSA	Reaction conditions as reported	Substituted benzimidazole derivatives	DBSA acts as a potent and less-toxic catalyst; applicable to the synthesis of different substituted benzimidazoles
Scheme 6	Benzene-1,2-diamine + Methyl 2,2,2-trichloroethanimidate	AcOH	Room temperature	2-(Trichloromethyl)-1H-benzimidazole	Reaction proceeds at room temperature; provides 2-trichloromethyl-substituted benzimidazole
Scheme 7	o-Phenylenediamine + Acetic acid / aromatic acids	NH_4Cl	EtOH; 80–90 °C	2-Methyl-1H-benzimidazole / substituted benzimidazoles	Green and economical approach using ammonium chloride as catalyst
Scheme 8	o-Phenylenediamine + Carboxylic acids	Zeolite HY	Solvent-free; microwave irradiation; 5–9 min; 160–560 W	2-Substituted benzimidazoles	Rapid microwave-assisted synthesis; solvent-free conditions; Zeolite HY catalyst; suitable for aromatic, aliphatic and heterocyclic carboxylic acids
Scheme 9	o-Phenylenediamine + Carboxylic acids	Silica gel	Solvent-free; microwave irradiation; 5–9 min; 160–560 W	2-Substituted benzimidazoles	Short reaction time, solvent-free conditions, and simple heterogeneous catalyst
Scheme 10	o-Phenylenediamine + Carboxylic acids	Alumina	Solvent-free; microwave irradiation; 5–9 min; 160–560 W	2-Substituted benzimidazoles	Rapid and economical microwave-assisted methodology using alumina as a heterogeneous catalyst

Copyright to IJARSCT
www.ijarsct.co.in
DOI: 10.48175/IJARSCT-40603

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 5, September 2026
Impact Factor: 8.2

Table: 9 Comparative Summary of Synthetic Methods

Method / Scheme	Catalyst / Promoter	Solvent	Temperature	Time	Major Advantage
Direct condensation	—	—	Reflux / heating	Relatively long	Straightforward synthesis from o-phenylenediamine and formic acid
Scheme 1	Pt/TiO_2	Ethanol	Reflux	3–7 h	Good yield and efficient catalytic system
Scheme 2	—	—	Reflux	24 h	Broad reaction with carboxylic acids
Scheme 3 – Phillips method	Conc. HCl	—	Heating	Long	Classical and well-established methodology
Scheme 4	YCl_3 / CH_3CN	As reported	RT / MW	Short	Clean reaction and good yield
Scheme 5	DBSA	As reported	As reported	—	Less-toxic catalyst and useful for substituted derivatives
Scheme 6	AcOH	—	Room temperature	Short	Mild reaction conditions
Scheme 7	NH_4Cl	Ethanol	80–90 °C	—	Green, economical and simple
Scheme 8	Zeolite HY	Solvent-free	Microwave	5–9 min	Very rapid and environmentally favorable
Scheme 9	Silica gel	Solvent-free	Microwave	5–9 min	Simple heterogeneous catalyst and short reaction time
Scheme 10	Alumina	Solvent-free	Microwave	5–9 min	Rapid, economical and solvent-free synthesis

X. CONCLUSION

Benzimidazole and its derivatives represent an important class of nitrogen-containing heterocyclic compounds with considerable significance in medicinal chemistry and pharmaceutical research. The literature survey demonstrates that a wide variety of synthetic approaches have been developed for the preparation of benzimidazole derivatives, mainly through the condensation of o-phenylenediamine with aldehydes, carboxylic acids, and their derivatives. Classical methods, including direct condensation and the Phillips method, are simple and reliable; however, they are often associated with prolonged reaction times, high temperatures, strong acidic conditions, moderate yields, and tedious work-up procedures.

Recent synthetic strategies have focused on overcoming these limitations through the use of heterogeneous catalysts, microwave irradiation, solvent-free conditions, and relatively environmentally benign catalytic systems. Catalysts such as TiO_2, Pt, YCl_3, DBSA, NH_4Cl, zeolite HY, silica gel, and alumina have demonstrated their usefulness in improving reaction efficiency. In particular, microwave-assisted and solvent-free methodologies provide significant advantages, including rapid reaction rates, reduced solvent consumption, simplified procedures, and improved energy efficiency.

Overall, the development of greener, faster, economical, and more efficient synthetic methodologies has considerably enhanced the preparation of benzimidazole derivatives. Future research should focus on designing recyclable catalysts, employing safer solvents or completely solvent-free conditions, reducing energy consumption, and developing scalable synthetic protocols. Such advances may facilitate the efficient synthesis of structurally diverse benzimidazole derivatives with potential applications in pharmaceutical, biological, and materials research.

REFERENCES

1. Phillips M.A. The formation of 2-substituted benzimidazoles. Journal of the Chemical Society, 1928.
2. Grimmett M.R. Imidazole and Benzimidazole Synthesis. Academic Press, London, 1997.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 5, September 2026
Impact Factor: 8.2

3. Preston P.N. Benzimidazoles and related systems. In: The Chemistry of Heterocyclic Compounds. Wiley-Interscience, New York, 1981.
4. Ladenburg A. Untersuchungen über Benzimidazole. Berichte der Deutschen Chemischen Gesellschaft, 1877.
5. Weidenhagen R., Wegner H. Über die Darstellung von Benzimidazolen aus o-Phenylendiamin und Aldehyden. Berichte der Deutschen Chemischen Gesellschaft, 1930.
6. Tandon V.K., Yadav D.B., Maurya H.K., Chaturvedi A.K., Shukla P.K. Design, synthesis and biological evaluation of benzimidazole derivatives. Bioorganic & Medicinal Chemistry, 2006.
7. Heravi M.M., Bakhtiari K., Tehrani M.H., Daroogheha Z. One-pot synthesis of benzimidazole derivatives using heterogeneous catalysts. Catalysis Communications, 2007.
8. Bahrami K., Khodaei M.M., Nejati A. Synthesis of benzimidazole derivatives under mild and environmentally friendly conditions. Green Chemistry, 2009.
9. Bhatnagar I., George M.V. Oxidation of substituted benzimidazoles and related heterocyclic compounds. Tetrahedron, 1968.
10. Singhal S., Khanna P., Panda S.S., Khanna L. Recent trends in the synthesis of benzimidazoles from o-phenylenediamine via nanoparticles and green strategies using transition metal catalysts. Journal of Heterocyclic Chemistry, 2019, 56, 2702–2729.
11. Sharma J., Soni P.K., Bansal R., Halve A.K. Synthetic approaches towards benzimidazoles by the reaction of o-phenylenediamine with aldehydes using a variety of catalysts: A review. Current Organic Chemistry, 2018, 22, 2280–2299.
12. Acharjee A., Das P., Das D., et al. Synthetic approaches to benzimidazoles from o-phenylenediamine: A literature review. Journal of Saudi Chemical Society, 2017.
13. Rushi K., et al. A convenient one-pot synthesis of 2-substituted benzimidazoles. Journal of Molecular Catalysis A: Chemical, 2006, 245, 8–11.
14. Wang L., et al. DMP-catalyzed synthesis of 2-substituted benzimidazole derivatives. Chinese Chemical Letters, 2009, 20, 893–897.
15. Kovvuri J., et al. An efficient synthesis of 2-substituted benzimidazoles via photocatalytic condensation of o-phenylenediamines and aldehydes. ACS Combinatorial Science, 2016, 18, 644–650.
16. Ghosh S., Subba Reddy K. Catalytic synthesis of substituted benzimidazoles from o-phenylenediamines and aldehydes. Tetrahedron Letters, 2010.
17. Venkateswarlu Y., et al. Lanthanum chloride catalyzed one-pot synthesis of 2-substituted benzimidazoles. Journal of Molecular Catalysis A: Chemical, 2007.
18. Lin S., Yang L. A simple and efficient synthesis of benzimidazoles using air as oxidant. Tetrahedron Letters, 2005.
19. Srinivasulu D., et al. Zinc triflate catalyzed synthesis of 2-substituted benzimidazoles. Synthetic Communications, 2008.
20. Zhao J.A., et al. Recent catalytic approaches for the synthesis of benzimidazole derivatives. Chinese Chemical Letters, 2017, 28, 1539–1546.
21. Kovvuri J., et al. Rose Bengal photocatalyzed condensation of o-phenylenediamines and aldehydes for benzimidazole synthesis. ACS Combinatorial Science, 2016, 18, 644–650.
22. Selective synthesis of benzimidazoles from o-phenylenediamine and aldehydes promoted by supported gold nanoparticles. Nanomaterials, 2020, 10, 2405.
23. Recent achievements in the synthesis of benzimidazole derivatives. RSC Advances, 2023. DOI: 10.1039/D3RA05960J.
24. Amalraj S.D., Palapetta S.C., Harichandran G. A facile one-pot synthesis, computational and molecular docking studies of benzimidazole and benzothiazole compounds using Amberlite IRA-400-Cl resin as green/reusable catalyst. Journal of Molecular Structure, 2022, 1268, 133704.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 5, September 2026
Impact Factor: 8.2

25. Shaibuna M., Hiba K., Shebitha A.M., et al. Sustainable and selective synthesis of benzimidazole scaffolds using deep eutectic solvents. Current Research in Green and Sustainable Chemistry, 2022, 5, 100285.
26. Gavali L.V., Wadhawa G.C., Shivankar V.S., et al. Review on synthesis of benzimidazole using green protocol. World Journal of Pharmacy and Pharmaceutical Sciences, 2016, 5(9), 624–656.
27. Wadhawa G.C., Shivankar V.S., Patil S.S., Gaikwad Y.A., Satere A.V., Rode B., Gill C.H., Gavali L.V. Synthesis of 2-substituted benzimidazoles and 1,5-disubstituted benzodiazepines using alum. Rasayan Journal of Chemistry, 2017.
28. Gill C.H., Wadhawa G.C., Gavali L.V., Shivankar V.S., et al. Synthesis of benzimidazole using waste magnesium oxide. Research Journal of Pharmacology and Pharmacodynamics, 2018, 10(3), 103.
29. Wadhawa G.C., Gaikwad Y.A., Shivankar V.S. Red mud catalyzed synthesis of benzimidazole, benzothiazoles and quinoxalines. International Journal of Advanced Science and Technology, 2020, 29(9s), 5286–5297.
30. Wadhawa G.C., Shivankar V.S., Patil S.S., Gill C.H., Gavali L.V. Synthesis of benzimidazole derivatives under solvent-free condition. National Conference on Recent Trends in Chemistry, 2020.
31. Valvi A.K., Nayak S.S., Shivankar V.S., Wadhawa G.C. Nano catalyst derived from plant Polyzygus tuberosus: A suitable catalyst for efficient and green synthesis of benzimidazole derivatives. Materials Today: Proceedings, 2022.
32. Wadhawa G.C., Valvi A.K., Mohite R.D., Patil D.D., Patil B., Gavit H.J. An organo-heterogeneous catalyst for synthesis of benzimidazole derivatives using nanoparticles synthesized from plant extract and supported on Montmorillonite K10. Materials Today: Proceedings, 2022, 58, 764–768.
33. Sasane N.A., Valvi A.K., Meshram G.S., Wadhawa G.C. Synthesis of benzimidazole derivatives in an aqueous medium and reflux conditions catalyzed by glycine at pH 5.2 under microwave approach. Materials Today: Proceedings, 2022, 58, 769–772.
34. Valvi A.K., Gavit H.J., Nayak S.S., Shivankar V.S., Wadhawa G.C. Synthesis of benzimidazole and benzothiazole derivatives using reusable waste stem of Trigonella foenum-graecum assisted zinc sulphide nanoparticles: A green and efficient solid acid catalyst. Materials Today: Proceedings, 2023, 73, 481–486.
35. Wadhawa G.C., Patil P.A., Patil P., Shivankar V.S., Nayak S.S., Gaikar P.S., Patil B.M. One-pot green synthesis of ZnS nanoparticles and its application for degradation of organic dye and synthesis of 2-benzimidazole derivatives. AIP Conference Proceedings, 2024.