IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 4, November 2026
Impact Factor: 8.2

Herb-Drug Interactions Documented in the Literature and their Clinical Significances

Dr. Bhagavan M. Kale[1], Drx. Mayuri Mokal[2], Umesh G Thale[3], Sachin J Ghodke[4], Kunal P Shirabhaye[5]

[1]Assistant Professor, Shri Goraksha College of Pharmacy and Research Centre, Khamgaon, Maharashtra, India.
[2]Principal, Shree Goraksha College of Pharmacy and Research Centre, Khamgaon, Maharashtra, India
[3,4,5]Student, Department of Pharmacy, Shri Goraksha College of Pharmacy and Research Centre, Khamgaon

Abstract : *Across the globe, patients managing chronic lifestyle conditions are increasingly turning to Complementary and Alternative Medicine (CAM)—especially herbal remedies and nutraceuticals— alongside their prescribed medications. Because many people assume that anything "natural" is inherently harmless, they often combine herbal supplements with conventional drugs without telling their doctors. This review brings together published clinical and experimental evidence on herb-drug interactions (HDIs) across four major lifestyle conditions: diabetes mellitus, psychiatric disorders, arthropathies, and cardiovascular disease. Specifically, we examine the pharmacokinetic and pharmacodynamic mechanisms behind interactions involving Trigonella foenum-graecum (fenugreek), Coccinia indica (ivy gourd), Panax ginseng, Allium sativum (garlic), Glycyrrhiza glabra (liquorice), and Allium cepa (onion). Reducing preventable adverse events will require stronger regulatory oversight, practical updates to medical education, and closer collaboration between traditional and modern healthcare systems.*

Keywords: Herb-drug interactions, Complementary and Alternative Medicine, Lifestyle disorders, Pharmacokinetics, Clinical significance, Integrative medicine

I. INTRODUCTION

Over the past two decades, the use of Complementary and Alternative Medicine (CAM) has grown steadily in both developed and developing countries (Eisenberg et al., 1998). Among the many alternative therapies available today, herbal medicine—consumed as home remedies, dietary supplements, and nutraceuticals—has gained particular traction. Easy over-the-counter access, lower out-of-pocket costs, rising disposable incomes, and the widespread belief that plant-based products are completely safe have all fueled this trend.

Yet clinical data tell a more cautious story. Studies from developed nations such as the United States have documented substantial numbers of adverse drug events and fatalities each year (Lazarou, 1998) when patients pair herbal supplements with standard therapies for chronic conditions like diabetes, hypertension, chronic stress, and joint disorders. Many of these complications stem from overlooked herb-drug, food-drug, drug-drug, and disease-drug interactions.

Clinicians and healthcare accreditation bodies can no longer afford to treat these interactions as rare edge cases. In routine practice, herb-drug interactions often go unrecognized or are dismissed as minor. Older adults, children, and immunocompromised patients are particularly vulnerable to serious side effects. Evaluating these risks is also inherently difficult: a single medicinal herb contains dozens of active phytochemicals rather than one isolated molecule, and researchers still lack a uniform, standardized framework to map every interaction pathway in the human body.

Causes of Herb-Drug Interactions

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 4, November 2026
Impact Factor: 8.2

The assumption that "natural" equals safe: Most patients view herbal teas, powders, and capsules as gentle and risk-free. As a result, they rarely mention these products during medical consultations, even when taking daily prescription drugs.

Gaps in pharmacokinetic and pharmacodynamic data: Unlike approved pharmaceuticals, many herbal products lack thorough human studies showing how the body absorbs, metabolizes, and clears them—particularly their effects on cytochrome P450 (CYP450) enzymes and Pglycoprotein transporters—making clinical outcomes hard to anticipate.

Complex and variable chemical profiles: Whole herbs contain a mixture of bioactive constituents. While some compounds act as natural buffers, concentrated commercial extracts can deliver high doses of potent, unstable molecules capable of triggering unexpected physiological reactions.

Limited clinical trials on dosing: Standardized dosage ranges, safe durations of use, and clear contraindications have not been established for many traditional formulations.

Inconsistent manufacturing and quality control: Commercial herbal products vary widely in purity and strength. Misidentified botanical species, unapproved plant substitutes, undeclared synthetic adulterants, and contamination with heavy metals or pesticide residues all heighten the danger of adverse reactions.

Types of Herb-Drug Interactions

Clinicians generally classify herb-drug interactions based on how they alter drug behavior in the body and what happens to the patient:

Synergistic and Antagonistic Interactions

- **Synergistic (Combined Effect):** The herb and the prescribed drug act in the same direction, amplifying the overall physiological response. While this can occasionally boost therapeutic benefit, it frequently pushes drug activity into a toxic range.
- **Antagonistic (Opposing Effect):** The herb interferes with absorption, speeds up drug clearance, or blocks receptor activity, leaving the patient with subtherapeutic drug levels and poor disease control.

Positive and Negative Interactions

- **Positive (Helpful):** Co-administration produces a favorable clinical outcome, such as improved drug bioavailability or fewer treatment-related side effects.
- **Negative (Harmful):** The combination leads to clinical harm, ranging from treatment failure to acute emergencies such as severe hypoglycemia or internal bleeding.

OBJECTIVES

This review aims to identify, explain, and help clinicians anticipate harmful or counterproductive biological interactions that arise when herbal remedies are taken alongside conventional prescription drugs, with particular focus on the safe management of chronic lifestyle disorders.

METHODS

We conducted a structured review of published literature examining clinically relevant herb-drug interactions in chronic lifestyle diseases. Searches were carried out across major biomedical and traditional medicine databases, including DHARA, MEDLINE, Wiley Online, Scopus, OvidSP, CINAHL, and ResearchGate. We screened 150 initial records, selecting clinical case reports, experimental investigations, and peer-reviewed reviews that documented interactions between widely used medicinal herbs and standard drugs prescribed for metabolic, psychiatric, musculoskeletal, and cardiovascular conditions.

RESULTS AND CLINICAL EVALUATION

From the 150 screened records—spanning clinical reports, experimental trials, editorials, and review papers—we grouped the most clinically notable interactions into four major lifestyle disease categories.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 4, November 2026
Impact Factor: 8.2




Figure 1: Overview of pharmacokinetic and pharmacodynamic mechanisms underlying herb-drug interactions across major lifestyle disorders, illustrating the interplay between botanical supplements and conventional medications.

1. Evaluating Herb-Drug Interactions in Diabetes

Diabetes mellitus remains one of the most widespread metabolic and lifestyle-related conditions worldwide. In an early ethnobotanical survey, Bever and Zahnd (1979) cataloged roughly 400 plant species with documented blood-glucose-lowering activity. Today, patients frequently supplement their diabetes prescriptions with bitter gourd (*Momordica charantia*), ivy gourd, jamun (*Syzygium cumini*), fenugreek, and *Gymnema sylvestre*.

- **Fenugreek (*Trigonella foenum-graecum*):** Known widely as *methi*, fenugreek seeds encourage pancreatic insulin release and slow gastric emptying because of their dense soluble fiber content. Although helpful for lowering fasting glucose and serum cholesterol, taking fenugreek alongside insulin or oral hypoglycemics (such as sulfonylureas or metformin) can trigger additive, clinically significant hypoglycemia. Its thick mucilaginous fiber can also bind to or delay the gut absorption of co-administered oral drugs.
- **Ivy Gourd (*Coccinia indica*):** Consumed both as a culinary vegetable and a classical Ayurvedic remedy, *C. indica* displays insulin-mimetic activity. Patients using ivy gourd extracts together with standard antidiabetic medications need regular blood glucose monitoring to avoid unexpected hypoglycemic episodes.

2. Assessment of Herb-Drug Interactions in Psychosis and Mood Disorders

Rapid urbanization and shifting socioeconomic pressures have contributed to rising rates of chronic stress, anxiety, and depressive disorders, driving higher prescription volumes for antidepressants and antipsychotics. Demographic projections had already estimated that India's older adult population would reach 113 million by 2016 (Chaubey and Aarti, 1999), pointing to a large cohort managing psychiatric symptoms alongside multiple chronic medications. At the same time, many individuals reach for traditional Indian and Chinese herbs—such as Ashwagandha and Ginseng—to ease stress, sharpen memory, and stabilize mood (Archana & Namasivayam, 1998).

- **Ginseng (*Panax ginseng* and *P. quinquefolium*):** Combining ginseng with tricyclic antidepressants (TCAs) or monoamine oxidase inhibitors (MAOIs) has been linked to insomnia, headaches, and mania-like episodes, and it can lower the seizure threshold in susceptible patients.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429 Volume 6, Issue 4, November 2026 Impact Factor: 8.2

- **Ashwagandha (*Withania somnifera*):** Valued for its adaptogenic and stress-relieving properties (Archana & Namasivayam, 1998), ashwagandha exhibits GABA-mimetic activity that can deepen central nervous system (CNS) depression when taken with benzodiazepines, barbiturates, or other sedatives.

3. Assessment of Herb-Drug Interactions in Arthropathies

According to the World Health Organization, more than 150 distinct conditions affect the joints and musculoskeletal system. While non-steroidal anti-inflammatory drugs (NSAIDs) and analgesics such as aspirin and paracetamol remain mainstays of pain control, prolonged use carries well-known risks of gastric ulceration, kidney injury, and liver toxicity. Seeking gentler long-term relief, many arthritis patients turn to botanical remedies such as garlic, guggulu (*Commiphora wightii*), and boswellia (*Boswellia serrata*).

- **Garlic (*Allium sativum*):** Alongside its everyday culinary role and topical use as a soothing paste for sore joints, concentrated oral garlic supplements inhibit platelet aggregation. Taking high-dose garlic together with aspirin, other NSAIDs, or prescription anticoagulants compounds antiplatelet activity and raises the risk of prolonged bleeding and gastrointestinal hemorrhage.

4. Assessment of Herb-Drug Interactions in Cardiovascular Diseases

Cardiovascular disease continues to be the leading cause of death globally, with hypertension alone responsible for roughly 9.4 million deaths—or 16.5% of all mortality—each year (Alwan, 2011). Patients living with coronary artery disease, cerebrovascular disease, or rheumatic heart conditions often turn to herbal supplements to help manage blood pressure, body weight, and cholesterol. These botanicals frequently exert cardiotonic, vasodilatory, or diuretic effects. However, serious complications can emerge when they are paired with prescription cardiac drugs, anticoagulants, or antiplatelet therapy (Cheng, 2000; Holbrook et al., 2005; Dunn and Macaulay, 2011).

- **Liquorice (*Glycyrrhiza glabra*):** Frequently used in traditional lozenges and decoctions for coughs and sore throats, liquorice contains glycyrrhizin, a compound that mimics aldosterone. Excess intake promotes sodium and water retention while depleting serum potassium (hypokalemia). Low potassium levels sharply increase the danger of digoxin toxicity and cardiac arrhythmias, while fluid retention directly counteracts antihypertensive therapy (Cheng, 2000).
- **Onion (*Allium cepa*):** Beyond its place in the daily diet, onion extracts possess natural antioxidant and antiplatelet properties. When consumed in concentrated supplement form alongside warfarin, clopidogrel, or aspirin, *Allium cepa* can amplify anticoagulant activity and elevate bleeding risk.

Table 1: Summary of Clinically Significant Herb-Drug Interactions in Lifestyle Disorders

Lifestyle Disorder	Herb (Botanical Name)	Conventional Drug Class	Mechanism & Clinical Outcome
Diabetes Mellitus	Fenugreek (*Trigonella foenumgraecum*)	Sulfonylureas, Insulin, Metformin	Additive blood-glucose lowering; soluble fiber may delay oral drug absorption.
Psychiatric Disorders	Ginseng (*Panax ginseng*)	Tricyclic Antidepressants (TCAs), MAOIs	Lowered seizure threshold; potential for insomnia, agitation, or mania-like symptoms.
Psychiatric Disorders	Ashwagandha (*Withania somnifera*)	Benzodiazepines, CNS Depressants	Enhanced sedation through additive GABAergic activity.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 4, November 2026
Impact Factor: 8.2

Arthropathies	Garlic (*Allium sativum*)	Aspirin, NSAIDs	Additive platelet inhibition; higher likelihood of gastrointestinal bleeding.
Cardiovascular Disease	Liquorice (*Glycyrrhiza glabra*)	Digoxin, Antihypertensives, Diuretics	Glycyrrhizin-driven potassium loss; increased digoxin toxicity and blunted blood pressure control.
Cardiovascular Disease	Onion (*Allium cepa*)	Warfarin, Antiplatelet agents	Amplified anticoagulant and antiplatelet response; heightened bleeding risk.

II. CONCLUSION

In both industrialized and developing healthcare settings, growing numbers of patients combine traditional herbal remedies with modern prescription medications—often unaware that natural products can alter drug safety and effectiveness. Reducing these preventable risks calls for practical updates to medical and pharmacy curricula, active engagement from healthcare accreditation bodies, and stronger enforcement of quality standards under laws such as the Drugs and Cosmetics Act and the Consumer Protection Act. Dedicated pharmacovigilance programs and research centers are equally needed to evaluate traditional formulations rigorously while respecting their clinical heritage.

As Hippocrates famously wrote, “Life is short and the art long; the occasion fleeting; experience fallacious; and judgment difficult.” Bridging the divide between traditional healing systems and contemporary pharmacology is essential if Integrative Medicine is to deliver truly safe, patientcentered care. Bringing both traditions under the shared discipline of evidence-based science—and building what Dr. Ramashekhar described as a “golden triangle” linking traditional medicine, modern medicine, and modern science—will require sustained cooperation among clinicians, researchers, industry leaders, and policymakers.

REFERENCES

1. Archana, R., & Namasivayam, A. (1998). Antistressor effect of *Withania somnifera*. *Journal of Ethnopharmacology*, 64(1), 91–93.
2. Alwan, A. (2011). *Global status report on non-communicable diseases 2010*. World Health Organization.
3. Bever, B. O., & Zahnd, G. R. (1979). Plants with oral hypoglycaemic action. *Quarterly Journal of Crude Drug Research*, 17(3–4), 139–196.
4. Chaubey, P. C., & Aarti. (1999). Demographic transition and health care of the elderly in India. *Indian Journal of Gerontology*, 13, 1–10.
5. Dunn, S. P., & Macaulay, T. E. (2011). Drug interactions with newer antiplatelet agents and herbal supplements. *Current Cardiology Reports*, 13(5), 415–422.
6. Eisenberg, D. M., Davis, R. B., Ettner, S. L., Appel, S., Wilkey, S., Van Rompay, M., & Kessler, R. C. (1998).
7. Trends in alternative medicine use in the United States, 1990–1997. *JAMA*, 280(18), 1569–1575.
8. Holbrook, A. M., Pereira, J. A., Labiris, R., McDonald, H., Douketis, J. D., Crowther, M., & Wells, P. S. (2005). Systematic overview of warfarin and its drug and food interactions. *Archives of Internal Medicine*, 165(10), 1095–1106.
9. Lazarou, J., Pomeranz, B. H., & Corey, P. N. (1998). Incidence of adverse drug reactions in hospitalized patients: A meta-analysis of prospective studies. *JAMA*, 279(15), 1200–1205.
10. Cheng, T. O. (2000). Herbal interactions with cardiac drugs. *Archives of Internal Medicine*, 160(6), 870– 871.