IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 5, November 2026
Impact Factor: 8.2

Evaluating the Impact of Strategic HR Intervention on Compassion Fatigue and Retention among Frontline Nursing Staff

Ms. Sulagna Panda[1], Prof. (Dr.) Bhawna Sharma Padroo[2], Dr. Sameer Kulkarni[3], Prof. (Dr.) Rajat Gera[4]
Research Scholar , Amity Business School, Amity University Mumbai[1]
Professor , Amity Business School & Director -IAD, Amity University Mumbai[2]
Associate Professor, Amity Business School, Amity University Mumbai[3]
Professor & Director, Amity Business School, Amity University Mumbai4
sulagna.panda@s.amity.edu, bsharma@mum.amity.edu
sakulkarni@mum.amity.edu, rgera@mum.amity.edu

Abstract: *Frontline nursing staff play a critical role in delivering quality healthcare services; however, continuous exposure to patient suffering, emotional distress, high workloads, and demanding working conditions often results in compassion fatigue. This condition adversely affects nurses' psychological well-being, professional performance, job satisfaction, and intention to remain in healthcare organisations. The present study aims to evaluate the impact of strategic Human Resource (HR) interventions on reducing compassion fatigue and improving employee retention among frontline nursing staff. It examines the effectiveness of organisational initiatives such as employee assistance programmes, psychological counselling, stress management training, flexible scheduling, supportive leadership, recognition, and professional development opportunities. The study further investigates the relationship between HR interventions, emotional well-being, organisational commitment, and nurses' retention intentions. By identifying effective HR practices, the research seeks to develop a sustainable human resource management framework that promotes workforce resilience, enhances employee satisfaction, and strengthens nursing workforce stability.*

Keywords: Compassion Fatigue, Strategic HR Interventions, Frontline Nursing Staff, Employee Retention, Healthcare Workforce Well-being

I. INTRODUCTION

The healthcare sector depends significantly on the dedication, professional competence, and emotional commitment of frontline nursing staff. Nurses frequently work in demanding environments where they encounter patient suffering, critical medical conditions, emergencies, and emotionally challenging situations. Continuous exposure to such circumstances can result in compassion fatigue, which refers to the physical, emotional, and psychological exhaustion experienced by healthcare professionals due to prolonged engagement with individuals experiencing pain and trauma. Compassion fatigue may negatively influence nurses' job satisfaction, work performance, interpersonal relationships, and overall quality of patient care.

In recent years, healthcare organisations have increasingly recognised the importance of strategic Human Resource Management (HRM) in addressing workforce-related challenges. Traditional HR practices focusing primarily on recruitment, compensation, and performance evaluation are no longer sufficient to address the complex psychological and emotional requirements of nursing professionals. Organisations must adopt employee-centred HR interventions that promote emotional resilience, psychological safety, work-life balance, and professional development.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429 Volume 6, Issue 5, November 2026 Impact Factor: 8.2

Strategic HR interventions such as counselling services, employee assistance programmes, stress management workshops, adequate staffing, flexible work schedules, supportive supervision, and recognition programmes can play an important role in reducing occupational stress and compassion fatigue. These initiatives may also strengthen organisational commitment, improve job satisfaction, and reduce employee turnover intentions.

The present study evaluates the impact of strategic HR interventions on compassion fatigue and retention among frontline nursing staff. It seeks to understand how organisational policies and employee welfare initiatives influence nurses' psychological well-being and willingness to continue working within healthcare institutions. The study ultimately aims to identify effective HR strategies that contribute to a healthier, more resilient, and sustainable nursing workforce.

Significance of the Study

The study is significant because it addresses the growing concern of compassion fatigue and employee turnover among frontline nursing professionals. It highlights the role of strategic HR interventions in improving nurses' psychological well-being, job satisfaction, and organisational commitment. The findings may assist hospital administrators, HR managers, and healthcare policymakers in developing employee-centred workplace policies. The research also emphasises the importance of counselling, workload management, supportive leadership, and employee recognition in strengthening workforce retention. Furthermore, the study may contribute to academic literature by establishing the relationship between strategic human resource practices, compassion fatigue management, and long-term nursing workforce sustainability.

Limitations of the Study

The study may face certain limitations that could influence the interpretation and generalisation of its findings. The research may be restricted to a selected geographical area and a limited number of healthcare institutions, thereby affecting its broader applicability. The responses collected from nursing professionals may involve personal perceptions and subjective experiences, which could introduce response bias. Differences in hospital infrastructure, patient workload, management practices, and organisational culture may also influence the results. Additionally, compassion fatigue and employee retention are influenced by several personal and professional factors beyond HR interventions. Time and resource constraints may further restrict the scope of the investigation.

Research Gap

Existing research has extensively examined occupational stress, burnout, compassion fatigue, and employee turnover among healthcare professionals. However, comparatively limited attention has been given to evaluating the integrated impact of strategic HR interventions on both compassion fatigue and retention among frontline nursing staff. Many studies concentrate on individual psychological support mechanisms without sufficiently examining organisational HR strategies such as flexible scheduling, workload distribution, employee recognition, counselling, and supportive leadership. Furthermore, the relationship between institutional interventions, emotional resilience, organisational commitment, and nurses' intention to remain employed requires further investigation. This study seeks to address these gaps through an integrated assessment of strategic HR practices and nursing workforce sustainability.

Statement of Research Problem

Frontline nursing staff frequently experience emotional exhaustion, psychological distress, and compassion fatigue due to continuous exposure to patient suffering, demanding work schedules, staff shortages, and high-pressure healthcare environments. These challenges may reduce job satisfaction, weaken organisational commitment, and increase employee turnover intentions. Despite the availability of various employee welfare initiatives, many healthcare organisations lack systematic and strategically integrated HR interventions to address these concerns effectively. The central research problem is to determine whether strategic HR interventions can significantly reduce compassion

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429 Volume 6, Issue 5, November 2026 Impact Factor: 8.2

fatigue and improve retention among frontline nursing staff. The study also examines which HR practices contribute most effectively to employee well-being and workforce stability.

Research Objectives

1. To examine the impact of strategic Human Resource (HR) interventions on reducing compassion fatigue among frontline nursing staff.

2. To evaluate the effectiveness of strategic HR interventions in improving job satisfaction, organisational commitment, and employee retention among frontline nursing staff.

Research Hypotheses

1. H_0 (Null Hypothesis): There is no significant relationship between strategic HR interventions and the reduction of compassion fatigue among frontline nursing staff.

H_1 (Alternative Hypothesis): There is a significant relationship between strategic HR interventions and the reduction of compassion fatigue among frontline nursing staff.

2. H_0 (Null Hypothesis): Strategic HR interventions have no significant impact on job satisfaction, organisational commitment, and employee retention among frontline nursing staff.

H_1 (Alternative Hypothesis): Strategic HR interventions have a significant positive impact on job satisfaction, organisational commitment, and employee retention among frontline nursing staff.

II. REVIEW OF LITERATURE

1. AbuAlRub, R. F., & Alghamdi, M. G. (2012)

AbuAlRub and Alghamdi (2012) examined the influence of leadership styles on nurses' job satisfaction and intention to remain employed in healthcare organisations in Saudi Arabia. The study highlighted the importance of effective leadership in creating a supportive working environment and improving professional satisfaction among nursing staff. It indicated that leadership behaviour plays an important role in influencing employee motivation, organisational commitment, and retention decisions. The research suggests that supportive and participative leadership practices can strengthen nurses' willingness to continue working within healthcare institutions. These findings establish leadership development as an important strategic HR intervention for improving nursing workforce retention.

2. Aiken, L. H., Clarke, S. P., Sloane, D. M., Sochalski, J., & Silber, J. H. (2002)

Aiken et al. (2002) investigated the relationship between hospital nurse staffing levels, patient mortality, nurse burnout, and job dissatisfaction. The study highlighted that inadequate nurse staffing increases professional workload and creates stressful working conditions, adversely affecting nurses' psychological well-being and job satisfaction. It further established the importance of maintaining appropriate nurse-to-patient staffing levels to ensure quality healthcare delivery. The findings suggest that excessive workload and insufficient organisational support can contribute to burnout and employee dissatisfaction. The study provides a foundation for strategic HR interventions focusing on workforce planning, workload distribution, and adequate staffing to improve employee well-being and retention.

3. Arimon-Pagès, E., Torres-Puig-Gros, J., Fernández-Ortega, P., & Canela-Soler, J. (2019)

Arimon-Pagès et al. (2019) examined the emotional impact of professional responsibilities and compassion fatigue among oncology nurses through a multicentre study. The research highlighted the emotional challenges experienced by nurses who regularly interact with patients suffering from serious illnesses and their families. Continuous exposure to patient distress, suffering, and end-of-life situations can negatively affect nurses' emotional well-being and professional performance. The study emphasised the importance of recognising compassion fatigue as a significant occupational concern within healthcare institutions. Its findings support the implementation of psychological counselling, emotional support programmes, and stress management initiatives as strategic HR measures to strengthen nurses' emotional resilience.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429 Volume 6, Issue 5, November 2026 Impact Factor: 8.2

4. Brown, P., Fraser, K., Wong, C. A., Muise, M., & Cummings, G. (2013)

Brown et al. (2013) conducted a systematic review to identify factors influencing the intention to stay and retention of nurse managers. The study examined organisational and professional factors affecting nursing leadership retention, including workplace relationships, leadership support, job satisfaction, and organisational conditions. The researchers highlighted that supportive management practices and positive organisational environments contribute significantly to employee retention. The review also indicated that retention requires a comprehensive organisational approach rather than depending exclusively on financial incentives. The findings are relevant to the present study because they demonstrate the importance of strategic HR policies, professional development, supportive leadership, and employee engagement in maintaining workforce stability within healthcare institutions.

5. Cavanagh, S. J. (1990)

Cavanagh (1990) examined the factors influencing nursing staff turnover and identified important determinants associated with nurses' decisions to leave their employment. The research contributed to understanding the relationship between job-related factors, professional satisfaction, and employee turnover in nursing organisations. It highlighted the importance of recognising workplace conditions and individual employment experiences when developing effective retention strategies. The study provides an important foundation for understanding nursing workforce instability and the need for systematic employee retention practices. Its findings are relevant to the present research because strategic HR interventions addressing job satisfaction, working conditions, professional expectations, and organisational support may help reduce turnover intentions among frontline nursing staff.

6. Coetzee, S. K., & Klopper, H. C. (2010)

Coetzee and Klopper (2010) conducted a concept analysis of compassion fatigue within nursing practice to explain its meaning, characteristics, and implications for healthcare professionals. The researchers identified compassion fatigue as a condition associated with prolonged exposure to patients' suffering and the emotional demands of providing compassionate care. The study explained how continuous emotional involvement may affect nurses' psychological health, professional relationships, and quality of care. It also highlighted the importance of recognising compassion fatigue as a distinct occupational concern requiring appropriate organisational attention. The research provides a theoretical foundation for introducing employee assistance programmes, psychological counselling, and resilience-building initiatives within healthcare organisations.

7. Duarte, J., & Pinto-Gouveia, J. (2016)

Duarte and Pinto-Gouveia (2016) investigated the effectiveness of a mindfulness-based intervention in reducing burnout and compassion fatigue symptoms among oncology nurses. The study explored how mindfulness practices could help nursing professionals manage occupational stress, emotional exhaustion, and challenging patient-care experiences. The findings highlighted the potential of mindfulness-based programmes in improving emotional regulation and psychological well-being among healthcare employees. The research suggested that structured interventions supporting self-awareness and stress management can contribute to healthier professional functioning. Its findings are relevant to the present study because they demonstrate how healthcare organisations can incorporate psychological wellness programmes into strategic HR policies to reduce compassion fatigue and support nursing staff retention.

8. Figley, C. R. (Ed.). (1995)

Figley (1995), in the edited book Compassion Fatigue: Coping with Secondary Traumatic Stress Disorder in Those Who Treat the Traumatized, discussed the psychological consequences of prolonged exposure to individuals experiencing traumatic situations. The work established an important theoretical understanding of compassion fatigue and secondary traumatic stress among professionals providing emotional and psychological support. It explained how continuous exposure to others' suffering can result in emotional exhaustion, reduced empathy, and psychological distress. The publication emphasised the importance of coping mechanisms, emotional awareness, and supportive interventions. This theoretical contribution is particularly relevant to the present research because it establishes the conceptual basis for developing organisational strategies to protect frontline nursing staff.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429 Volume 6, Issue 5, November 2026 Impact Factor: 8.2

9. Halter, M., Boiko, O., Pelone, F., Beighton, C., Harris, R., Gale, J., Gourlay, S., & Drennan, V. (2017)

Halter et al. (2017) conducted a systematic review of systematic reviews to examine the determinants and consequences of adult nursing staff turnover. The research identified several factors associated with nurses' decisions to leave healthcare organisations, including job dissatisfaction, workplace conditions, professional stress, and organisational factors. It also highlighted the consequences of nursing turnover, such as increased recruitment expenses, workforce shortages, and additional pressure on existing employees. The study demonstrated that nursing retention is a complex organisational issue requiring comprehensive management strategies. Its findings support the present research by establishing the need for strategic HR interventions that improve workplace satisfaction, employee well-being, and organisational commitment.

10. Halter, M., Pelone, F., Boiko, O., Beighton, C., Harris, R., Gale, J., Gourlay, S., & Drennan, V. (2017)

Halter et al. (2017) examined interventions designed to reduce adult nursing staff turnover through a systematic review of systematic reviews. The study explored organisational approaches aimed at improving employee retention and reducing workforce instability in healthcare institutions. It highlighted the importance of workplace improvements, supportive management, professional development, and employee engagement in addressing nursing turnover. The research suggested that retention initiatives should be developed through an integrated organisational approach rather than relying on isolated interventions. The findings are directly relevant to the present study because they provide evidence supporting strategic HR planning, employee welfare programmes, and organisational support mechanisms to improve retention among frontline nursing professionals.

III. RESEARCH METHODOLOGY

Research Topic: Evaluating the Impact of Strategic HR Intervention on Compassion Fatigue and Retention among Frontline Nursing Staff

1. Research Design

The present study adopts a descriptive and analytical research design to evaluate the impact of strategic Human Resource (HR) interventions on compassion fatigue and employee retention among frontline nursing staff. The descriptive approach helps understand existing HR practices, working conditions, and levels of compassion fatigue among nursing professionals. The analytical approach examines the relationship between HR interventions, psychological well-being, job satisfaction, organisational commitment, and employee retention intentions. A quantitative research approach will be used to collect and analyse primary data through a structured questionnaire.

2. Nature of Data

The study will be based on both primary and secondary data.

Primary Data: Primary data will be collected directly from frontline nursing staff through structured questionnaires. The questionnaire will include questions relating to HR interventions, compassion fatigue, job satisfaction, organisational support, and employee retention.

Secondary Data: Secondary data will be collected from research journals, academic books, published research papers, healthcare reports, government publications, and relevant online resources.

3. Sampling Method

The study will adopt a convenience sampling technique to select respondents from frontline nursing staff working in hospitals and healthcare institutions. This method is selected because it facilitates the collection of responses from nursing professionals who are accessible and willing to participate in the research. Efforts will be made to include nurses from different departments and professional backgrounds to obtain varied perspectives.

4. Sample Size

The study will consider a total sample size of 100 respondents, comprising frontline nursing professionals employed in selected healthcare institutions. The respondents will include nurses working in different departments, such as emergency care, intensive care units (ICU), general wards, and other patient-care departments.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429 Volume 6, Issue 5, November 2026 Impact Factor: 8.2

5. Geographical Scope

The research will be conducted among frontline nursing staff working in selected hospitals and healthcare institutions in the chosen geographical area. The study will focus on their experiences concerning organisational HR practices, compassion fatigue, and employee retention.

6. Research Instrument

A structured questionnaire will be used as the primary research instrument. It will consist of multiple-choice questions and five-point Likert scale statements.

The questionnaire will cover the following dimensions:

Demographic and professional profile of respondents.

Awareness and availability of strategic HR interventions.

Level of compassion fatigue and emotional exhaustion.

Job satisfaction and organisational commitment.

Intention to remain employed within the healthcare organisation.

7. Tools for Data Analysis

The collected data will be classified, tabulated, and analysed using appropriate statistical techniques.

Percentage analysis will be used to understand the demographic profile and response distribution.

Chi-square tests will be applied to examine the association between selected HR interventions, compassion fatigue, and employee retention.

Tables and graphical representations will be used to present the research findings.

8. Hypothesis Testing

The study will test the following hypotheses:

H_{01}: There is no significant relationship between strategic HR interventions and the reduction of compassion fatigue among frontline nursing staff.

H_{11}: There is a significant relationship between strategic HR interventions and the reduction of compassion fatigue among frontline nursing staff.

H_{02}: Strategic HR interventions have no significant impact on job satisfaction, organisational commitment, and employee retention among frontline nursing staff.

H_{12}: Strategic HR interventions have a significant positive impact on job satisfaction, organisational commitment, and employee retention among frontline nursing staff.

9. Period of Study

The research will be conducted during the selected academic research period. The primary data collection, analysis, interpretation, and preparation of findings will be completed within the stipulated research schedule.

10. Ethical Considerations

Participation in the study will be voluntary. Respondents will be informed about the purpose of the research, and their identities and responses will be kept confidential. The collected information will be used exclusively for academic and research purposes. Appropriate care will be taken to avoid collecting personally identifiable information unnecessarily.

11. Sample Selection and Respondent Profile

A total of 100 frontline nursing staff will be selected for the study. The respondents will be drawn from different healthcare departments to ensure representation of nurses working under varied professional and emotional conditions.

The proposed respondent distribution is as follows:

Particulars	***Proposed respondents***
General Ward Nurses	25
ICU Nurses	20
Emergency Department Nurses	20

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 5, November 2026
Impact Factor: 8.2

Operation Theatre Nurses	15
Other Patient-Care Nurses	20
Total	100

Important methodological clarification: The above is a proposed distribution, not actual collected data. The final distribution should reflect the respondents who participate in the survey.

12. Inclusion and Exclusion Criteria

Inclusion criteria:

- Registered nursing professionals working in hospitals.
- Nurses directly involved in patient-care activities.
- Professionals having at least six months of work experience.
- Nurses willing to participate voluntarily.
- Exclusion criteria:
- Administrative and non-clinical hospital employees.
- Nursing students and interns.
- Nurses not directly involved in patient care.
- Individuals unwilling to participate in the study.

13. Measurement of Research Variables

The study will examine the following variables:

Independent variables	***Dependent variables***
Psychological counselling	Compassion fatigue
Flexible working schedules	Employee retention
Employee assistance programmes	Job satisfaction
Supportive leadership	Organisational commitment
Stress management programmes	Emotional well-being
Employee recognition	Retention intention

For measuring compassion fatigue, the researcher may consider the Professional Quality of Life Scale (ProQOL Version 5). A 2025 study has also developed and validated a Compassion Fatigue Assessment Scale specifically for Indian nurses.

IV. DATA ANALYSIS AND INTERPRETATION USING CHI-SQUARE METHOD

Research Topic: Evaluating the Impact of Strategic HR Intervention on Compassion Fatigue and Retention among Frontline Nursing Staff

Sample Size: 100 Respondents

Statistical Tool: Chi-Square Test

Level of Significance: 5% (0.05)

1. Introduction to Data Analysis

The collected data from 100 frontline nursing staff will be analysed using the Chi-Square statistical method to examine the relationship between strategic Human Resource (HR) interventions, compassion fatigue, job satisfaction, organisational commitment, and employee retention. The analysis will focus on testing the two research objectives and their corresponding hypotheses. For the purpose of this illustrative analysis, assumed responses have been used to

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429 Volume 6, Issue 5, November 2026 Impact Factor: 8.2

demonstrate the application of the Chi-Square test. The observed frequencies will be compared with expected frequencies to determine whether statistically significant associations exist between the selected variables.

2. Objective 1

To examine the impact of strategic Human Resource (HR) interventions on reducing compassion fatigue among frontline nursing staff.

Hypothesis 1

H_{01} (Null Hypothesis): There is no significant relationship between strategic HR interventions and the reduction of compassion fatigue among frontline nursing staff.

H_{11} (Alternative Hypothesis): There is a significant relationship between strategic HR interventions and the reduction of compassion fatigue among frontline nursing staff.

Table 1: Relationship between HR Interventions and Compassion Fatigue

The following assumed data represent respondents' perceptions of the availability of HR interventions and their experience of compassion fatigue.

HR Intervention Availability	*High Compassion Fatigue*	*Low Compassion Fatigue*	*Total*
Adequate HR Support	15	35	50
Inadequate HR Support	35	15	50
Total	50	50	100

Calculation of Expected Frequency

Formula:

$$E = \frac{RT \times CT}{GT}$$

Where:

E = Expected Frequency

RT = Row Total

CT = Column Total

GT = Grand Total

Expected frequencies:

HR Intervention Availability	*High Compassion Fatigue*	*Low Compassion Fatigue*
Adequate HR Support	25	25
Inadequate HR Support	25	25

Chi-Square Calculation

$$\chi^2 = \sum \frac{(O - E)^2}{E}$$

Observed (O)	*Expected (E)*	$O - E$	$(O - E)^2/E$
15	25	-10	4
35	25	10	4
35	25	10	4
15	25	-10	4

Copyright to IJARSCT
www.ijarsct.co.in
DOI: 10.48175/IJARSCT-41033

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 5, November 2026
Impact Factor: 8.2

Total			16.00

O = Observed Frequency; E = Expected Frequency.

Degrees of Freedom

$$df = (r-1)(c-1)$$

$$df = (2-1)(2-1) = 1$$

Calculated Chi-Square Value = 16.00

Table Value at 5% significance level and 1 degree of freedom = 3.841

Significance Level = 0.05

Interpretation

The calculated Chi-Square value of 16.00 is greater than the critical table value of 3.841. Therefore, the null hypothesis (H_{01}) is rejected and the alternative hypothesis (H_{11}) is accepted. The findings indicate a statistically significant association between the availability of strategic HR interventions and compassion fatigue among frontline nursing staff. Respondents reporting adequate HR support show a lower proportion of high compassion fatigue (30%) compared with those reporting inadequate HR support (70%). This suggests that organisational support, counselling facilities, stress management programmes, and supportive working conditions may contribute to reducing compassion fatigue.

H_{01} Rejected

3. Objective 2

To evaluate the effectiveness of strategic HR interventions in improving job satisfaction, organisational commitment, and employee retention among frontline nursing staff.

Hypothesis 2

H_{02} (Null Hypothesis): Strategic HR interventions have no significant impact on job satisfaction, organisational commitment, and employee retention among frontline nursing staff.

H_{12} (Alternative Hypothesis): Strategic HR interventions have a significant positive impact on job satisfaction, organisational commitment, and employee retention among frontline nursing staff.

Table 2: Relationship between HR Interventions and Employee Retention

The following assumed data examine whether the availability of adequate HR interventions is associated with nurses' intention to remain employed in their present healthcare organisation.

HR Intervention Availability	*Intend to Stay*	*Do Not Intend to Stay*	*Total*
Adequate HR Support	40	10	50
Inadequate HR Support	20	30	50
Total	60	40	100

Calculation of Expected Frequency

Formula:

$$E = \frac{RT \times CT}{GT}$$

Expected frequencies:

HR Intervention Availability	*Intend to Stay*	*Do Not Intend to Stay*
Adequate HR Support	30	20
Inadequate HR Support	30	20

Chi-Square Calculation

Copyright to IJARSCT
www.ijarsct.co.in



DOI: 10.48175/IJARSCT-41033



IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 5, November 2026
Impact Factor: 8.2

$$\chi^2 = \sum \frac{(O-E)^2}{E}$$

Observed (O)	*Expected (E)*	*O − E*	*(O − E)²/E*
40	30	10	3.33
10	20	-10	5.00
20	30	-10	3.33
30	20	10	5.00
Total			16.66

Degrees of Freedom

$$df = (r-1)(c-1)$$

$$df = (2-1)(2-1) = 1$$

Statistical Results:

Calculated Chi-Square Value = 16.66

Table Value at 5% significance level and 1 degree of freedom = 3.841

Significance Level = 0.05

Interpretation

The calculated Chi-Square value of 16.66 is greater than the critical table value of 3.841. Therefore, the null hypothesis (H_{02}) is rejected and the alternative hypothesis (H_{12}) is accepted. The findings indicate a statistically significant association between strategic HR interventions and employee retention intentions among frontline nursing staff. The analysis reveals that 80% of respondents receiving adequate HR support intend to remain employed, compared with only 40% of respondents reporting inadequate HR support. This indicates that supportive leadership, employee recognition, flexible scheduling, professional development, and psychological assistance may contribute to improving nurses' organisational commitment and retention intentions.

H_{02} Rejected

4. Summary of Hypothesis Testing

The results of both Chi-Square tests are summarised below:

Particulars	*Objective 1*	*Objective 2*
Sample Size	100	100
Calculated Chi-Square	16.00	16.66
Table Value	3.841	3.841
Degrees of Freedom	1	1
Significance Level	5%	5%
Null Hypothesis	Rejected	Rejected
Alternative Hypothesis	Accepted	Accepted

5. Overall Findings of Statistical Analysis

Based on the above Chi-Square analysis, the following findings emerge:

1. There is a statistically significant association between adequate HR interventions and lower levels of compassion fatigue among frontline nursing staff.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429 Volume 6, Issue 5, November 2026 Impact Factor: 8.2

2. Respondents receiving adequate HR support report comparatively lower compassion fatigue than those experiencing inadequate organisational support.
3. There is a statistically significant association between strategic HR interventions and nurses' intention to remain employed in their healthcare organisations.
4. Employee assistance programmes, psychological counselling, supportive leadership, and flexible working arrangements may contribute to improved employee well-being.
5. Adequate HR support is associated with stronger retention intentions among frontline nursing professionals, highlighting the importance of employee-centred HR policies.

6. Conclusion of Hypothesis Testing

The statistical analysis demonstrates that both null hypotheses are rejected at the 5% level of significance. The findings suggest that strategic HR interventions are significantly associated with compassion fatigue and employee retention intentions among frontline nursing staff. Healthcare organisations should therefore consider strengthening employee welfare programmes, psychological support systems, workload management, leadership practices, and professional development opportunities. Such initiatives may help create a supportive working environment and improve nursing workforce stability.

Important research note: All frequencies above are assumed figures prepared for illustrative academic analysis, not actual survey findings. They must be replaced with the responses collected from the 100 respondents before submission. Also, Objective 2 includes job satisfaction and organisational commitment, which have not been separately tested in Table 2. Separate cross-tabulations and Chi-Square tests should be conducted for these variables if all three dimensions are to be statistically evaluated. The analysis establishes association rather than causal impact.

Challenges in Managing Compassion Fatigue and Employee Retention among Frontline Nursing Staff

1. Emotional Exhaustion and Psychological Stress

Frontline nursing professionals regularly encounter patients suffering from critical illnesses, traumatic injuries, and life-threatening medical conditions. Continuous exposure to pain, death, and emotional distress can result in compassion fatigue and psychological exhaustion. Nurses may experience reduced empathy, emotional detachment, anxiety, and decreased motivation towards their professional responsibilities. These conditions can adversely affect their personal well-being and quality of patient care. Managing emotional exhaustion becomes challenging when healthcare institutions lack adequate psychological counselling, employee assistance programmes, and stress management initiatives. Therefore, organisations must develop structured emotional support systems to strengthen nurses' psychological resilience and professional effectiveness.

2. Inadequate Staffing and Excessive Workload

Insufficient nursing staff and increasing patient numbers create significant workload-related challenges within healthcare institutions. Nurses are frequently required to work extended shifts, manage multiple patients, and perform additional administrative responsibilities. Such demanding conditions reduce opportunities for adequate rest and recovery, increasing physical and emotional exhaustion. Excessive workload may also contribute to workplace dissatisfaction, reduced efficiency, and compassion fatigue. Healthcare organisations often face difficulties in maintaining an appropriate nurse-to-patient ratio due to financial limitations and recruitment challenges. Strategic workforce planning, systematic duty allocation, adequate recruitment, and flexible scheduling are essential to reduce workload pressure and improve employee satisfaction.

3. Lack of Effective HR Support and Organisational Recognition

Many healthcare organisations face challenges in implementing effective HR interventions specifically designed to address the psychological and professional needs of nursing staff. Limited access to counselling, employee assistance programmes, professional development opportunities, and recognition initiatives may create feelings of organisational neglect. When nurses believe that their contributions are not adequately acknowledged, their motivation and organisational commitment may decline. Additionally, inadequate communication between management and nursing

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429 Volume 6, Issue 5, November 2026 Impact Factor: 8.2

professionals can create dissatisfaction and weaken workplace relationships. Healthcare institutions must establish employee-centred HR policies, transparent grievance mechanisms, recognition programmes, and supportive leadership practices to improve workplace experiences and strengthen professional commitment.

4. High Employee Turnover and Retention Difficulties

Employee turnover is a major challenge for healthcare organisations because experienced nursing professionals may leave their positions due to excessive workload, emotional exhaustion, limited career advancement, inadequate compensation, and unsatisfactory working conditions. Frequent resignations increase recruitment and training expenses while placing additional responsibilities on existing nursing staff. This may create a continuous cycle of staff shortages, increased workload, and declining employee satisfaction. Retaining qualified nursing professionals therefore requires more than financial incentives. Organisations must introduce effective retention strategies, career development opportunities, flexible working arrangements, competitive compensation, and supportive management practices to encourage nurses to continue their professional association with healthcare institutions.

5. Difficulty in Measuring and Monitoring Compassion Fatigue

Measuring compassion fatigue among nursing professionals presents another significant organisational challenge because emotional exhaustion and psychological distress are subjective experiences. Individual nurses may respond differently to similar workplace situations depending on their professional experience, coping mechanisms, personal circumstances, and emotional resilience. Some employees may hesitate to disclose psychological difficulties due to fear of professional judgement or organisational consequences. Furthermore, healthcare institutions may lack standardised assessment procedures and regular monitoring systems. The absence of reliable measurement mechanisms makes it difficult for HR departments to identify affected employees and evaluate intervention effectiveness. Regular psychological assessments, confidential feedback systems, and validated measurement instruments can help address this challenge.

V. REMEDIES AND SUGGESTIONS

1. Implementation of Psychological Counselling and Employee Wellness Programmes

Healthcare organisations should establish structured psychological counselling and employee wellness programmes to address compassion fatigue among frontline nursing staff. Regular counselling sessions, stress management workshops, mindfulness activities, and emotional resilience training can help nurses manage occupational stress and emotional exhaustion. Employee Assistance Programmes (EAPs) should provide confidential psychological support to employees experiencing workplace-related difficulties. Organisations should also encourage open communication about mental well-being without creating fear of discrimination or professional judgement. These initiatives can improve emotional stability, strengthen coping mechanisms, and promote a healthier working environment.

2. Effective Workforce Planning and Workload Management

Healthcare institutions should adopt systematic workforce planning to maintain adequate nurse-to-patient ratios and ensure equitable distribution of responsibilities. Excessive working hours, continuous night shifts, and insufficient rest periods should be addressed through effective duty scheduling and shift rotation. Flexible working arrangements and planned leave policies can help nursing professionals maintain a healthy work-life balance. Hospital management should regularly evaluate staffing requirements according to patient volume and departmental workload. Adequate recruitment and deployment of nursing professionals can reduce excessive work pressure, minimise physical and emotional exhaustion, and improve the quality of patient care.

3. Strengthening Strategic HR Policies and Employee Recognition

Human Resource departments should introduce employee-centred policies that recognise the professional contribution and emotional responsibilities of nursing staff. Regular performance appreciation, financial incentives, transparent promotion policies, grievance redressal mechanisms, and professional development opportunities can improve employee motivation. Healthcare organisations should also encourage participative decision-making and maintain effective communication between nursing staff and management. Supportive leadership practices can create a positive

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429 Volume 6, Issue 5, November 2026 Impact Factor: 8.2

organisational culture where employees feel valued and respected. These measures can strengthen organisational commitment, improve job satisfaction, and encourage nursing professionals to maintain long-term employment relationships with healthcare institutions.

4. Developing Comprehensive Employee Retention Strategies

Healthcare organisations should implement comprehensive retention strategies to reduce employee turnover and retain experienced nursing professionals. Competitive compensation, career advancement opportunities, specialised training, mentorship programmes, and supportive working conditions should form important components of HR policies. Management should conduct periodic employee satisfaction surveys and exit interviews to identify the reasons behind nursing staff turnover. Special attention should be given to employees working in high-pressure departments such as intensive care units and emergency services. A structured employee retention framework can improve professional satisfaction, reduce recruitment costs, maintain institutional knowledge, and ensure continuity in healthcare delivery.

5. Regular Assessment and Monitoring of Compassion Fatigue

Healthcare institutions should introduce systematic mechanisms to identify and monitor compassion fatigue among nursing professionals. Periodic psychological assessments using validated instruments, such as the Professional Quality of Life Scale (ProQOL), can help identify compassion fatigue, burnout, and compassion satisfaction. Confidential employee feedback systems should be established to encourage nurses to communicate their concerns without hesitation. HR departments should evaluate the effectiveness of wellness initiatives through regular assessments and employee satisfaction surveys. Data-driven monitoring can help management identify vulnerable employee groups, introduce corrective interventions, and assess improvements in psychological well-being and retention intentions.

VI. CONCLUSION

The present study, titled “Evaluating the Impact of Strategic HR Intervention on Compassion Fatigue and Retention among Frontline Nursing Staff,” highlights the importance of strategic Human Resource Management in addressing the emotional, psychological, and professional challenges experienced by nursing professionals. Frontline nurses play an essential role in healthcare delivery; however, continuous exposure to patient suffering, demanding working conditions, excessive workloads, and emotional responsibilities can contribute to compassion fatigue, reduced job satisfaction, and increased employee turnover intentions.

The study emphasises that traditional HR practices focusing primarily on recruitment, compensation, and performance evaluation are insufficient to address the complex requirements of nursing professionals. Healthcare organisations must adopt comprehensive HR interventions, including psychological counselling, employee assistance programmes, flexible working schedules, stress management training, supportive leadership, employee recognition, and professional development opportunities. Such initiatives can help create a positive organisational environment and strengthen employees' emotional resilience.

The illustrative Chi-Square analysis, based on assumed data from 100 respondents, indicates a statistically significant association between adequate HR support and lower compassion fatigue. Similarly, the analysis identifies a significant association between strategic HR interventions and nurses' intention to remain employed within their healthcare organisations. Both null hypotheses were rejected at the 5% level of significance. These results demonstrate the potential importance of employee-centred HR strategies in improving nursing workforce stability.

Furthermore, the study identifies major challenges, including emotional exhaustion, inadequate staffing, excessive workload, insufficient organisational recognition, employee turnover, and difficulties in measuring compassion fatigue. Addressing these challenges requires systematic workforce planning, regular psychological assessments, effective communication, and continuous evaluation of HR policies.

In conclusion, strategic HR interventions should be considered an essential component of sustainable healthcare workforce management. Healthcare institutions must recognise nursing professionals not merely as service providers but as valuable human resources whose psychological well-being directly influences organisational effectiveness and patient care quality. A supportive, inclusive, and employee-oriented HR framework can contribute to reducing

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 5, November 2026
Impact Factor: 8.2

compassion fatigue, strengthening organisational commitment, and improving long-term retention among frontline nursing staff.

REFERENCES

1. AbuAlRub, R. F., & Alghamdi, M. G. (2012). The impact of leadership styles on nurses' satisfaction and intention to stay among Saudi nurses. Journal of Nursing Management, 20(5), 668–678. https://doi.org/10.1111/j.1365-2834.2011.01320.x

2. Aiken, L. H., Clarke, S. P., Sloane, D. M., Sochalski, J., & Silber, J. H. (2002). Hospital nurse staffing and patient mortality, nurse burnout, and job dissatisfaction. JAMA, 288(16), 1987–1993. https://doi.org/10.1001/jama.288.16.1987

3. Arimon-Pagès, E., Torres-Puig-Gros, J., Fernández-Ortega, P., & Canela-Soler, J. (2019). Emotional impact and compassion fatigue in oncology nurses: Results of a multicentre study. European Journal of Oncology Nursing, 43, Article 101666.

4. Brown, P., Fraser, K., Wong, C. A., Muise, M., & Cummings, G. (2013). Factors influencing intentions to stay and retention of nurse managers: A systematic review. Journal of Nursing Management, 21, 459–472. https://doi.org/10.1111/j.1365-2834.2012.01352.x

5. Cavanagh, S. J. (1990). Predictors of nursing staff turnover. Journal of Advanced Nursing, 15(3), 373–380.

6. Coetzee, S. K., & Klopper, H. C. (2010). Compassion fatigue within nursing practice: A concept analysis. Nursing & Health Sciences, 12(2), 235–243. https://doi.org/10.1111/j.1442-2018.2010.00526.x

7. Duarte, J., & Pinto-Gouveia, J. (2016). Effectiveness of a mindfulness-based intervention on oncology nurses' burnout and compassion fatigue symptoms: A non-randomized study. International Journal of Nursing Studies, 64, 98–107.

8. Figley, C. R. (Ed.). (1995). Compassion fatigue: Coping with secondary traumatic stress disorder in those who treat the traumatized. Brunner/Mazel.

9. Halter, M., Boiko, O., Pelone, F., Beighton, C., Harris, R., Gale, J., Gourlay, S., & Drennan, V. (2017). The determinants and consequences of adult nursing staff turnover: A systematic review of systematic reviews. BMC Health Services Research, 17, Article 824. https://doi.org/10.1186/s12913-017-2707-0

10. Halter, M., Pelone, F., Boiko, O., Beighton, C., Harris, R., Gale, J., Gourlay, S., & Drennan, V. (2017). Interventions to reduce adult nursing turnover: A systematic review of systematic reviews. The Open Nursing Journal, 11, 108–123. https://doi.org/10.2174/1874434601711010108

11. Hayes, L. J., O'Brien-Pallas, L., Duffield, C., Shamian, J., Buchan, J., Hughes, F., Laschinger, H. K. S., & North, N. (2006). Nurse turnover: A literature review. International Journal of Nursing Studies, 43(2), 237–263.

12. Hunsaker, S., Chen, H. C., Maughan, D., & Heaston, S. (2015). Factors that influence the development of compassion fatigue, burnout, and compassion satisfaction in emergency department nurses. Journal of Nursing Scholarship, 47(2), 186–194. https://doi.org/10.1111/jnu.12122

13. Joinson, C. (1992). Coping with compassion fatigue. Nursing, 22(4), 116–121.

14. Lake, E. T. (2002). Development of the practice environment scale of the Nursing Work Index. Research in Nursing & Health, 25(3), 176–188.

15. Manzanedo-Moreno, M., Salcedo-Sampedro, C., Abajas-Bustillo, R., Martin-Melón, R., Fernández Peña, R., & Ortego-Maté, C. (2025). Interventions to reduce compassion fatigue in nursing: A systematic review and meta-analysis of controlled studies. International Nursing Review, 72(2), Article e70023. https://doi.org/10.1111/inr.70023

16. Maslach, C., & Leiter, M. P. (2016). Understanding the burnout experience: Recent research and its implications for psychiatry. World Psychiatry, 15(2), 103–111. https://doi.org/10.1002/wps.20311

17. Pressley, C., & Garside, J. (2023). Safeguarding the retention of nurses: A systematic review on determinants of nurses' intentions to stay. Nursing Open, 10, 2842–2858. https://doi.org/10.1002/nop2.1588

18. Stamm, B. H. (2010). The concise ProQOL manual (2nd ed.). ProQOL.org.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 5, November 2026
Impact Factor: 8.2

19. Zhang, H., Xia, Z., Yu, S., Shi, H., Meng, Y., & Dator, W. L. (2025). Interventions for compassion fatigue, burnout, and secondary traumatic stress in nurses: A systematic review and network meta-analysis. Nursing & Health Sciences, 27(1), Article e70042. https://doi.org/10.1111/nhs.70042

20. Pressley, C., & Garside, J. (2023). Safeguarding the retention of nurses: A systematic review on determinants of nurses' intentions to stay. Nursing Open, 10, 2842–2858. https://doi.org/10.1002/nop2.1588