IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 6, November 2026
Impact Factor: 8.2

Dietary Fat Quality and Human Health: Evidence-Based Study

Mr. Rehan Shaikh[1], Dr. Bhagwan M. Kale[2], Pathan Taufik Khan Rafik Khan[3], Pathan Sami Khan Yousuf Khan[4], Shaikh Faizan Rafique[5]

[1] Assistant Professor, [2] Principal, [3],[4],[5] Students

Shri Goraksha College of Pharmacy and Research Centre, Khamgaon, Maharashtra, India

Abstract : *Dietary fats are essential components of human nutrition, providing energy and supporting cell membrane structure, hormone production, and the absorption of fat-soluble vitamins. However, the type and quality of fat consumed may influence cardiovascular and metabolic health. This review article examines the health effects of saturated, unsaturated, and trans fats, with attention to dietary cholesterol and evidence discussed in research and dietary guidelines from 2015 to 2026. Its objectives are to compare the effects of different fats on blood lipid profiles, inflammation, metabolic health, body weight, and cardiovascular risk, and to summarize evidence-based recommendations for healthier fat consumption. The article reviews published studies, guideline summaries, and related evidence concerning monounsaturated fatty acids (MUFA), polyunsaturated fatty acids (PUFA), omega-3 and omega-6 fatty acids, saturated fats, trans fats, and dietary cholesterol. The evidence summarized indicates that diets containing unsaturated fats, particularly MUFA and PUFA, are associated with improved LDL and HDL cholesterol profiles and other beneficial cardiovascular and metabolic effects. Industrial trans fats are are associated with adverse lipid changes, inflammation, inf and increased cardiovascular risk, while the effects of saturated fats depend partly on the food source and the nutrient replacing them. Dietary cholesterol alone is described as having a limited effect on blood cholesterol for most people compared with overall dietary fat quality. The article concludes that replacing unhealthy fats with healthier unsaturated-fat sources, limiting saturated fat, and avoiding trans fats are important elements of a balanced dietary pattern for supporting long-term health and reducing chronic disease risk.*

Keywords: Dietary fats; unsaturated fats; saturated fats; trans fats; dietary cholesterol; monounsa (MUFA); polyunsaturated fatty acids (PUFA); LDL cholesterol; HDL cholesterol; ca metabolic health; inflammation; nutrition guidelines.

Copyright to IJARSCT
www.ijarsct.co.in

DOI: 10.48175/568

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 6, November 2026
Impact Factor: 8.2

INTRODUCTION:

Dietary fats agential part of human nutrition and play a major role in energy production, cell structure, hormone synthesis, and the absorption of fat-soluble vitamins (A, D, E, and K). However, not all fats have the same effect on health.The quality of dietary fat consumed is more important than the total quantity. Scientific evidence shows that replacing unhealthy fats with healthier fats can reduce the risk of cardiovascular disease,obesity,diabetes,andinflammation.This study examines different types of dietary fats, their effects on human health, and evidence-based recommendations for healthy fat consumption.The way consumers obtain nutrition information has changed substantially in the past two decades. Use of the internet and social media has grown rapidly, and these are now among the leading sources of information for health and wellness. Perhaps due to access to more information than ever, including conflicting information of uncertain and variable quality, many consumers are more confused than ever.Nutrition scientists are trusted content experts . Consequently, it is essential that they effectively communicate research findings to policy makers, authoritative bodies and the general public in order for consumers to make sound, evidence-based dietary decisions. In addition, communicating scientific findings can be viewed as a civic duty and has been suggested to be included in formal academic training . Yet, scientists' ability to communicate is less than other professionalsWithin the nutrition community, one example of suboptimal communication between scientists and the public is the continued demonization and general avoidance of dietary fat . For years, an emphasis of nutrition communication was to balance calorie intake and energy expenditure and decrease dietary fat.




Fig.1 Dietary Guideline

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 6, November 2026
Impact Factor: 8.2

Reductions in total dietary fat were recommended to reduce saturated fat as well as due to the energy density of lipids and the overall goal to reduce caloric intake . Partly as a result, low-fat, high-carbohydrate diets were recommended in 1980 and thereafter for weight loss and reducing cardiovascular disease (CVD) risk. However this led to unintended consequences. The focus on reducing total fat resulted in increased consumption of refined carbohydrates and added sugars, and avoidance of nutrient-dense foods rich in healthy unsaturated fats such as nuts, seeds, avocados and vegetable oils. Subsequently, fat consumption has decreased while carbohydrate intake has increased as percentage of calories, which has been accompanied by significant increases in total energy intake and obesity rates in the United States.

While single nutrient targets have worked well for treating diseases of deficiency, this has been problematic for addressing chronic diseases . In contrast, specific foods and overall dietary patterns can substantially affect chronic disease

Risk. However, by attributing their effects to single nutrients, foods with very different physiological effects can become conflated and contribute to consumer confusion. A global survey found that 95% of respondents knew that vitamins were needed for a healthy diet, but only 41% knew certain fats were essential nutrients .

In recent decades, we have gained substantial knowledge regarding the role of broad classes of nutrients and foods in majorchronic diseases.

In addition, challenges in communicating science clearly to the public contributes to consumer confusion and, as a result, may have public health consequences. This paper summarizes an Experimental Biology Scientific Session for which the goals were to discuss the history of recommendations for dietary fat and evidence showing health benefits of unsaturated fats from plant sources in order to provide context for the strategies described to improve nutrition science communications to the public.

OBJECTIVE:

- To evaluate the role of different types of dietary fats (saturated, unsaturated, and trans fats) on overall human health based on recent scientific evidence (2015–2026).
- To analyze the impact of dietary fat quality on serum lipid profile, including LDL, HDL, triglycerides, and total cholesterol levels.
- To determine whether dietary cholesterol directly contributes to heart disease, using updated guidelines and research findings.
- To compare the health effects of saturated fats vs. unsaturated fats, especially their influence on cardiovascular risk.
- To assess scientific evidence regarding trans fats and their association with inflammation, metabolic diseases, and cardiac risk.
- To study the recommended daily intake of healthy fats according to global guidelines (WHO, AHA, etc.) and understand their importance in diet.
- To identify the beneficial roles of mono-unsaturated and poly-unsaturated fats in reducing chronic disease risk.

Copyright to IJARSCT
www.ijarsct.co.in
DOI: 10.48175/568

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 6, November 2026
Impact Factor: 8.2

- To review clinical trials and epidemiological studies published between 2015 and 2026 that examine the link between fat quality and health outcomes.
- To explore how fat quality affects metabolic processes, including insulin sensitivity, inflammation markers, and body weight.
- To provide evidence-based recommendations on optimal fat consumption for improving human health and preventing diseases.

LITERATURE REVIEW:

1. Shift in Understanding of Dietary Cholesterol (2015–2026)

Earlier guidelines linked dietary cholesterol to heart disease. However, the 2015–2020 Dietary Guidelines for Americans removed cholesterol as a "nutrient of concern," stating that food cholesterol has minimal impact on blood cholesterol in most individuals.

Studies (Berger et al., 2018; Carson et al., 2016) found no significant association between dietary cholesterol intake and cardiovascular disease (CVD) risk.

Conclusion: Total cholesterol from food is not a major driver of heart disease; fat quality matters more.

2. Saturated Fat and Heart Disease: A Controversial Area

Research post-2015 challenged the old belief that saturated fat directly causes heart disease.

A meta-analysis by de Souza et al. (2015) found no strong evidence linking saturated fat with CVD mortality.

However, replacing saturated fat with refined carbohydrates showed no benefit, while replacing it with unsaturated fats reduced CVD risk (Li et al., 2015; Mozaffarian & Ludwig, 2016).

WHO and AHA still recommend limiting saturated fat to <10% of total calories.

Conclusion: Saturated fat is not equally harmful across foods; context and replacement nutrient matter.

3. Health Benefits of Unsaturated Fats

Evidence strongly supports the protective effects of mono-unsaturated (MUFA) and poly-unsaturated fats (PUFA):

3.1 MUFA – Olive Oil

- Studies including the PREDIMED Trial (2018) showed that diets rich in olive oil decrease:
- LDL oxidation
- Inflammatory markers
- CVD incidence

3.2 Omega-3 PUFA – Fish Oils

- EPA and DHA have been linked to:
- Reduced triglycerides
- Lower arrhythmia risk
- Anti-inflammatory effects
- Improved endothelial function (Nicholls et al., 2019)

3.3 Omega-6 PUFA

Despite controversy, research confirms omega-6 fats lower LDL and reduce heart-disease risk when replacing saturated fats.

Conclusion: Unsaturated fats are consistently associated with improved cardiovascular and metabolic health.

4. Trans Fats and Their Strongly Established Harm

Industrial trans fats are universally recognized as harmful.

They increase LDL, decrease HDL, and cause systemic inflammation.

WHO eliminated industrial trans fats globally under the REPLACE Initiative (2018–2023).

Multiple studies show trans fats significantly increase CVD and all-cause mortality.

Conclusion: Trans fats are unequivocally harmful and should be completely eliminated.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 6, November 2026
Impact Factor: 8.2

5. Fat Quality and Metabolic Health
Recent studies explore the impact of fat type on metabolic markers:
High unsaturated fat diets improve insulin sensitivity (Rosqvist et al., 2019).
Saturated fat increases liver fat accumulation, increasing NAFLD risk.
MUFA-rich diets improve gut microbiota, lowering inflammatory pathways.
Conclusion: Fat quality affects more than heart health—it also influences insulin signaling, fatty liver, gut health, and inflammation.
6. Dietary Fat Quality, Obesity & Weight Regulation
Evidence suggests fat type affects body weight:
Diets higher in MUFA and PUFA improve satiety and metabolic rate.
Trans fats and saturated fats (from processed foods) are linked to visceral fat accumulation.
Mediterranean diet trials show weight loss despite high fat intake.
Conclusion: Fat quality influences weight and body composition more than total fat quantity.

Total Fat:

The percentage of energy consumed as fat can vary widely, and the diet can still meet energy and nutrient needs. Current recommendations from various organizations regarding fat intake in adults are summarized in Table .

Dietary guidelines from the World Health Organization and the Dietary Reference Intakes recommend a total fat intake between 20 and 35% of total calories The minimum of 20% is to ensure adequate consumption of total energy, essential fatty acids, and fat-soluble vitamins and prevent atherogenic dyslipidemia (low high-density lipoprotein cholesterol (HDL-C), high triglyceride-rich lipoproteins) which occurs with low-fat, high carbohydrate diets and increases risk of coronary heart disease .

The maximum of 35% was based on limiting saturated fat and also the observation that individuals on higher fat diets consume more calories, resulting in weight gain . No Tolerable Upper Intake Level was set for total fat because there is no intake level for which there is an adverse event Of note, the 2015 Dietary Guidelines Advisory Committee placed emphasis on the types and quality of foods consumed and did not set an upper limit for total fat based on the lack of supporting evidence .

This was reflected in the Dietary Guidelines for Americans 2015–2020, which emphasizes types of fat within the context of a healthy dietary pattern .

Total fat is an essential macronutrient required for normal growth, development, and maintenance of human health. It provides a concentrated source of energy, supplying approximately 9 kcal per gram, which is more than twice the energy provided by carbohydrates and proteins.

Dietary fats play a vital role in the absorption of fat-soluble vitamins such as vitamins A, D, E, and K, and are important for hormone production, cell membrane structure, brain function, and insulation of body organs.

Total fat in the diet includes different types of fatty acids, mainly saturated fats, unsaturated fats (monounsaturated and polyunsaturated fats), and trans fats. The quality and quantity of fat consumed greatly influence overall health.

Healthy fats obtained from foods such as nuts, seeds, fish, olive oil, and avocados are associated with beneficial effects on heart health and inflammation, whereas excessive intake of saturated and trans fats may increase the risk of obesity, cardiovascular diseases, diabetes, and other metabolic disorders.

Copyright to IJARSCT
www.ijarsct.co.in
DOI: 10.48175/568

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 6, November 2026
Impact Factor: 8.2

According to nutritional guidelines, fats should contribute a balanced proportion of daily caloric intake. Emphasis is placed not only on reducing excessive fat consumption but also on improving dietary fat quality by replacing harmful fats with healthier unsaturated fats. Therefore, understanding total fat and its impact on human health is important in developing evidence-based dietary recommendations and promoting long-term well-being.

Human Health:

Human health is one of the most fundamental aspects of life and is considered the greatest wealth a person can possess. The World Health Organization defines health as "a state of complete physical, mental and social well-being and not merely the absence of disease or infirmity.

This definition highlights that health is a holistic concept that goes beyond treating illness. It involves maintaining a balanced lifestyle, preventing disease, and ensuring mental and emotional stability. The health of an individual is influenced by a wide range of factors.

These include genetics, nutrition, personal hygiene, physical activity, environmental conditions, socioeconomic status, and access to quality healthcare services.

In today's fast-paced world, lifestyle-related disorders such as diabetes, hypertension, obesity, and cardiovascular diseases are increasing rapidly. At the same time, infectious diseases, antimicrobial resistance, and emerging viral

outbreaks continue to challenge global health systems.

The field of pharmacy plays a crucial role in safeguarding human health.

Pharmacists are not only responsible for dispensing medicines but also for ensuring their safe and effective use.

They contribute to drug discovery, formulation development, clinical research, and patient counseling.

Through pharmaceutical care, pharmacists help reduce medication errors, improve patient compliance, and promote rational use of drugs.

The development of vaccines, antibiotics, and advanced drug delivery systems has significantly increased human life expectancy and quality of life.

In recent years, there has been growing interest in preventive healthcare and traditional systems of medicine. Herbal and natural products are being explored for their therapeutic benefits, fewer side effects, and cost-effectiveness.

Public health awareness, sanitation, vaccination drives, and health education programs also play a major role in disease prevention and health promotion.

Maintaining human health is not the responsibility of healthcare professionals alone. It requires active participation from individuals, communities, governments, and researchers. Advances in pharmaceutical sciences, biotechnology, and medical research continue to provide new solutions for existing and emerging health problems.

The ultimate aim of all healthcare-related fields is to ensure that every individual can live a long, healthy, and productive life.

In conclusion, human health is a dynamic and multifaceted concept that forms the foundation of personal and national development.

Protecting and improving health through scientific research, responsible healthcare practices, and public awareness is essential for building a healthier society.?

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 6, November 2026
Impact Factor: 8.2

Alcohol:

Alcohol consumption impacts nearly every organ system, causing short-term impairment and long-term chronic diseases. Immediate effects include lowered inhibition, slurred speech, and impaired coordination, while chronic misuse increases risks of liver disease, cancer, heart disease, brain damage, and mental health issues like anxiety and depression. No level of consumption is considered entirely safe, with risks proportional to amount.

Short-Term Side Effects (Acute Consumption):

Nervous System: Relaxation, euphoria, slurred speech, **Central** slowed reflexes, and impaired judgment.

Digestive System: Nausea, vomiting, diarrhea, and irritation of the stomach lining.

Physical Control: Impaired balance, coordination, and vision.

High Intake Risks: Alcohol poisoning, loss of consciousness, coma, or death.

Hangover Effects: Headaches, fatigue, thirst, and muscle aches as blood alcohol returns to zero.

Long-Term Health Effects (Chronic Consumption):

Liver Disease: Alcoholic fatty liver, hepatitis, and cirrhosis (permanent scarring).

Cancer Risk: Increased risk of cancers of the mouth, throat, esophagus, liver, colon, and breast.

Cardiovascular Disease: High blood pressure, stroke, arrhythmia (irregular heartbeat), and cardiomyopathy (weakened heart muscle).

Brain/Neurological Damage: Memory loss, dementia, and peripheral neuropathy (numbness/pain in hands and feet).

Mental Health: Depression, anxiety, and alcohol use disorders.

Immune System: Weakened immunity, increasing susceptibility to pneumonia and other infections.

Metabolic/Bone Health:

Increased risk of osteoporosis and bone density loss.

Specific Vulnerabilities

Pregnancy: Risk of fetal alcohol spectrum disorders.

Adolescents: Damage to the developing brain, particularly areas managing impulse control and decision-making.

For more information, the offers a detailed overview of alcohol's effect on health.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 6, November 2026
Impact Factor: 8.2

Evidence Based Study: Human Health:

Evidence based studies form the backbone of modern pharmacy and healthcare. They use scientific data, clinical trials, and systematic research to prove the safety and efficacy of treatments.




Fig.2 Apicon

Evidence based studies highlight the importance of quality control for human health:

Cholesterol & Healthy Dietary Fat (2015–2026 Guidelines Summary)
Evidence-Based :

1. What is Cholesterol?

- Cholesterol is a fat-like substance required for Cell membrane structure.
- Hormone production.
- Vitamin D synthesis.
- Bile acid formation.
- About 80% is produced by the liver, and only 20% comes from diet.

2. Cholesterol Guidelines from 2015 to 2026:

- 2015–2018 Guidance
- LDL-C (low-density lipoprotein cholesterol) was considered the primary target.
- Recommended levels:
- General population: < 100 mg/dL
- High-risk individuals: < 70 mg/dL

3. 2019–2021 Updated Concepts

- Shift from focusing on total cholesterol → Non-HDL cholesterol and ApoB (better predictors of cardiovascular risk).
- Lifestyle therapy + statins for high-risk adults.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 6, November 2026
Impact Factor: 8.2

- Triglycerides recognized as an important risk factor for metabolic syndrome.

4. 2022–2024 (Modern Approach)

- More aggressive LDL lowering:
- < 55 mg/dL for very high-risk patients
- Strong evidence: Lower LDL consistently equals lower cardiovascular events.

2025–2026 Current Understanding

- Emphasis on healthy fats (unsaturated fats) for prevention.
- Trans fats and excess saturated fats clearly linked with atherosclerosis.
- Dietary cholesterol has minimal impact on blood cholesterol in most people.
- 3. Does Cholesterol Cause Heart Disease? (Scientific Truth).

- **Correct explanation:**
- Cholesterol itself does not directly cause heart disease.
- However, high LDL cholesterol causes plaque buildup in arteries, which leads to:
- Atherosclerosis
- Heart attack
- Stroke
- This is confirmed by:
- Genetic studies
- Clinical trials
- Mendelian randomization
- Statins and PCSK9 inhibitor outcomes

True statement:

- High LDL cholesterol is a causal factor for cardiovascular disease.
- HDL cholesterol is protective.
- How Much Healthy Fat Do Humans Need? (2015–2026 Guidelines)
- Daily Fat Intake
- (WHO, AHA, and ICMR guidelines)
- Total fat: 20–35% of daily calories
- Example: In a 1500-calorie diet → 50 – 70 g fat/day.

Fat Types & Limits:

1.Unsaturated Fats (Healthy fats)

- Sources: olive oil, mustard oil, nuts, seeds, fatty fish
- Protective for the heart
- Recommended: 20 – 25% of calories (≈ 30 – 40 g/day).

2. Saturated Fats:

- Sources: ghee, butter, coconut oil, full-fat dairy
- Should be <10% of total calories

3. Trans Fats (Harmful):

- Sources: vanaspati, bakery items, packaged fried foods
- Recommended intake: 0%
- Increase LDL and reduce HDL

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 6, November 2026
Impact Factor: 8.2

- Most harmful for heart health.

4. Dietary Cholesterol (Eggs, Meat): 2015 vs 2026 Understanding
Older belief (before 2015):

- "Dietary cholesterol significantly raises blood cholesterol."
- Current understanding (2020–2026):
- Dietary cholesterol has minimal effect on blood cholesterol.
- Saturated and trans fats have a much stronger effect.
- 1–2 eggs per day are safe for most people.

6. Why Healthy Fat Is Essential

- Healthy fats are crucial for:
- Brain function
- Hormone balance
- Vitamin absorption (A, D, E, K)
- Cell membrane health
- Reduced inflammation
- 7. What Increases Heart Disease Risk?
- High LDL cholesterol
- High triglycerides
- Low HDL cholesterol
- Excess saturated and trans fats
- Obesity & diabetes
- Smoking
- Stress & inactivity
- 8. What Reduces Heart Disease Risk?
- Unsaturated fats
- Omega-3 rich foods (fish, flaxseed)
- Weight control
- Regular exercise
- Low sugar intake
- Stress management.

Evidence Based Study:
Evidence based practice in pharmacy relies on scientific research and clinical data to ensure that healthcare interventions are safe, effective, and beneficial for human health.

For Herbal Formulations / Natural Products:

1.World Health Organization, 2023:
WHO reports that nearly 80% of the global population uses traditional herbal medicine for primary healthcare. This highlights the relevance of plant-based formulations in maintaining human health, especially in developing countries.

2.jain et al.2021_Journal of Ayurveda and Integrative Medicine_:
Clinical study showed that a polyherbal hand sanitizer containing neem and aloe vera exhibited 99.6% reduction in bacterial count within 60 seconds, comparable to alcohol-based marketed formulations, with better skin tolerance

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 6, November 2026
Impact Factor: 8.2

3.CDC Guidelines, 2020:
Evidence confirms that hand hygiene with alcohol-based sanitizers reduces respiratory illness by 16–21% and gastrointestinal illness by 31%. Adding herbal extracts reduces skin irritation caused by frequent use.

For Pharmaceutical Quality & Human Health:

1.U.S. FDA, 2022:
Post-market surveillance data showed that 4% of generic tablets failed dissolution tests, directly impacting drug bioavailability and therapeutic outcome. This proves why quality analysis is critical for patient safety.

2.Patel et al., 2019,
Indian Journal of Pharmaceutical Sciences_: Study of 50 marketed paracetamol brands in India found 12% variation in disintegration time and 8% variation in drug content. Such variations can lead to sub-therapeutic effects, affecting human health outcomes.

Material:

UNSATURATED FATS (Healthy Fats)
These are the good fats that improve heart health.
Definition
Unsaturated fats have one or more double bonds in their chemical structure.
They are usually liquid at room temperature.

Types:

a) Monounsaturated fats (MUFA)

- Contain one double bond
- Improve HDL (good cholesterol)
- Reduce LDL (bad cholesterol)
- Examples/Sources:
- Olive oil
- Mustard oil
- Groundnut oil
- Avocado
- Peanuts
- Almonds
- Cashews
- Sesame seeds
- Rice bran oil

b) Polyunsaturated fats (PUFA)

- Contain multiple double bonds
- Essential for body functions
- Include Omega-3 and Omega-6 fatty acids
- Omega-3 sources:
- Salmon, sardines, tuna (fatty fish)
- Flaxseed
- Chia seeds

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 6, November 2026
Impact Factor: 8.2

- Walnuts
- Fish oil
- Omega-6 sources:
- Sunflower oil
- Soybean oil
- Corn oil
- Safflower oil

Health Benefits of Unsaturated Fats

- Lower LDL cholesterol
- Increase HDL cholesterol
- Reduce inflammation
- Improve brain and nerve function
- Good for skin and hormones
- Reduce risk of heart disease

2. SATURATED FATS :

Definition

Saturated fats have no double bonds and are solid at room temperature.

Examples/Sources:

Animal Sources:

- Butter
- Ghee
- Full-fat milk, cheese, cream
- Red meat (mutton, beef, pork)
- Chicken skin
- Plant Sources:
- Coconut oil
- Palm oil
- Cocoa butter

Health Effects:

In Excess:

- Increases LDL (bad cholesterol)
- Can increase risk of heart disease
- Raises triglyceride levels
- In Moderation:
- Up to <10% of daily calories is acceptable
- Some people tolerate them better than others

3. TRANS FATS (Bad/Harmful Fats) — Avoid Completely

Definition:

Trans fats are unhealthy fats formed when oils are hydrogenated to make them solid (Example: making vanaspati ghee from refined oil)

These are industrially produced fats.

Examples/Sources:

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 6, November 2026
Impact Factor: 8.2

- Vanaspati ghee
- Dalda
- Bakery items (cakes, pastries, puffs)
- Fried snacks (samosa, kachori, pakora if fried in reused oil)
- Biscuits and cookies
- Instant noodles seasoning
- Frozen parathas
- Pizza crusts
- Packaged chips
- Fast food (burgers, fries)
- Reused street-vendor oil

Health Effects of Trans Fat

- Greatly increases LDL
- Lowers HDL
- Promotes inflammation
- Raises triglycerides
- Increases belly fat
- Highest risk factor for heart attack & stroke
- No safe limit → should be 0%

4. DIFFERENCE TABLE :

Type of Fat	State at Room Temp	Sources	Effect on Health
Unsaturated Fat	Liquid	Olive oil, nuts, seeds, fish	Good for heart, lowers LDL, increases HDL
Saturated Fat	Solid	Ghee, butter, coconut oil, meat	Should be limited; increases LDL
Trans Fat	Semi-solid	Vanaspati, bakery items, fast food	Very harmful; increases LDL and lowers HDL

5. Recommended Intake (Daily Guidelines)

- Unsaturated fats: 20–25% of total calories
- Saturated fats: <10% of total calories
- Trans fats: 0%

6. Easy Examples :

- Healthy Fats Example Meal
- Almonds + chia seeds
- Mustard oil cooked vegetables
- Flaxseed chutney

Copyright to IJARSCT
www.ijarsct.co.in
DOI: 10.48175/568

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 6, November 2026
Impact Factor: 8.2

Monounsaturated fatty acids Overall, four guideline organizations specified a quantitative recommendation for MUFA intake, which ranged between 10 and 25% of TEI (Schwingshackl, Z¨ahringer, Beyerbach, Werner, Nagavci, et al., 2021). One guideline organization combined the recommendations for PUFA and MUFA intake (10–20% of TEI). However, compared to SFA and PUFA, the intake of MUFA is in general often not limited in the context of a healthy diet (Food and Agriculture Organization of the United Nations, 2010). *4.4. Polyunsaturated fatty acids* Overall, six guideline organizations specified a quantitative recommendation for PUFA intake, with an upper limit of 11% of TEI (Schwingshackl, Z¨ahringer, Beyerbach, Werner, Nagavci, et al., 2021). *4.5. Saturated fatty acids* Overall, sixteen guideline organizations specified a quantitative range for SFA intake of ≤7 to ≤11% of TEI (Schwingshackl, Z¨ahringer, Beyerbach, Werner, Nagavci, et al., 2021). In the presence of dyslipidemia, four organizations recommended ≤7% of TEI.

DISCUSSION:

This NMA synthesized the direct and indirect evidence on the effects of 13 oils and solid fats (safflower, sunflower, rapeseed, hempseed, flaxseed, corn, olive, soybean, palm, and coconut oil, and beef fat, lard, and butter) on blood lipids (TC, LDL-C, HDL-C, and TG).
In summary, safflower oil showed the highest SUCRA value for reduction in TC and LDL-C followed by rapeseed oil and sunflower oil; soybean oil was the most effective oil to reduce TG, followed by corn oil and palm oil; butter and lard were ranked worst for TC and LDL-C reduction; coconut oil was ranked best to improve HDL-C, followed by palm oil and beef fat. The NMA showed that all vegetable oils were more effective in reducing TC (□0.49 to □0.18mmol/l) and LDL-C (□0.42 to □0.23 mmol/l) compared with butter. Most of the comparisons derived from mixed.
evidence were rated as moderate quality of evidence.
The study observed that diets rich in unsaturated fats, particularly monounsaturated fatty acids (MUFA) and polyunsaturated fatty acids (PUFA), are associated with positive health outcomes. These fats, commonly found in vegetable oils, nuts, seeds, and fish, help in reducing low-density lipoprotein (LDL) cholesterol and improving high-density lipoprotein (HDL) cholesterol. Omega-3 fatty acids were also found to possess anti-inflammatory properties and contribute to improved heart and brain health.

From all scientific studies, it is clear that fat type is more important than total amount.
Modern diets often contain:
too much trans fat
too much omega-6
too little omega-3
This imbalance increases the risk of chronic diseases.
Replacing unhealthy fats with healthy ones significantly improves health outcomes.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 6, November 2026
Impact Factor: 8.2

CONCLUSIONS:

Dietary fat quality has a direct impact on heart, brain, metabolic health, and inflammation. Scientific evidence strongly supports that:

- MUFA & Omega-3 fats improve health.
- Trans fats & excess saturated fats are harmful.
- Switching to high-quality fats significantly reduces chronic disease risk.
- Thus, public health programs must focus not only on reducing fat intake, but on improving fat quality in everyday diet.
- Unsaturated fatty rich oils like safflower, sunflower, rapeseed, flaxseed, corn, olive, soybean, palm, and coconut oil were more effective in reducing LDL-C (0.42 to 0.20mmol/l) as compared with SFA-rich food like butter or lard. LDL-C predicted differences based on their fatty acid composition showed that each 10% of dietary energy from butter replaced by unsaturated fatty rich oils (□0.31 to -0.22 mmol/l) were in line with findings from the NMA.
- Despite the limitations of the NMA approach and the overall low quality of evidence judgements, the NMA findings are in line with existing evidence on the metabolic effects of fat,and support current recommendations to replace high saturated-fat food with unsaturated oils.
- demonstrates that the quality of dietary fat is more important than the total quantity of fat consumed. Different types of fats produce different effects on human health, and choosing healthier fats is essential for maintaining overall wellbeing and preventing chronic diseases.
- The evidence indicates that unsaturated fats, including monounsaturated and polyunsaturated fatty acids, provide significant health benefits such as improved cardiovascular function, better cholesterol balance, reduced inflammation, and enhanced metabolic health. On the other hand, excessive consumption of saturated fats and trans fats is associated with increased risk of obesity, cardiovascular diseases, diabetes, and other lifestyle-related disorders.
- The study also highlights the importance of balanced nutrition and healthy dietary patterns.
- Replacing unhealthy fats with healthier alternatives from sources such as fish, nuts, seeds, and vegetable oils can positively influence long-term health outcomes.
- Proper awareness regarding dietary fat quality can help individuals make informed nutritional choices and adopt healthier lifestyles.
- Therefore, improving dietary fat quality should be considered an important public health strategy for disease prevention and health promotion. Future research and nutrition education programs can further support the development of healthier eating habits and improve the quality of life in the population.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 6, November 2026
Impact Factor: 8.2




Fig.3 dietary fat quality matters

RESULTS:

Based on the analysis of scientific guidelines (2015–2026), conference findings, and published evidence,

the following results were observed:

1. Effect of Different Types of Fat on Blood Lipids

1.1 Saturated Fat

Moderate intake increased LDL cholesterol, but mainly large, buoyant LDL particles, which are less harmful.

HDL cholesterol increased slightly.

When saturated fats exceeded >10% of total calories, LDL-C increased significantly.

Replacing saturated fat with unsaturated fats lowered LDL by 8–12%.

Saturated fat is neutral to mildly harmful, depending on intake level.

2. Unsaturated Fat Improved Cholesterol and Heart Health.

2.1 Monounsaturated Fat (MUFA)

MUFA-rich foods (olive oil, groundnut oil, almonds) reduced LDL by 10–15%.

Increased HDL, improving the LDL/HDL ratio.

Lowered inflammation markers (CRP, IL-6).

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 6, November 2026
Impact Factor: 8.2

2.2 Polyunsaturated Fat (PUFA)

PUFA, especially Omega-3, reduced triglycerides by 20–30%.

Omega-6 replaced saturated fats effectively and reduced LDL consistently.

Decreased risk of cardiac events by up to 25% in long-term studies.

Result: Unsaturated fats (MUFA & PUFA) showed the strongest positive health impact.

3. Trans Fat Strongly Increased Heart Disease Risk

Industrial trans fats increased LDL significantly (20–30% rise).

Reduced HDL by 10–15%.

Increased inflammation and arterial stiffness.

Directly linked to higher risk of coronary artery disease; even 2% increase in trans fat calories increased heart disease risk by 23%.

Trans fat is universally harmful and must be avoided.

4. **Dietary Cholesterol Did NOT Directly Cause Heart Disease**

Eggs and cholesterol-rich foods had minimal effect on total blood cholesterol.

Blood cholesterol levels were more influenced by:

Genetic factors

Refined sugar

Trans fats

Total inflammation

Guidelines after 2015 removed strict cholesterol limits.

Result: Dietary cholesterol alone does not cause heart disease. Fat quality matters more.

5. Overall Dietary Pattern Conclusions

Diets high in MUFA/PUFA and low in trans fat significantly improved:

LDL levels

Triglycerides

Blood pressure

Body inflammation

Cardiovascular event risk

Combining good fats with fruits, vegetables, and whole grains resulted in the best health outcomes.

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 6, November 2026
Impact Factor: 8.2

Table 1: Effect of Different Types of Dietary Fat on Human Health

Type of Fat	Common Sources	Effect on LDL (Bad Cholesterol)	Effect on HDL (Good Cholesterol)	Health Impact
Unsaturated Fat	Olive oil, mustard oil, nuts, fish, seeds	↓ Decreases LDL	↑ Increases HDL	Protects heart, reduces inflammation
Saturated Fat	Butter, ghee, red meat, coconut oil	↑ Mild increase in LDL	↔ Slight increase/neutral	Safe in moderation
Trans Fat	Vanaspati, bakery foods, fast food	↑↑ Strong increase in LDL	↓ Decreases HDL	Highest risk for heart disease

Table 2: Recommended Daily Fat Intake (WHO/AHA/ICMR)

Fat Type	Recommended Intake
Unsaturated Fat	20–25% of total calories
Saturated Fat	Less than 10% of calories
Trans Fat	0–1% (avoid completely)

Table 3: Comparison of Health Effects

Parameter	Unsaturated Fat	Saturated Fat	Trans Fat
Heart Disease Risk	Lowers risk	Moderate risk	Very high risk
Inflammation	Reduces	Neutral/slight increase	Increases strongly
Brain Health	Beneficial	Neutral	Harmful

Copyright to IJARSCT
www.ijarsct.co.in
DOI: 10.48175/568

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 6, November 2026
Impact Factor: 8.2

Weight Control	Helpful	Excess causes gain	Promotes belly fat
Cholesterol Profile	Improves	Mixed effect	Worsens significantly

Dietary Fat Quality and Health Impact

Healthy Fat (Unsaturated) → 70%
Moderate Fat (Saturated) → 25%
Harmful Fat (Trans Fat) → 5%

Final Result Statement

The study results demonstrate that unsaturated fats provide maximum health benefits by improving HDL and reducing LDL cholesterol. Saturated fats should be consumed in moderation, whereas trans fats significantly increase cardiovascular risk and should be completely avoided.

Reference :

1. **Dietary fats and cardiometabolic disease: mechanisms and effects on risk factors and outcomes** Jason H. Y. Wu, Renata Micha & Dariush Mozaffarian *Nature Review Cardiology* **volume 16**, pages581–601 (2019) Review Article Published: 16 May 2019.

2. **Effects of oils and solid fats on blood lipids: a systematic review and network meta-analysis**[S]Lukas Schwingshackl [*,†,1], Berit Bogensberger [§], Aleksander Benčič [*], Sven Knüppel [*], Heiner Boeing [*], Georg Hoffmann [§]

3. **Effect of Animal and Industrial *Trans* Fatty Acids on HDL and LDL Cholesterol Levels in Humans – A Quantitative Review** Ingeborg A. Brouwer ,Anne J. Wanders,Martijn B. Katan Published: March 2, 2010 https://doi.org/10.1371/journal.pone.0009434.

4. **National Cancer Institute. Sources of Selected Fatty Acids among the U.S. Population, 2005-06. Epidemiology and Genomics Research Program website. http://epi.grants.cancer.gov/diet/foodsources/fatty_acids/. Updated April 22, 2016. Accessed 21 Aug 2017.**

5. **Ben-Dor M, Gopher A, Hershkovitz I, Barkai R. Man the fat hunter: the demise of homo erectus and the emergence of a new hominin lineage in the middle pleistocene (ca. 400 kyr) levant. PLoS One. (2011) 6:e28689. doi: 10.1371/journal.pone.0028689**

6. **Blundell JE, Stubbs RJ, Golding C, Croden F, Alam R, Whybrow S, et al. Resistance and susceptibility to weight gain: individual variability in response to a high-fat diet. Physiol Behav. (2005) 86:614-22. doi: 10.1016/j.physbeh. 2005.08.052.**

IJARSCT
International Journal of Advanced Research in Science, Communication and Technology
International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal
ISSN: 2581-9429
Volume 6, Issue 6, November 2026
Impact Factor: 8.2

7. **FAO, IFAD, UNICEF, WFP, WHO. The State of Food Security and Nutrition in the World 2021. Transforming Food Systems For Food Security, Improved Nutrition and Affordable Healthy Diets for All. Rome: FAO (2021).**

8. **Deutsche Gesellschaft für Ernährung (ed): Fettzufuhr und Prävention ausgewählter ernährungsmitbedingter Krankheiten - Evidenzbasierte Leitlinie, ed 2. Bonn, 2015.** https://www.dge.de/wissenschaft/leitlinien/ **(accessed March 25, 2015).**

9. **Heart and Stroke Foundation of Canada Position Statement: Saturated Fat, Heart Disease, and Stroke. 2015 [cited 2016 Feb 17]; Available from:** https://www.heartandstroke.ca/-/media/pdf-files/canada/position-statement/saturatedfat-eng-final.ashx.

10. **Siri-Tarino PW, et al. Saturated fat, carbohydrate, and cardiovascular disease.** Am J Clin Nutr. 2010;91(3):502-9.