

# Diversity and Distribution of Agricultural Insect Pest in Some Selected Areas of Indapur (Pune) and Phaltan (Satara) Tehsil, Maharashtra, India

Rupesh B. Yadav<sup>1</sup>, Prajakta D. Khaire<sup>2</sup>, Suryakant V. Maske<sup>3</sup>

Thakur College of Science & Commerce, Kandivali East, Mumbai, Maharashtra, India<sup>1</sup>

Tuljaram Chaturchand College Arts, Commerce & Science, Baramati, Maharashtra, India<sup>2</sup>

Shri Vijaysinha Yadav Arts & Science College, Peth Vadgaon, Kolhapur, Maharashtra, India<sup>3</sup>

**Abstract:** A survey was conducted to investigate the agricultural insect pest diversity in Indapur and Phaltan Tehsil. The actual survey was carried out during 15 Sep. 2020 to 15 March 2021. Throughout study period, total 25 species of agricultural insect pests were collected & identified from 29 different sampling sites of 7 villages. They are belonging to 4 orders that is Orthoptera, Hemiptera, Coleoptera and Lepidoptera, while included in 12 families that is Gryllidae, Gryllotalpidae, Acrididae, Tettigoniidae, Pyrgomorphidae, Pentatomidae, Pseudococcidae, Coreidae, Chrysomelidae, Scarabaeidae, Cupedidae, and Sphingidae. The reported species were recorded from different habitat like Agricultural field, Residential area, Water Reservoirs, Woodland & Forest, Thorny Scrub and Grassland area. According to IUCN Red list, 1 species is critically Endangered, 2 species are Endangered, 1 species is Vulnerable, 5 species are Near Threatened, 14 species are Data deficient, and 2 species are Least Concern. The insect pest cause on an average 15–20% yield losses in food and cash crops. Due to heavy of Insecticides use in agricultural field, the natural biocontrol agent's numbers are drastically declined in study area. Suggestions for future improvement include, Integrated Pest Management (IPM) program through the ecological pest management practices.

**Keywords:** Diversity, Distribution, Agricultural Insect Pest, Indapur, Phaltan, etc.

## I. INTRODUCTION

India is the world's largest popular agriculture countries in the globe and the largest producer of crop wheat, rice, sugarcane, vegetables, groundnut, fruits and cotton. Food plants of the world are damaged by more than 10,000 species of insects, 100,000 diseases, 30,000 species of weeds and 1000 species of nematodes (Hall, 1995; Dhaliwal et al., 2007). Insect cause damage to plants either directly or indirectly in their attempt to source food, and almost all the portions, viz. Roots, bark, shoots, stem, leaves, buds, flowers and fruits of plant are attacked and injured by insects (Atwal and Dhaliwal, 2015). Many of the insect pests such as aphids, caterpillar, grasshopper, locusts, whiteflies, leafhopper, mole cricket, thrips and some bugs etc. are damage the crop. There are many different types of insect pest included in different orders.

Losses because of the insect pest in Indian agriculture have been assessed from time to time (Singh et al., 2014; Dhaliwal & Arora, 2015). Extensive surveys carried out during early 1960s revealed that fruit, cotton, rice, and sugarcane suffered 25, 18, 10 and 10% yield losses, respectively (Pradhan, 1964). The number of insect pest damage the crop and loss of the yield. The agriculture field of India is presently suffering an annual loss of about Rs. 8, 63,884 million due to insect pest (Dhaliwal et al., 2010). The heavy crops are loss from insect pests, the farmers practice large amounts of pesticides (Aktar et al., 2009). But, both the quantity of food loss due to pests and the cost of pest control in terms of money and human health are significant (Pimentel and Greiner, 1997). Sucking insect pests and defoliators like mirid bug, whitefly, aphids, mealy bug, plant hoppers, shoot fly, and the defoliating tobacco caterpillar, leaf miner and leaf folder *Cnaphalocnoides medinalis* (Guenee) have emerged as major pests (Chakra-barty, 2015).

Insects have a directly influence on agricultural food production by sucking out plant juices, chewing the leaves of crop

plants, stems or leaves and spreading plant pathogens, boring within the roots. They feed on natural fibers, destroy wooden building materials, ruin stored grain and accelerate the process of decay. (Yang et al. 2014) Pest control is the best achieved with an integrated pest management plan using a range of biological, chemical, physical, mechanical or cultural control methods. Selective insecticides can be employed with fewer effects on natural enemies. Common predators of insect pest include praying mantis, ladybird beetles, spiders, true bugs, earwigs, ground beetles, hoverfly larvae and lacewings (Cork, A., Dobson, et al., 2009).

Considering the above importance of the study, the present investigation was undertaken to know the diversity and distribution of agricultural insect pest in some selected areas of Indapur (Pune) and Phaltan (Satara) Tehsil.

## II. MATERIALS AND METHODS

### A. Study Area-

Indapur is located at 18.1187° N, 75.0234° E. It has an average elevation of 527 meters. and Phaltan is located at 17.9935° N, 74.4304° E. It has an average elevation of 558 meters. The climate of Indapur is quiet hot place as it receives scanty rainfall an inland climate of Maharashtra. The Phaltan climate is an inland of Maharashtra. The temperature is relatively high range between 12°C to 45°C. The observation of agricultural insect pest was carried out in twice a day (morning and evening) from two places of sampling site of Indapur and Phaltan Tehsil. The total study period 15 Sep. 2020 to 15 March 2021. The places include Dalaj no. 1, Dalaj no.2, Dalaj no. 3, Kalewadi, Bhadalwadi, Pilewadi, Bhigwan, Takalwade, Pimprad, Nimblak, area of Indapur and Phaltan city.

### B. Sampling Site-

Throughout the Indapur and Phaltan Tehsil, extensive survey was carried out to cover maximum study area, actual collection sites shown in table No. 1- collection site along with their GPS location and elevation of Indapur and Phaltan city.

### C. Collection of the Insect-

The regular collection of agricultural insects made during period of 15 Nov 2020 to 15 March 2021 using hand picking and insect net. The photography of the specimen was carried out by using DSLR camera Canon 760 D with 18-55 lance. The insects were anesthetized using jar containing cotton wad dipped chloroform. The insects then sun-dry & preserved and entomological pins were used for spread of the insect on entomological board.

### D. Taxonomic Identification-

Agricultural insect pest identification was done by using the available literature, Research article and identification keys.

## III. RESULT AND DISCUSSION

During present study, a total 25 species of agricultural insect pest belonging to 4 order and 12 family were collected and identified (Table No. 2). Out of 25 species specimen, 1 species belonging to family Gryllidae. In order Orthoptera, 15 species from family Gryllotalpidae, 3 species from family Acrididae, 7 species from family Tettigoniidae, and 2 species from family Pyrgomorphidae, 3 species from family Pentatomidae.

**Table No. 1-** Agricultural insect pests sampling sites along with GPS location and elevation.

Sr. No.	Name of Village	Habitat	Latitude	Longitude	Elevation (In Meters)
1	Dalaj No. 3	Residential area	18°13'33.9"N	74°49'29.1"E	510
2	Dalaj No. 3	Grassland area	18°13'43.6"N	74°49'34.1"E	505
3	Dalaj No. 3	Residential area	18°13'45.6"N	74°49'39.2"E	503
4	Dalaj No. 3	Agricultural area	18°13'53.3"N	74°49'40.0"E	500
5	Dalaj No. 3	Agricultural area	18°13'37.6"N	74°49'42.7"E	503
6	Dalaj No. 3	Agricultural area	18°13'28.3"N	74°49'31.1"E	509
7	Dalaj No. 1	Grassland area	18°14'00.9"N	74°48'54.4"E	507

8	Dalaj No.1	Grassland area	18°14'24.0"N	74°48'48.1"E	502
9	Dalaj No. 1	Residential area	18°14'03.1"N	74°49'12.4"E	502
10	Dalaj No. 2	Agricultural area	18°14'04.0"N	74°48'11.6"E	505
11	Dalaj No. 2	Agricultural area	18°13'27.8"N	74°47'59.7"E	529
12	Dalaj No. 2	Grassland area	18°13'59.4"N	74°48'15.8"E	508
13	Kalewadi	Residential area	18°24'27.8"N	74°41'00.3"E	515
14	Pilewadi	Agricultural area	18°12'20.2"N	74°47'20.5"E	558
15	Bhadalwadi	Agricultural area	18°13'39.0"N	74°46'13.5"E	526
16	Bhadalwadi	Grassland area	18°13'45.8"N	74°46'27.3"E	522
17	Bhadalwadi	Agricultural area	18°13'45.8"N	74°46'21.3"E	521
18	Bhadalwadi	Residential area	18°13'40.1"N	74°46'28.0"E	521
19	Bhigwan	Residential area	18°18'06.9"N	74°45'18.0"E	507
20	Takalwade	Residential area	17°59'35.3"N	74°33'10.2"E	547
21	Takalwade	Agricultural area	17°59'29.1"N	74°33'06.6"E	547
22	Takalwade	Residential area	17°59'41.9"N	74°33'12.2"E	545
23	Takalwade	Grassland area	17°59'39.2"N	74°33'21.2"E	546
24	Nimblak	Agricultural area	19°09'15.0"N	74°40'16.7"E	654
25	Nimblak	Grassland area	19°09'16.0"N	74°40'34.1"E	646
26	Nimblak	Agricultural area	19°09'17.1"N	74°40'42.8"E	644
27	Pimprad	Agricultural area	17°58'59.8"N	74°31'47.0"E	558
28	Pimprad	Agricultural area	17°58'40.5"N	74°32'00.1"E	559
29	Pimprad	Residential area	17°58'43.0"N	74°31'53.3"E	559

In order Hemiptera, 2 species from family Pseudococcidae, 1 species from family Coreidae. 1 species from family Chrysomelidae. In order Coleoptera, 2 species from family Scarabidae, 1 species from family Cupedidae. 1 species from family Spingidae. In order Lepidoptera were collected. The photographs of the observed agricultural insect pest are given in plate.

In Order Orthoptera, *Telegryllus emma* commonly known as a field cricket is black to dark brown insect, about 25mm long. They have large heads and wings that are folded flat against the black. They have long antennae and two hairy prongs sticking out of the end of the abdomen. In family Gryllotalpidae, *Neocurtilla hexadactyla* are called northern mole cricket. The northern mole cricket has yellowish brown with a dark prothorax. 3 or 4 claws, short wings, long tail like extension and brawny legs. In family Acrididae, *Phaulacridium marginale* are called New Zealand grasshopper. Wings are small, size were often unfortunately small. They are colored to match background vegetation and three pairs of leg. *Schistocerca americana* is also called as American grasshopper. The body of this pest generally yellow brown in color. The wings are pale with large brown spot. They are slender grasshopper with long wings. White strips on the tegmina, pronotum and head are distinct. Another grasshopper *Oedaleus infernalis* shows cylinder body shape. This pest usually has a light brown body with dark brown stripes and small size. In family Tettigoniidae, *Microcentrum rhombifolium* shows pale green color. Veins on the wings look like the veins on a leaf. The wings are longer than the body and overlap each other when closed. The tip of the wings comes to a point much like a willow leaf. *Pterophylla camellifolia* are leaf green in color, length from 1.5 to 2.5 inches, long antennae and their veins on their oval shaped.

**Plate No. 1 - Photographs of Agricultural Insect Pest from Study Area**



The *Tettigonia viridissima* commonly known as Great green bush cricket are green with orangey brown stripes running the length of the body, long wings. It can be up to 3-4 cm long. Antennae are very long and thin. The *Letana intermedia* shows green color with or without dark, medium sized, slender, antennae and 3 pairs of leg are present. *Conocephalus maculatus* are smaller size, slanting shaped head, antennae are long, tegmina and wings are white in color and without any spots.

In family Pyrgomorphidae, *Chrotogonus oxypterus* are brown pale, ventrally with black spots on thorax and metasoma. Antennae darker towards tip, hind femur with black spots. Head is sub conical, straight, cerci short, conical, stumpy. *Tagasta apludi* shows yellowish brown in color and brown spots in body. Antennae are short, 3 pairs of leg and head is pointed.

The survey was conducted to check the abundance of agricultural insect pest in Indapur and Phaltan Tehsil. The present study indicates the agricultural insect pests are increase in number as compared to biological agents. In order Orthoptera species was highly observed as compared to order Hemiptera, Coleoptera, Lepidoptera. The most of the order Hemiptera in green stink bug are found in maize plant. The order Coleoptera in different species found in different areas. Some caterpillars are found in plants and vegetables.

In farmer's point of view agricultural insect pest like grasshopper, leafhopper, army worm, locusts, beetles, bugs, some caterpillar damages the agricultural crops and forestry causing significant losses to farmers. Examples of agricultural insect pest like brown marmorated stink bug can cause widespread damage to fruit and vegetable crops. Aphids are damaging crops by the direct action of sucking sap, but also harming them. Insects are responsible for two major forms of damage to crops. First, there is the direct injury they cause to the plants as they feed on tissues. Second, there is the indirect damage in which the insect itself does little or no harm but transmits a bacterial viral, fungal infection into a crop. So, pesticides and herbicides are applied to agricultural land to control pests that disrupt crop production. Pesticide can be toxic to a host of other organisms like beneficial insects and non-target plants and also damage soil biomass. They are best for the agricultural use of vermicompost organic fertilizer production and also the conservation of biocontrol agents.

#### IV. SUMMARY AND CONCLUSION

A survey was conducted to investigate the diversity and distribution of Agricultural insect pest in Indapur and Phaltan Tehsil. The actual survey was carried out during 15 Sep. 2020 to 15 March 2021. Throughout study period, total 25 species of agricultural insect pests were collected & identified from 29 different sampling sites of 7 villages. They are belonging to 4 orders and 12 families. The reported species were assessed on IUCN Red list to know the status. The insect pest cause on an average 15–20% yield losses in food and cash crops. Due to heavy of Insecticides use in agricultural field, the natural biocontrol agent's numbers are drastically declined in study area.

Suggestions for future improvement include, Integrated Pest Management (IPM) program through the ecological pest management practices.

**Table 2:** Checklist of Agricultural Insect Pests in Indapur and Phaltan Tehsil.

Sr. No.	English Name	Scientific Name	IUCN Status	Habitat
	<b>Order – 1 Orthoptera</b>			
	<b>Family – 1 Gryllidae</b>			
1	Emma field cricket	<i>Teleogryllus emma</i>	EN	GA & RA
	<b>Family - 2 Gryllotalpidae</b>			
2	Northern mole cricket	<i>Neocurtilla hexadactyla</i>	DD	WR & RA
	<b>Family – 3 Acrididae</b>			
3	New Zealand grasshopper	<i>Phaulacridium marginale</i>	DD	GA
4	American grasshopper	<i>Schistocerca americana</i>	DD	GA & WF
5	Grasshopper	<i>Oedaleus infernalis</i>	EN	GA
	<b>Family – 3 Tettigoniidae</b>			

6	Broadwinged katydid	<i>Microcentrumrhombifolium</i>	DD	TS & WF
7	Great green bush cricket	<i>Tettigonia viridissima</i>	LC	GA& AF
8	Bush cricket	<i>Letana intermedia</i>	DD	GA
9	Spotted meadow katydid	<i>Conocephalus maculatus</i>	LC	GA & RA
10	True katydid	<i>Pterophylla camellifolia</i>	CR	TS & WF
	<b>Family – 4 Pyrgomorphidae</b>			
11	Deccan grasshopper	<i>Chrotogonus oxypterus</i>	DD	WF
12	Grasshopper	<i>Tagasta apludi</i>	DD	GS & RA
	<b>Order – 2 Hemiptera</b>			
	<b>Family – 5 Pentatomidae</b>			
13	Brown marmorated stink bug	<i>Halyomorpha halys</i>	DD	AF & RA
14	Green stink bug	<i>Chinavia hilaris</i>	NT	AF, WF & RA
15	Spined soldier bug	<i>Podisus maculiventris</i>	NT	AF & RA
	<b>Family – 6 Pseudococcidae</b>			
16	Cotton mealy bug	<i>Phenacoccus solenopsis</i>	DD	AF
17	Red cotton stainer	<i>Dysdercus cingulatus</i>	DD	AF & RA
	<b>Family – 7 Coreidae</b>			
18	Western conifer seed bug	<i>Leptoglossus occidentalis</i>	NT	
19	Squash bug	<i>Anasa tristis</i>	DD	AF
	<b>Order -3 Coleoptera</b>			
	<b>Family – 8 Chrysomelidae</b>			
20	Leaf beetle	<i>Monolepta signata</i>	NT	AF
	<b>Family – 9 Scarabaeidae</b>			
21	Southern masked chafer	<i>Cyclocephala lurida</i>	DD	AF & WF
22	African black beetle	<i>Heteronychus arator</i>	DD	AF & GA
23	Coconut rhinoceros beetle	<i>Oryctes rhinoceros</i>	VL	AF, WF & RA
	<b>Family – 10 Cupedidae</b>			
24	Beetle	<i>Tenomerga cinerea</i>	DD	AF
	<b>Order – 4 Lepidoptera</b>			
	<b>Family – 11 Sphingidae</b>			
25	Oleander hawk moth	<i>Daphnis nerii</i>	NT	AF & WF

**Abbreviations**

LC – Least Concern

VL – Vulnerable

RA – Residential area

TS – Thorny Scrub

CR – Critically Endangered

EN – Endangered

WR – Water Reservoirs

GA – Grassland area

NT – Near Threatened

DD – Data Deficient

WF – Woodland and Forest

AF – Agricultural field

**REFERENCES**

- [1] Agarwala B K (1979), "Some aspects of aphid (Homoptera: Insecta), studies in Sikkim and Bhutan," Ph. D. thesis University of Calcutta, India, P.383.
- [2] Ambrose, E (1984) Research and development in banana crop protection (excluding sigatoka in the English-speaking Caribbean). *Fruits* 39, 234- 47.
- [3] Arora, R. and Dhawan, A K. 2013 Climate change and insect pest management Dhawan,
- [4] K., Singh, B., Bullar, M. B. and Arora, R. (eds). *Insect Pest Management*. Scientific publishers, Jodhpur. PP. 44-60.
- [5] Bana J K, Sharma H, Kumar Sushil, Singh P. Impact of weather parameters on population dynamics of oriental fruit fly, *Bactrocera dorsalis* (Hendel) (Diptera: Tephritidae) under South Gujarat mango ecosystem. *Journal of Agrometeorology*. 2017; 19: 78- 80.
- [6] Chakrabarty, P.K. 2015. Pest Management for sustainable agriculture. Sharma, D. R., Kumar, S. and Singh, (Eds). *Entomology for sustainable agriculture. Invited papers—Fourth congress on Insect science*. April, 16- 27, INSAIS, PAU, Ludhiana. P.1.
- [7] Cork, A., Dobson, H., Grzywacz, D., Hodges, R., Orr, A. and Stevenson, P. (2009) Review of pre- and post- Harvest Pest Management for pulses with special References to East and Southern Africa. Natural Resources Institute, University Extension, East Lansing.
- [8] CPCRT, 2015. CPCRT „Red Alert“ against Invasive pests of coconut. *The New Indian Express*, 6 January, Kerala, 2015. PP. 1- 3.
- [9] DAE (Department of Agricultural Extension). 2011. *Krishi Dairy 2011*. Published by Agriculture Information Service, Kharmarbari, Dhaka. PP. 214.
- [10] Dhaliwal, G.S., Arora, R. and Dhawan, A. K. (2004) crop losses due to insect pests in Indian agricultural: An update. *Indian j. Ecol.* 31(1): 1- 7.
- [11] Dillon, M.E., Wang, G. and Huey, R.B. 2010. Global metabolic impacts of recent climate warming. *Nature*, 467: 704 – 707.
- [12] Fletcher TB. *Some South Indian Insects*. Superintendent Government Press, Madras. 1914, 565.
- [13] Lanjar, A.G., Talpur, M.A., Khuhro, R.D. and Qureshi, K.H. 2002. Occurrence and abundance of grasshopper species on rice. *Pakistan Jour of Applied Sci.* 2 (7): 763 – 767.
- [14] Lee, J.C.T. and Landis, D.A. (2001) *Natural Enemies in field crops: A Guide to biological control*. Michigan State University Extension, East Lansing.
- [15] Nagrare VS, Kranthi S, Biradar VK, Zade NN, Sangode V, Kakde G. *Phenacoccus solenopsis* Tinsley (Hemiptera: Pseudococcidae), on cotton in India. *Bulletin of Entomological Research*. 2009; 99:537- 541.
- [16] Nair, M.R.G.K. 1970. *Insects and mites of crops in India*. New Jack printing works private Limited, Bombay. PP.32.
- [17] Palaniswami, MS, Pillai KS, Nair RR, Mohandas C. A new cassava pest in India. *Cassava Newsletter*. 1995; 19:6-7.
- [18] Patel KB, Saxena SP, Patel KM. Fluctuation of fruit fly-oriented damage in mango in relation to major abiotic factors. *Hort Flora Research Spectrum*. 2013; 2: 197- 201.
- [19] Pimentel D. In: Pimentel, D. (Ed.), *Biological invasions economic and environmental costs of alien plant, animal and microbe species*, second ed. CRC Press, Boca Raton, Florida, 2011, 463.
- [20] Pradhan, S. (1964) Assessment of losses caused by insect pests of crops and estimation of insect population. In: N.C. Pant (ed.) *Entomology in India*, Entomological Society of India, New Delhi, PP.17- 58.
- [21] Prasanna BM, Joseph Huesing E, Regina Eddy, Virginia M.P. *Fall armyworm in Africa: A Guide for Integrated Pest Management*, First Edition. Mexico, 2018.
- [22] Rejmanek M, Richardson DM. What makes some conifers more invasive? *Proceedings of the Fourth International conifer conference*, 2000.
- [23] Usmani M.K., Nayeem M.R. and Akhtar M.H., 2012. Field observations on the incidence of Grasshopper fauna as a pest of paddy and pulses. *European Journal of Experimental Biology*, 2 (5): 1912- 1917.
- [24] Varghese Devi. Seasonality and sampling of the mango shoot borer, *Chlumetia transversa* Walker (Lepidoptera: Noctuidae) pest Management in horticultural Ecosystems 1998; 4:16-20.

- [25] Viraktamath, C.A.2002. Alien invasive insects and mite pests and weeds in India and their management. *Microensica supplementia*, 6:67- 83.
- [26] Aldrich, J. A. 1988 Chemical ecology of the heteroptera. *Annual Review of Entomology*33:211-238.
- [27] Lamb." *Economic Entomology in the tropics*". Academic press (1974): 195.