

# Formulation and Evaluation of Antimicrobial Polyherbal Hand Wash

Jadhav Trupti<sup>1</sup>, Gaikwad Rutuja<sup>2</sup>, Navale Dnyaneshwari<sup>3</sup>, Tamboli Kashyab<sup>4</sup>, Khaire Akshada<sup>5</sup>

Students, Samarth Institute of Pharmacy, Belhe, Maharashtra, India<sup>1,3,4,5</sup>

Department of Pharmacology, Samarth Institute of Pharmacy, Belhe, Maharashtra, India<sup>2</sup>

**Abstract:** The aim of present work was to prepare formulations of polyherbal handwash from the metholonic extracts of leaves of *Mimosa pudica* (touch me not), *Azadirachta indica* (Neem) and lemon juice. Two formulations of hand wash were prepared and the formulations were evaluated for physical property like appearance pH and viscosity. The antimicrobial activity of prepared formulations of hand wash was checked against skin pathogens *Bacillus subtilis*, *Staphylococcus aureus*, *Psuedomonas aeruginosa* and *Escherichia coli* by agar diffusion method. The results revealed that prepared herbal hand wash formulations showed significant zone of inhibition compared with standard antibiotic drug (Amoxicillin). So these plant materials can be used in the preparation of herbal hand wash on commercially scale.

**Keywords:** Polyherbal hand wash ,antimicrobial activity, mimosa pudica and azadirachta indica

## I. INTRODUCTION

Skin being the most exposed part of our body requires protection from skin pathogens. The hands of Health Care workers (HCWs) are the primary routes of transmission of multidrug resistant pathogens and infection to the patients. Hence, it brings up the use of antiseptic for hand wash purpose. Many of the chemical antiseptics are now available in market as alcohol based sanitizers, chlorhexidine products etc. These soaps or solutions help to reduce health care associated transmission of contagious diseases more effectively but they have some shortcomings or adverse effects. Their frequent use can lead to skin irritation and also resistant among pathogens. Organisms such as *Staphylococcus aureus*, *Pseudomonas* spp., *Klebsiella pneumonia* & *Proteus vulgaris* are some of the skin pathogens<sup>1</sup>. Hand washing is an important way to help fight the spread of disease. Hand washing removes visible dirt from hands and reduce the number of harmful microorganisms. Harmful bacteria and viruses such as, *E. coli* and *Salmonella* can be carried by people, animals or equipment and transmitted to food<sup>2</sup>

*Mimosa pudica* L. (Mimosaceae) also referred to as touch me not, live and die, shame plant and humble plant is a prostrate or semi-erect subshrub of tropical America and Australia, also found in India heavily armed with recurved thorns and having sensitive soft grey green leaflets that fold and droop at night or when touched and cooled. It majorly possesses antibacterial, antivenom, antifertility, anticonvulsant, antidepressant, aphrodisiac, and various other pharmacological activities. The herb has been used traditionally for ages, in the treatment of urogenital disorders, piles, dysentery, sinus, and also applied on wounds.



Figure: *Mimosa pudica* L.



Figure: *Azadirachta indica* A.

***Azadirachta indica*** A. Juss, (Neem tree), from the Meliaceae family, also known as Margosa or Indianlilac. Various parts of the Neem tree have been used as traditional Ayurvedic medicine in India. Neem oil, the bark and leaf extracts have been therapeutically used as folk medicine to control leprosy, intestinal helminthiasis, respiratory disorders and constipation and also as a general health promoter. Neem leaves possess a wide spectrum of antibacterial action against gram-negative and gram-positive microorganisms.

**Lemon Juice:** which is obtained from fruits of *Citrus limon* L. belonging to the family *Rutaceae* is traditionally used for the purpose of cleaning due to its disinfectant properties. Lemon juice is also used as a short term preservative in some food preparations. Lemon juice is used in Indian medicinal systems because of the anti microbial properties of lemon. It is also used to add taste to many food preparations.

## II. MATERIAL AND METHOD

### Collection of Plant Material

The plant of *Mimosa pudica* L and *Azadirachta indica* A were collected from the garden area of NGSMIPS, PUNE

### Preparation of Herbal Leaf Extract

The collected plants *Mimosa pudica* L and *Azadirachta indica* A leaves are taken and coarsely powdered. 10 grams of coarsely powdered leaves of both plants were soaked in 200 ml of methanol and kept for maceration for about 3-5 days. After maceration the extract is filtered and the filtrate was collected and used for making hand wash.

### Preparation of Herbal Hand Wash Formulation

#### Formulation 1(F-1)

In this formulation the hand wash was prepared using 20 ml of methanolic extract filtrate. To this this formulation was prepared by adding 20 ml of lemon juice to 20 ml of methanolic extract filtrate of *Mimosa pudica* L and *Azadirachta indica* A leaves. The remaining ingredients include all the same as mentioned above in formulation 1. The formulations of both F1 volume is made up to 100ml with purified water.

#### Formulation 2(f-2)

This formulation was prepared by adding 20 ml of lemon juice to 20 ml of methanolic extract filtrate of *Mimosa pudica* L and *Azadirachta indica* A leaves.

The remaining ingredients include all the same as mentioned above in formulation 1

The formulations of both F1 and F2 show in below table.

**Table 1:** Formulation of Polyherbal Hand Wash

Ingredients	Quality
Methanolic extract of <i>mimosa pudica</i> L and <i>azadirachta indica</i>	20ml
Purified water	100ml
Sodium lauryl sulphate	6gms
Glycerine	40ml
Methyl paraben	0.3gms
Rosemerry oil	5ml

**Table 2:** Formulation of Poly Herbal Hand Wash F-2

Ingredient	Quantity
<i>Rosemerry oil</i>	5 ml
<i>Methanolic extract of mimosa pudica L and azadirachta indica</i>	20ml
<i>lemon water</i>	20ml
<i>sodium lauryl sulphate</i>	6gms
<i>Glycerine</i>	40ml
<i>methyl paraben</i>	0.3gms
<i>Purified water qs</i>	100ml

**Evaluation of Antimicrobial Activity**

The screening of anti-microbial efficacy of the formulated poly herbal hand wash was performed on various microorganisms by using agar plate method as per standard procedure. Four sterile petri plates were taken for testing the anti-microbial activity against four different microorganisms, Staphylococcus aureus. The plates were filled with nutrient agar solution and allowed for solidification. After solidification the microorganisms from the subculture were inoculated into the nutrient agar media and three cavities were made in it. The first cavity is filled with standard antibiotic amoxicillin, second one with herbal hand wash without lime water (F-1) and third cavity is filled with herbal hand wash with lime water (F-2). It was taken care that sample should be placed at the level of cavity. The plates are placed in incubator at 37°C to test the activity. After 24 hours the plates were observed for the formation of zone of inhibition. From the zone of inhibition the anti-microbial activity of formulation is estimated .


**III. RESULT AND DISCUSSION**

The prepared formulation of polyherbal hand wash were subjected for physical evaluation and antimicrobial efficacy.

- **Appearances:** The prepared two for formulation of hand wash appear as greenish brown and greenish yellow colour.
- **PH:** The ph of formulation was measured by digital ph meter. The ph of two formulation was found to be 6.31 and 6.24
- **Viscosity:** The viscosity of hand wash was determined by using Brookfield viscometer. 50ml of herbal hand wash is taken into 100ml of beaker and the tip of viscometer was dipped into the beaker containing hand wash formulation and its viscosity was measured. The viscosity of F-1 and F-2 was found.
- **Antimicrobial Activity:** The Anti-microbial efficacy of the formulations of Polyherbal Hand Wash was tested on Staphylococcus aureus by agar plate technique. The results of zone of inhibition showed that the hand wash prepared from methanol extract of the combined plant materials shown significant antimicrobial activity. The hand wash prepared with lemon juice (F-2) showed little higher activity than the formulation prepared without lemon juice (F-1)

**Antimicrobial Activities of Polyherbal Hand Wash Formulation**

organism	Zone of inhibition in Cms		
	Std. drug	F-1	F-2
Bacillus subtilus	1.2	3.4	3.6
Staphylococcus aureus	2.9	3.4	4.2
Pseudomonas aeruginosa	2.7	3.8	4.4
Escherichia coli	1.9	3.2	3.8



Plates showing zone of inhibition for different organism

#### IV. CONCLUSION

The results suggest that methanolic extract of *Azadiracta indica*, *Mimosa pudica* and their combinations with lemon water are capable of giving superior zone of inhibition to protect against the skin pathogens. This might be a rational basis for use of herbs in preparation of hand wash and use of these compounds in making antiseptic lotions or soaps in place of chemicals.

#### REFERENCES

- [1]. Shweta SP, Yuvraj JM, Shrinivas KM. Formulation and evaluation of herbal hand wash. Int J U Pharm and Bio Sciences. 4(2); 2015:30-33.
- [2]. Jyothi MJ, AV Praveen Kumar AV, Mohanalakshmi S, Prathyusha S. Formulation and evaluation of poly herbal hand wash. Int J Pharm. 2(2); 2012: 39-43. 6.
- [3]. Minakshi GJ, Kamat DV, Kamat SD. Evaluation of herbal handwash formulation. Natural Product Jyothi MJ, AV Praveen Kumar AV, Mohanalakshmi S, Prathyusha S. Formulation and evaluation of poly herbal hand wash. Int J Pharm. 2(2); 2012: 39-43. 6.
- [4]. Ali HA, Shehab NG, Rasool BK. Formulation and evaluation of herbal hand wash from Matricaria chamomilla flower extracts. Int J Res Ayu and Pharm. 2(6); 2011:1811-1813 Radiance. 7(5); 2008:413-415.
- [5]. Mashood AH, Satheesh B, Natarajan, Gousuddin M. Formulation, Evaluation and Antibacterial Efficiency of Herbal Hand Wash Gel. Int J Pharm Sci Rev Res. 25(2); 2014: 120-124
- [6]. Tamilarasi T and Ananthi T. Phytochemical Analysis and Anti Microbial Activity of Mimosa pudica Linn. Res. J. Chem. Sci. 2(2); 2012: 72-74.
- [7]. Jyothi MJ, AV Praveen Kumar AV, Mohanalakshmi S, Prathyusha S. Formulation and evaluation of polyherbal hand wash. Int J Pharm. 2(2); 2012: 39-43