

Active Pharmaceutical Ingredient and Impurity Profiling Paracetamol

D. A. Patil, S. K. Bais, Makardhwaj Bansode

Fabtech College of Pharmacy, Sangola, Solapur, Maharashtra, India

Abstract: *The first paracetamol was created in 1877, or even 1852. In both the US and Europe, it is the most frequently prescribed drug for pain and fever. It is listed as one of the Essential Medicines by the World Health Organization. Tylenol and Panadol are just a couple of the brand names for the generic drug paracetamol, which is also sold under numerous names. It received more than 5 million prescriptions in 2020, ranking it as the 118th most popular drug in the country. Acetaminophen, usually referred to as paracetamol, is a drug used to treat fever and mild to moderate discomfort. Tylenol and Panadol are examples of popular brand names. The benefits of paracetamol use for fever are unclear because, at a typical dose, it only modestly lowers body temperature; in that regard, it is inferior than ibuprofen. Acute mild migraines may be helped by paracetamol, however recurring tension headaches may only be minimally relieved. Ibuprofen is superior to paracetamol in terms of effectiveness for post-surgical pain management. The medicine of choice for lowering fever is paracetamol. It's unclear whether paracetamol has any advantages in musculoskeletal problems like osteoarthritis and backaches. When used to treat mild to moderate pain, paracetamol can help with headaches, muscle pains, minor arthritic pain, toothaches, and discomfort from the common cold, the flu, sprains, and dysmenorrhea. It is particularly advised for acute mild to moderate pain because there is little evidence to support its use in the treatment of chronic pain. At a dose of 4gram per day, paracetamol may marginally elevate systolic blood pressure in hypertensive patients. Although it's uncertain whether paracetamol is the real reason of this increase, studies show that children of mothers who used paracetamol for an extended period of time during pregnancy had higher rates of asthma, developmental, and reproductive issues. A maximum of three to four grammes per day is advised for adults. Toxicity from higher doses, including liver failure, is a possibility. Most drug overdoses in the US, UK, Australia, and New Zealand are caused by paracetamol poisoning, which is the leading cause of acute liver failure in the Western world.*

Keywords: Paracetamol

Objective:

1. To synthesise paracetamol from p-amino phenol, an antipyretic drug
2. Describing paracetamol
3. To use chromatography to evaluate the impurity profile

I. INTRODUCTION

Module 1:

Laboratory Safety:

Every time a worker is in a laboratory, basic safety regulations for behaviour should be followed. Below is a collection of many of the most prevalent safety guidelines.

- You are personally responsible for your safety.
- Consistently adhere to the right methods.

Never cut corners.

- If you created a mess, accept responsibility and clean it up.
- Keep your workspace tidy and organised
- Ensure that access to emergency exits and equipment is unobstructed and simple.
- While working, stay awake and alert.

- Be aware of the locations of eyewash stations, fire extinguishers, and lab safety showers.
- Be familiar with emergency exits.
- Prevent all chemical contact with the skin and yuca.
- Limit your exposure to all chemicals.
- Horseplay will not be permitted.

Assume that all compounds are very dangerous even if their toxicity is upermitte

- Be sure to protect your eyes.
- Use Fire Safety Procedures
- Do safely handle glassware.
- Retain notes.
- Be Sure to Glow Safely.

Don closed-toed footwear.

Don't forget to use electrical Safety

Introduction to Hazardous Chemicals and MSD

In Australian workplaces, chemical risks are a significant issue for occupational health and Safety. A variety of specialists must work together to manage chemical dangers. The need of collaborating with OHS experts to guarantee the use of a variety of talents is emphasise they are frequently connected to Major Hazard Facilities (MHF), chemical dangers are under the purview of all OHS practitioners. The management of bulk chemical or explosive transport, storage, and handling is typically outside the purview of the generalist OHS professional. In Australian workplaces, chemical risks are a significant issue for occupational health and safety. A variety of specialists must work together to manage chemical dangers. The need of collaborating with OHS experts to guarantee the use of a variety of talents is emphasised. Although they are frequently connected to Major Hazard Facilities (MHF), chemical dangers are under the purview of all OHS practitioners. The management of bulk chemical or explosive transport, storage, and handling is typically outside the purview of the generalist OHS professional.

Definition

Any element, chemical compound, or combination of elements and/or compounds is referred to as a chemical. The operational definition of hazardous substances is "substances that, upon worker exposure, may have a negative impact on health" (Safe Work Australia, 2010b). the word Emergency response organisations use the term "hazardous material" to describe substances that require special handling.

MSDS

A material safety data sheet is a technical document that offers thorough information about a controlled product, including information on the product's potential health consequences. Review of the risks associated with the handling, storage, or usage of the product. Protective measures for exposed workers. Emergency action

Introduction to Handling of Chemical and Safety Requirements

Chemicals kept in lab vials ought to be labelled with the day, month, and year they were initially received and opened. Only the substances that will be actively used should be stored, at the most. For work with particular chemical dangers, adhere to the following chemical storage recommendations. The programme seeks to give laboratory personnel the instruction, knowledge, and assistance they need to perform their jobs safely at home and in the lab. Large bottles of acids should be kept on low shelves, on trays, or in cabinets designated "corrosives" or "acids." Separate organic acids, flammable, and combustible compounds from oxidising acids. To move acid bottles, use a trolley or bottle carriers. Keep a list of all the substances you have on hand at work. Identify safe chemical usage, such as handling guidelines, protective gear requirements, mixing guidelines, and washing guidelines in the event of spills or skin contact. Eat, drink, and smoke should not be done near Chemicals. There are many different physical and health concerns associated with toxic chemicals and substances. Keep a list of all the substances that are kept on hand at work. Request safety data sheets from your chemical supplier and keep them on hand for reference. Eat, drink, and smoke should not be done near chemicals

Safety Requirements

- Helmets
- Goggles
- Glovees
- Masks
- Safety goggles
- Guards.
- Shields
- A lab coat
- First-aid supply
- Extinguishers for fire
- A chemical hood

II. MODULE 2: LABORATORY TECHNIQUES

Glass, the material used to construct laboratory equipment, is resistant to the majority of chemicals. Typically, two types of glass are utilised to create laboratory equipment. These are both very combustible types of glass: soda-lime glass and borosilicate glass. Soda-lime glass should be heated and cooled gently because abrupt heating and chilling might cause it to shatter. Heating soda, limestone, and silica at a temperature of 300–400 °C produces soda-lime. Breakage is avoided by annealing followed by consistent cooling and a gentle reheat. Borosilicate glass requires an oxygen natural gas flame to work because it does not soften below 700–800°C. The natural gas flame is produced by burning natural gas combined with oxygen. This glass has a low coefficient of expansion, making equipment manufactured with it resistant to abrupt temperature fluctuations

General Techniques

The book performs a fair job of covering an A to Z of laboratory procedures despite its 1964 copyright. The majority of the more modern methods mentioned are probably irrelevant to the work of the typical high school teacher (e.g., HPLC, TLC, mass spectrometry, etc.). The chapters on the application and maintenance of analytical balances and volume. Glassware are excellently written. It also contains details on calibration procedures and National Bureau of Standards tolerances. Although not all of the topics will be covered by the normal High school instructor, they are all well written and easy to understand. This publication is “intended for analytical chemists, laboratory technicians, beginners, and students,” according to the author. The book’s age subtracts from any potential advantages. A glaring omission in the distillation portion is the absence of glassware with conventional taper-r grindings.

Chromatography

In both synthetic and biological chemistry, molecules are created by a series of reactions involving intermediates. Although the molecules (benzene, phenol, and aniline) are similar to one another, they have different physical and chemical characteristics that can be utilised to distinguish them from one another.

Physical Properties:-


1. Molecular weight
2. Boiling point (in case both are liquid, as in this case)
3. Freezing point
4. Crystallization
5. Solubility
6. Density

Chemical Properties

1. Functional Group, for example, phenol has –OH where as aniline has NH₂
2. Reactivity towards other reagent to form complex.

Extraction Technique


- Extraction, as the term is used pharmaceutically, involves the separation of medicinally. Active portions of plant or animal tissue from the inactive or inert components. Products so obtained
- from plants are relatively. Impure liquids, semisolids or powders intended only for oral or external use.
- One or more components in the mixture are not thermally stable.
- The components in the mixture have a very high or low boiling point requiring vacuum or Cryogenic distillation, which is very energy intensive.
- The boiling points of the components are very close or they form azeotropes • Two
- Components with very different boiling points have to be separated at the same time.
- The components to be separated (pollutants or valuable products) are only a small fraction Of the mixture.



Crystallization:

A common separation technique used in the chemical and pharmaceutical industries is crystallisation. Many crystallization-related phenomena are still not fully understood. Research on crystallisation is becoming increasingly popular as consumers demand products with consistent quality (purity, crystal size, etc.). The process of creating a crystalline substance from a liquid, gas, or amorphous substance is known as crystallisation. The crystal lattice serves as the foundation for the very regular interior structure of crystals. In the process industry, crystallisation is crucial since it produces a solid product while also purifying it. The most promising crystals are translucent, have sharp edges, and are ideally between 0.1 and 0.4mm Size. Techniques used to grow crystals include solvent evaporation. Vapour diffusion, sublimation, solvent/non-solvent diffusion, and gradual cooling of the solution. As long as the sample is almost insoluble in the ether, the solvent combination $\text{CH}_2\text{Cl}_2/\text{EtOH}$ is an excellent choice. At the boundary where the solvents slowly diffuse into one another, crystals develop size.

CRYSTALLIZATION PROCESS




Distillation:

Distillation is the process of selective evaporation and condensation used to separate a component or components from a liquid mixture. The most fundamental technique for separating liquid mixtures and purifying liquids is distillation. Distillation entails heating a liquid to boiling, gathering the vapours, and condensing them back into the liquid state. Through this,

The separation of the mixture's liquids, which have boiling points that differ by a few degrees. Liquid separation from non-volatile components, • Cleaning up the liquid. A method of separating components from a liquid mixture is distillation. Processes are dependent on the liquid mixtures' vapour pressure properties. Packed columns have led to increased use in distillation, especially for low-volatility products such as paint and varnish.

Overview of distillation:-

A physical separation process called distillation uses the vaporisation of various constituents. It is employed in the refinery sector to convert crude oil into petroleum distillates. Since at least a few thousand years ago, alcohol has been concentrated through distillation, albeit in its most basic form. Vapors are created as the liquid heats up and rise to the distilling head, where a thermometer is frequently positioned. The liquid drips to the receiving flask, where the distillate is collected by the action of gravity.



Simple Distillation

- A technique for separating mixtures in a boiling liquid combination based on differences in their volatilities.
- Under normal pressure, it is possible to distill compounds with simple structures that are easily flammable and resistant to their own boiling heat. This method can be used to purify substances including hydrocarbons, alcohols, esters, small-molecule fatty acids, and amines.
- Liquids with various boiling points can be successfully separated using straightforward distillation. The component of the combination that boils at the lowest temperature will be the component with the richest vapours as the liquid being distilled is heated. Until every component of the initial mixture has been separated, the procedure can be repeated. After applying heat to a sample mixture, the components are instantly cooled by the action of cold water in a condenser.
- Instead of distilling the liquid combination as shown in the above picture, simple distillation entails heating it to the boiling point and then condensing it.

Fractional distillation:-

- By heating chemical mixtures to a temperature where one or more fractions of the mixture evaporate, chemical compounds are separated. Distillation is used to fractionate.
- Typically, the boiling temperatures of the constituent parts differ by no more than 25 °C (45 °F) from one another.

- Fractional distillation can be used to clean up combinations of liquids whose boiling points are relatively near to one another. The difference between the mixture's boiling points is often less than 80 °C, for example, acetone (b.p. 56 °C) and methyl alcohol (b.p. 65 °C).

Raoult's Rule :-

The vapour pressure of a solvent above a solution is equal to the vapour pressure of the pure solvent at the same temperature scaled by the mole fraction of the solvent present, according to Raoult's law:

$$P_{\text{solution}} = \chi_{\text{solvent}} P_{\text{solvent}}$$

P = atmospheric Pressure


X = number of moles

Vacuum Distillation

Vacuum distillation is the process of distilling under low pressure, which enables the purification of chemicals that are difficult to distil at ambient pressures or is done merely to save time and energy. Using this method, substances are separated based on the differences in their boiling points. Low pressure distillation or vacuum distillation. Organic substances that decompose at or below their boiling points are employed for it. Consider glycerol. The temperature at which the total vapour pressure equals the atmospheric pressure is known as a liquid's boiling point. This implies that the boiling point of the liquid can be decreased by lowering the pressure. We can get an organic substance to boil at a lower temperature if it breaks down at its boiling point. The substance boils at a considerably lower temperature and distils over undecomposed under reduced pressure. This is the purification procedure used to remove contaminants from non-volatile solid materials such as resin, stain, and others. Consider glycerol.

Steam Distillation

A separation technique called steam distillation involves distilling water along with other volatile and non-volatile ingredients. The separation technique of steam distillation is used to extract or purify components that are sensitive to temperature, such as natural aromatic compounds. A separation technique called steam distillation is employed to extract or purify components that are sensitive to temperature, such as natural aromatic compounds. The distillation apparatus is given steam or water to lower the compounds' boiling points. These substances ought to be water insoluble. The purification of organic molecules can benefit from steam distillation. The vapour from the distillation of organics is then condensed into a liquid. Due to the immiscibility of water and organics, the resultant liquid often consists of two phases: water and the organic substance. Organic distillate.



Rotary Evaporation

A rotary evaporator, commonly known as a "rotovap" or "rotovap," is a tool used in labs to gently and effectively evaporate solvents from samples. An illustration of a conventional rotary evaporator may be seen on the right. Rotating

an evaporator under steady heating and particular pressures can create a thin film (vacuum or atmospheric). Effective evaporation results in the vapours being separated, collected, and condensed. Application: Concentration vapour extraction Crystallization in particular materials that are temperature-sensitive The most efficient and significant tool in the biological, biochemical, pharmaceutical, chemical, food, and fragrance industries for research and development, educational purposes, and pilot testing is the rotary evaporator.

- Other Techniques:
- Boiling point:

The temperature at which a liquid's saturated vapour pressure equals the air pressure outside is known as the boiling point. When liquids are heated at constant pressure, like at atmospheric pressure, their vapour pressure, r , rises according to the amount of heat applied. The temperature at which a liquid's vapour pressure equals the pressure of the surrounding atmosphere is known as the boiling point of the liquid.

Melting point

The temperature at which a substance transforms from a solid to a liquid is known as its melting point. The solid and liquid phases are in equilibrium at the melting point. Pressure affects a substance's melting point, which is typically reported at a standard pressure such 1 atmosphere or 100 kPa. Wikipedia. The temperature at which a solid transforms from a solid to a liquid under atmospheric pressure is known as the melting point. Both the solid and liquid states coexist at the melting point because these two phases, solid and liquid, are in equilibrium.

Sublimation:-

When heated, a solid transforms immediately into a gas without first becoming liquid. Used to separate non-volatile contaminants such naphthalene, benzoic acid, anthracite, and camphor from volatile organic molecules. If the impurities present do not sublime, this process can be used to purify substances that do. When heated, many organic substances that are solids initially turn into vapour without ever becoming liquids. When the vapour is cooled, a solid is produced immediately. Sublimation is a method for separating a combination of two substances. Solid plus heat equals gas or vapour

(Heat = ΔH_{sub})

Chemical Test :-

The following characteristics are not generally included in chemical tests: PH, hardness, presence of a selected group of chemical metres, biocides, and highly poisonous symbols. The degree at which a liquid reaches its boiling point and becomes a gas. The pH is a unit of measurement and a gauge of the relative acidity or alkalinity of water. The answer is indicated by a PH 07 value. Alkalinity is indicated by neutron values of 2-1 and higher, while acidity is indicated by neutron values of 6-9. & lower.

Reaction Workup:-

The reaction work-up, which happens right away after the reaction is finished, is an important stage in this sequence. The work-up refers to a material purification procedure that typically takes place in a separatory funnel. Adding solutions to the funnel. Either to extract or combine with. In order to separate the finished product from excess reagent, catalysts, side products, solvers, or chemicals derived from side reactions.

III. MODULE 3

API Technology:


With customers ranging from big pharma to virtual companies, we have a long history of providing reliable service. Our manufacturing and development service. Routinely support for APIs.

Drugs that require advanced. My highly effective chemicals are created using knowledge in synthesis, which calls for specialised infrastructure and handling capabilities.

**Overview of Benzimidazole, Intermediate and Fine Chemical Industry:**

The drug paracetamol (PA) is frequently used. It is an analgesic that is used to treat minor discomfort and lower fever in individuals (antipyretic). Despite being used as an analgesic and an antipyretic, paracetamol is a crystalline solid with a melting point of 169–171 °C and no discernible anti-inflammatory properties. In the United States and many other nations, acetaminophen overdose is the most frequent cause of hepatotoxicity and abrupt liver failure. High dosages of paracetamol may cause undesired side effects in the body, but at recommended dosages, it has no adverse effects and is widely used because of its exceptional therapeutic properties and strict control over quality.


Acetylation of para amino phenol with acetic anhydride occurs in an aqueous medium to produce paracetamol. Only the amino group is acetylated in aqueous media out of the two groups present (amino and hydroxy). However, in aqueous media, the hydroxy group gets acetylated.

**PARACETAMOL****Unit Processes Synthesis Benzimidazole**

They exhibit a wide spectrum of biological and therapeutic uses and have been employed in the production of drugs and medicinal chemistry. For the industrial manufacturing (or thought of as such) of paracetamol from benzene, four alternative pathways were explored.

- Nitration of chlorobenzene is Route 1.
- Route 2: Phenol nitration
- Route 3: Nitrobenzene reduction.
- Hydroquinone is acetamidated in route 4.

By nitrating phenol with sodium nitrate, extracting the required p-nitro phenol from the ortho-byproduct, and reducing the nitro group with sodium borohydride, paracetamol can be made quickly and easily. After that, acetic anhydride is used to acetylate the resulting p-aminophenol. Since the phenol is a strong activator in this process, only a minor nitration condition is needed. Although the industrial method is similar, sodium borohydride reduction is replaced with hydrogenation.



Optimization of Organic reaction and process:

Increasing the efficiency of organic reactions can significantly improve synthetic chemistry by enabling quick, repeatable, and scalable development. Chemistry Numerous processes, such as condensation, heterocyclic creation, and catalyst-catalyzed cross coupling, have been studied in the context of microwaves.

Industrial process and scale up technique industrial manufacturing method and flow of charts:

1. The state of the pharmaceutical industry, including information on formulations, natural products, and bulk drugs India's position in relation to developed countries
2. Scale-up techniques, including techniques for process control, productivity maximisation, and process optimization
3. Case studies focusing on the logic behind the selection of routes, raw materials, process control techniques, pollution control procedures, etc. for selected drugs' chemical technologies
4. Data collection during pilot plant trials, preparation of flow diagrams, material balance sheets, and technical data sheets are all aspects of the chemical technology of some drugs.
5. Process technologies for a few natural products of interest to businesses.
6. For instance, 4-hydroxyisoleucine
7. Scale-up strategies for industrial pharmacy, typical SOPs for various dosage forms, and in-process control techniques
8. 7.manufacturing equipment: Tools used in the production of large-scale drugs.
9. Formulation-related equipment used in pharmaceutical manufacturing.

Polymorphism In Benzimidazole

The first paracetamol was created in 1877, or even 1852. In both the US and Europe, it is the most frequently prescribed drug for pain and fever. It is listed as one of the Essential Medicines by the World Health Organization. Tylenol and Panadol are just a couple of the brand names for the generic drug paracetamol, which is also sold under numerous names.

The biological effects of paracetamol include gastric ulcer, antihypertensive, bleeding propensity, pregnancy, and more. The efficiency of acetaminophen Paracetamol offers only limited pain relief for persons with knee or hip osteoarthritis, according to high-quality research. Its effectiveness in treating tension-type headaches is supported by evidence of fair quality.

Method and Materials

As it melts Open capillary tubes were used to determine the points of the organic compounds, and they were not adjusted. TLC, or thin layer chromatography, was used. On the manually made silica Gel G-coated glass slides (Rankem). DRS 8000 Spectrometer was used to record IR spectra in the IR Affinity-1 FTIR Shimadzu. Using Tetramethylsilane as the internal standard, ¹H NMR spectra were captured at 400 MHz using a Bruker Avance 400 spectrometer. The variation in thermal enthalpy Shimadzu's DSC-60 (Differential Scanning Calorimetry) was used to record the temperature, and the output was recorded as mW.

Chirality in API Industry, Resolution of racemate, asymmetric synthesis, few case studies:-

Louis Pasteur, a French scientist and biologist, made the discovery of chiral chemistry in 1848 when he manually separated the two isomers of sodium ammonium tartrate for the first time (1, 2). It took approximately a century, though, for researchers to discover that the phenomenon of chirality is crucial to the health of both plants and animals as well as to the pharmaceutical, agricultural, and other chemical sectors. The majority of alkaloids, hormones, nucleosides, proteins, enzymes, amino acids, carbohydrates, and carbohydrates are all chiral substances. In the pharmaceutical industry, 88% of the most recent medications are sold as racemates, which are composed of an equimolar mixture of two enantiomers, and 56% of the drugs now in use are chiral products (3-5). All natural compounds have a single enantiomeric form, as opposed to chiral artificial products. For instance, all natural amino acids are the l-isomer (levorotatory), and all natural sugars (carbohydrates) are the d-isomer (dextrorotatory). Despite

sharing the same chemical structure, the majority of enantiomers of racemic medications show distinct variances in pharmacology, toxicology, pharmacokinetics, metabolism, etc. Now it is clear how chiral medicines interact with the biological environment. In order to remove the undesirable isomer from the preparation, determine the best course of action for the patient, and establish the proper therapeutic monitoring, it is crucial to enhance chiral separation and analysis of racemic pharmaceuticals in both the pharmaceutical industry and the clinic.

Chirality

Chirality Defined

A molecule's capacity to exist in two asymmetric forms that are not superposable mirror images of one another without altering its atomic structure, atom-to-atom connections, or bond ordering is known as chirality. Fig. (1a) (1a). This phenomenon typically results in what might be called left-hand and right-hand copies of the same molecule due to a change in the three-dimensional orientation of four separate substituents connected to a single core atom.

Enantiomers are the names given to these two different forms of the molecule. There will always be at least one substituent attached to the chiral atom that cannot be superimposed when various versions are attempted to be combined. The Cahn-Ingold-Prelog system, or simply the R/S notation, is used to distinguish between the two enantiomers, as advised by the International Union of Pure and Applied Chemistry (IUPAC) [4] Fig (1b). The D / L nomenclature, which stands for dextrorotatory (clockwise) and levorotatory (counter-clockwise) optical rotation of polarised light, is only used for amino acids and sugars, two different sorts of molecules. As a result of the R/S notation for chirality and the +/- symbol for optical rotation [5], this practise is no longer widely used. In achiral contexts, enantiomers are typically said to have identical physical characteristics. With the rotation of plane polarised light being the one exception. In reality, chiral circularly polarised light has left- and right-handed components that make up plane polarised light. The phenomenon of optical rotation is caused by slight variations in how chiral molecules interact with these components. Achiral molecules are those that can be superimposed on their mirror images.




Fig. (1)

a) basic chirality.

These two molecules cannot completely superimpose despite sharing the identical atoms and atom-atom interactions. As a result, they are known as enantiomers and can be seen in three dimensions as mirror images. The main...

A racemate, also known as a racemic mixture, is a sample that contains both enantiomers of a substance in equal amounts. Enantiomer separation can be achieved via chiral chromatographic techniques as well as kinetic and thermodynamic resolution. One enantiomer may be preferred over the other by stereoselective chemical processes such as asymmetric synthesis (also known as chiral synthesis) or stereoselective enzymes.

Chirality can still exist as increasingly complicated compounds are taken into account, with more complex substituents connected to a single or several chiral centres. These molecules are referred to as diastereoisomers when they are not

enantiomers of one another. Similar to how varied physical characteristics like melting and boiling temperatures are possible for these molecules

To determine precedence and hence nomenclature when a possible chiral centre has two identical atoms linked to it, the following “level” of atoms must be taken into account. The carbon-oxygen substituent has higher precedence than the carbon-hydrogen substituent in identifying the R or S, for instance, if two carbons are connected to the chiral centre and one of them just has hydrogens attached while the other has an oxygen atom attached.

Paracetamol: brief Overview of QA/QC.GMP guidance in API Manufacturing (Q7 Q7A & Q11)

A quality assurance component that can be described is quality management since it is committed to fostering trust that quality requirements will be met. The focus of quality control is on ensuring that the requirements for quality are met by quality management.

The goal of the ICH Guideline 9.7 Good Manufacturing Practice Guideline for API is to provide guidance on GMP for the production of API under a suitable quality management system.


Additionally, it aims to guarantee that API uphold the standards of quality and purity that they stand for in processing.

Paracetamol:-

Acetaminophen, usually referred to as paracetamol, is a drug used to treat fever and mild to moderate discomfort. Tylenol and Panadol are examples of popular brand names.

Principal:-

The synthesis of the amide basically merely entails running the reaction at a specific temperature with the right catalyst. From p-aminophenol, paracetamol is made by acetylating it with acetic anhydride while using 3–4 drops of concentrated sulfuric acid as a catalyst.




Requirements:

Chemicals:- Acetic anhydride, p-amino phenol, Distilled water.

Glassware:- Conical flask, water bath etc

Mechanism



Procedure

1. Put 4.4 grammes of p-amino phenol in a conical flask with 12 millilitres of distilled water.
2. Stir in 4.8 ml of acetic anhydride.
3. Vigorously shake the reaction mixture and heat it on a water bath until an almost clear solution forms.
4. Use the ice bath to cool the conical flask.
5. Wash the product through a filter with cold water and then recrystallize it in hot water.
6. The product's melting point is 169°C.

Formula: P-amino phenol:- C₆H₇NO

P-acetyl amino phenol :- C₈H₉NO₂

Molar mass: 151.163g/mol

- Boiling point: 420°C
- Melting point: 169°

Uses of Paracetamol:-

1. Lower fever
2. lessen minor to moderate discomfort
3. Headaches
4. Minor headache
5. Low back discomfort
6. Gastritis or peptic ulcers
7. Propensity to bleed
8. Aspirin allergy
9. Viral contamination
10. Pregnancy
11. Osteoarthritis
12. Muscle ache.

IV. MODULE 4: EXPERIMENTAL TECHNIQUES

Purification and impurity control paracetamol:

By recrystallizing the substance from water, the material can be made pure. This method is based on the fact that paracetamol dissolves very well in hot water (5 g/100 cm³) but not very well in cold water (1.4 g/100 cm³). Insoluble contaminants can be filtered off when the crude solid is heated in water and dissolves. Impurities in paracetamol Acetaminophen, usually referred to as paracetamol, is a drug used to treat fever and mild to moderate discomfort. Tylenol and Panadol are examples of popular brand names.

Below is a list of the reference standards for paracetamol API, along with its pharmacopoeia, non-pharmacopoeia impurities, and stable isotopes.

Packaging and Repackaging of paracetamol:

The manufacturer's stability Guarantee is void if a medicine is taken out of its original packing and repackaged into a compliance aid. Despite this, stability research has only been done on a small number of medications. Aspirin, atenolol, clozapine, frusemide, paracetamol, prochlorperazine, and sodium valproate were repackaged into compliance aids. 8-14 However, no research has been done on the stability of these repackaged medications for a full year.

Calculation:

For Theoretical yield:-

P-amino phenol :-C₈H₉NO₂ P-acetyl amino phenol C₆H₇NO

109 gm/mol = 151gm/mol 4.4gm/mol = x

X = 151×4.4/108 X = 6.095gm

Theoretical yeild : 6.095gm

Practical yeild. : 3.008gm

$$\begin{aligned}\% \text{ Practical yeild} &= \text{Practical yeild} / \text{Theoretical yeild} \times 100 \\ &= 3.008 / 6.095 \times 100 \\ &= 49.35\%\end{aligned}$$

Result: The result synthesis of Benzimidazole was found to be as follows:

1. Theoretical yeild : 6.095gm
2. Practical yeild. : 3.008gm
3. % Practical yeild : 49.35%
4. Melting point. : 169

V. CONCLUSION

One of the most beneficial biological activities of paracetamol can be found. Numerous medical conditions, including gastric ulcers, pregnancy, and muscle tension, are treated with paracetamol. This review's presentation of effective and affordable processes for producing paracetamol by acetylating p-amino phenol with acetic anhydride in aqueous medium enables chemists to obtain first-hand knowledge of the process for producing paracetamol and Develop procedures for the mass manufacture of paracetamol, which will be highly helpful to chemists and professionals in this field. And this can expand over time to create novel, environmentally friendly, economically sound protocols for the large-scale manufacturing of significant pharmacophores based on synthetic paracetamol.

REFERENCES

- [1]. Rapid Chromatographic Techniques for Preparative Separation with Moderate Resolution." Still, W. C.; Kahn, M.; Mitra, A. J. Org. Chem. 1978, 43 (14), 2923-5.
- [2]. Preparative Reversed-Phase Flash Chromatography, a Convenient Method for the Workup of Reaction Mixtures." Kuhler, T. C.; Lindsten, G. R. J. Org. Chem. 1983, 48 (20), 3589-3591.
- [3]. Mullin, J.W., Gaska, C.: "The growth and dissolution of potassium sulphate crystals in a fluidizedBed crystallizer", Can. J. Chem. Eng. 47 (1969) 47, 483-489.
- [4]. Jones, A. G., Mullin, J.W.: "Programmed cooling crystallization of potassium sulphate solutions", Chem. Eng. Sci. 29 (1974) 1, 105-118.
- [5]. Crystals and Crystal Growing, Alan Holden and Phylis Singer, Anchor Books- Doubleday, New York, 1960
- [6]. The Growth of Single Crystals, R. A. Laudise, Solid State Physical Electronics Series, Nick Holonyak, Jr. Editor, Prentice-Hall, Inc., 1970
- [7]. Wankat, P. C.: "Separation Process Engineering", Chapman and Hall, London, 2006.
- [8]. Ghosal, S.K., Sanyal, S.K. and Dutta, S., Introduction to Chemical Engineering, Tata McGraw Hill Book Co. (2004).
- [9]. Gilliland, E. R., Multicomponent Rectification: estimation of number of theoretical Plates as a function of reflux ratio, Ind. Eng. Chem., 32, 1220-1223 (1940).
- [10]. Hines, A. L.; Maddox, R. N., Mass Transfer: Fundamentals and Applications, Prentice Hall; 1 Edition (1984).
- [11]. Molokanov, Y. K., Korabline, T. R., Mazuraina, N. I. And Nikiforov, G. A., An Approximate Method for Calculating the Basic Parameters of Multicomponent Fractionation, International Chemical Engineering, 12(2), 209 (1972).
- [12]. Vidarsson, Gestur; Dekkers, Gillian; Rispen, Theo (2014). "IgG subclasses and allotypes: from structure to effector functions". Frontiers in Immunology. 5: 520
- [13]. Finkelman, Fred D. (September 2007). "Anaphylaxis: Lessons from mouse models". Journal of Allergy and Clinical Immunology. 120 (3): 506–515.
- [14]. Whitten, Kenneth W.; Gailey, Kenneth D.; Davis, Raymond E. (1992). General chemistry (4th ed.). Saunders College Publishing. P. 475

- [16]. Boiani, M.; Gonzalez, M. Imidazole and Benzimidazole derivatives as chemotherapeutic agents. *Mini-Rev. Med. Chem.* 2005, 5, 409–424
- [17]. Narasimhan, B.; Sharma, D.; Kumar, P. Benzimidazole: A medically important heterocyclic moiety. *Med. Chem. Res.* 2012, 21, 269–283
- [18]. Bansal, Y.; Silakari, O. The therapeutic journey of benzimidazoles: A review. *Bioorg. Med. Chem.* 2012, 20, 6208–6236.
- [19]. Lidstrom, P.; Tierney J.; Wathey, B.; Westman, J. *Tetrahedron*. 2001, 57, 9225. (b) Perreux, L.; Loupy, A. J. *Tetrahedron*. 2001, 57, 9199.
- [20]. Liu, J. F.; Lee, J.; Dalton, A. M.; Bi, G.; Yu, L.; Baldino, C. M.; Brown, M. *Tetrahedron Lett.* 2005, 46, 1241. (b) Liu, J. F.; Yohannes, D. *Biorg. Med. Chem. Lett.* 2006, 16, 686.
- [21]. Patil A, Ganguly S, et al “A Systematic Review of Benzimidazole Derivatives as an Antiulcer Agent” *Rasayan, J Chem.*, 2008, 1(3):447-460
- [22]. JuttaKockler, Sherryl Robertson, Denise Hope, Alison Haywood, Beverley Glass
- [23]. JoaquínIsac-García, ... HenarMartínez-García, in *Experimental Organic Chemistry*, 2016 reference
- [24]. H. Jaeschke, M.L. Bajt, in *Comprehensive Toxicology*, 2010 reference 24. Basant Giri, in *Laboratory Methods in Microfluidics*, 2017 reference
- [25]. HaniehGhadimi, ... IS Amiri, in *Electrochemistry of Dihydroxybenzene Compounds*.
- [26]. <https://slideplayer.com/amp/15785699/>